



You have downloaded a document from  
**RE-BUŚ**  
repository of the University of Silesia in Katowice

**Title:** Induction of aromatic ring: Cleavage dioxygenases in *Stenotrophomonas maltophilia* strain KB2 in cometabolic systems

**Author:** Danuta Wojcieszńska, Urszula Guzik, Izabela Greń, Magdalena Perkosz, Katarzyna Hupert-Kocurek

**Citation style:** Wojcieszńska Danuta, Guzik Urszula, Greń Izabela, Perkosz Magdalena, Hupert-Kocurek Katarzyna. (2011). Induction of aromatic ring: Cleavage dioxygenases in *Stenotrophomonas maltophilia* strain KB2 in cometabolic systems. "World Journal of Microbiology and Biotechnology" (2011, no. 4, s. 805-811), DOI: 10.1007/s11274-010-0520-6



Uznanie autorstwa - Użycie niekomercyjne - Licencja ta pozwala na kopiowanie, zmienianie, remiksowanie, rozprowadzanie, przedstawienie i wykonywanie utworu jedynie w celach niekomercyjnych. Warunek ten nie obejmuje jednak utworów zależnych (mogą zostać objęte inną licencją).



UNIwersYTET ŚLĄSKI  
W KATOWICACH



Biblioteka  
Uniwersytetu Śląskiego



Ministerstwo Nauki  
i Szkolnictwa Wyższego

# Induction of aromatic ring: cleavage dioxygenases in *Stenotrophomonas maltophilia* strain KB2 in cometabolic systems

Danuta Wojcieszńska · Urszula Guzik ·  
Izabela Greń · Magdalena Perkosz ·  
Katarzyna Hupert-Kocurek

Received: 28 May 2010 / Accepted: 27 July 2010 / Published online: 10 August 2010  
© The Author(s) 2010. This article is published with open access at Springerlink.com

**Abstract** *Stenotrophomonas maltophilia* KB2 is known to produce different enzymes of dioxygenase family. The aim of our studies was to determine activity of these enzymes after induction by benzoic acids in cometabolic systems with nitrophenols. We have shown that under cometabolic conditions KB2 strain degraded 0.25–0.4 mM of nitrophenols after 14 days of incubation. Simultaneously degradation of 3 mM of growth substrate during 1–3 days was observed depending on substrate as well as cometabolite used. From cometabolic systems with nitrophenols as cometabolites and 3,4-dihydroxybenzoate as a growth substrate, dioxygenases with the highest activity of protocatechuate 3,4-dioxygenase were isolated. Activity of catechol 1,2- dioxygenase and protocatechuate 4,5-dioxygenase was not observed. Catechol 2,3-dioxygenase was active only in cultures with 4-nitrophenol. Ability of KB2 strain to induce and synthesize various dioxygenases depending on substrate present in medium makes this strain useful in bioremediation of sites contaminated with different aromatic compounds.

**Keywords** Nitrophenols · Cometabolism · Dioxygenase · Induction

## Introduction

Under aerobic conditions aromatic compounds, with the exception of those with deactivated substituent group,

undergo degradation through oxygenetic ring fission by catechol 1,2- or 2,3-dioxygenase as well as protocatechuate 3,4- or 4,5-dioxygenase. Presence of deactivated substituents with strong electrophilic properties makes the attack of oxygen on aromatic ring impossible (Pitter 1985). Since the nitro group is a powerful deactivating substituent aerobic degradation of these substrates occurs generally through the reduction as well as regrouping and elimination of the nitrate group following aromatic ring cleavage (Blasco et al. 1999; Schenzle et al. 1997). Transformation of polynitrated aromatic compounds by  $F_{420}$ -dependent enzymatic complex and hydride  $\delta$ -complex formation was also shown (Ebert et al. 1999). Some microorganisms degrade nitrophenols through aromatic ring fission by dioxygenases without prior removal of the nitrate group. However degradation of nitrophenols occurs most often through their hydroxylation to benzenetriol, which is the ring fission substrate for hydroquinone 1,2-dioxygenase (Chauhan et al. 2000; Pakala et al. 2007), Walia et al. (2002) have shown activity of nitrocatechol 2,3- dioxygenase catalyzing transformation of 4-nitrocatechol into 3-nitro-2-hydroxy-6-oxo-hexa-2,4-dienoic acid in *Pseudomonas putida* strain. In other study Kieboom et al. (2001) have observed induction of 2,3- dioxygenase in *Nocardia* strain S3 in the presence of 3-nitrocatechol. It has been shown that this enzyme exhibits absolute specificity to catechol C-3 substituents (Kieboom et al. 2001) and clearly suggests dependence between inductor and the activity of 2,3- dioxygenase. Cassidy et al. (1999) observed degradation of nitrophenols by *Sphingomonas* sp. strain UG30 after induction of hydroquinone dioxygenase by pentachlorophenol. Pentachlorophenol-induced enzyme transformed benzenetriol formed during *p*-nitrophenol degradation.

Because of high toxicity and resistance to biodegradation, mononitrophenols are insufficient source of carbon and energy for microorganisms and degradation of these

D. Wojcieszńska · U. Guzik (✉) · I. Greń · M. Perkosz ·  
K. Hupert-Kocurek  
Faculty of Biology and Environment Protection,  
Department of Biochemistry, University of Silesia,  
Jagiellonska 28, 40-032 Katowice, Poland  
e-mail: urszula.guzik@us.edu.pl

compounds requires presence of additional growth substrate, which increases biomass concentration and stimulates degradation of toxic chemicals. As the compounds similar in structure to transformed nitroaromatics accelerate their degradation, for mononitrophenols degradation other phenolic compounds are frequently used as growth substrate (Cho et al. 2000; Leung et al. 1997, 1999; Wan et al. 2007).

In previous works activity of various dioxygenases isolated from strain KB2 together with the ability of this strain to cometabolic transformation of nitroaromatics was observed (Greń et al. 2010; Guzik et al. 2009). Therefore in the present work an attempt to determine activity level of dioxygenases after their induction by different aromatic compounds was made. The main aim of this work was to check the activity of dioxygenases isolated from cultures with nitrophenols and verification how cometabolic conditions influenced synthesis of these enzymes.

## Methods

### Bacterial strain and growth conditions

Strain KB2 is a gram-negative, aromatic compounds—degrading bacterium that was isolated from activated sludge of a sewage treatment plant in Bytom—Miechowice in Poland as described previously (Guzik et al. 2009). On the basis of morphological and physiochemical characteristics and 16S rRNA gene sequence analysis was identified as *Stenotrophomonas maltophilia* (NCBI accession number DQ230920).

Liquid cultures of *Stenotrophomonas maltophilia* strain KB2 were grown in mineral salts medium (MSM) contained 3.78 g Na<sub>2</sub>HPO<sub>4</sub>·12H<sub>2</sub>O, 0.5 g KH<sub>2</sub>PO<sub>4</sub>, 5 g NH<sub>4</sub>Cl, 0.2 g MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.01 g yeast extract and 1 ml of trace element solution (TES) per liter. TES contained 3.82 g FeSO<sub>4</sub>·7H<sub>2</sub>O, 0.3 g CoSO<sub>4</sub>·7H<sub>2</sub>O, 0.08 g MnSO<sub>4</sub>·H<sub>2</sub>O, 0.14 g ZnSO<sub>4</sub>·7H<sub>2</sub>O, 0.006 g H<sub>3</sub>BO<sub>3</sub>, 0.04 NaMoO<sub>4</sub>·2H<sub>2</sub>O, 0.08 g NiSO<sub>4</sub>·7H<sub>2</sub>O, 0.003 g CuSO<sub>4</sub>·5H<sub>2</sub>O, 0.15 g Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>·18H<sub>2</sub>O, 0.006 Na<sub>2</sub>WO<sub>4</sub>·2H<sub>2</sub>O, 10 ml 32% HCl in 1 l of the solution. The pH was adjusted to 7.1.

In studies on cometabolic transformation of nitrophenols, as well as induction of enzymes, 3 mM benzoate (BA), 4-hydroxybenzoate (4-HB), 3,4-dihydroxybenzoate (3,4-DHB) as growth substrates and 2-nitrophenol (2-NP), 3-nitrophenol (3-NP) or 4-nitrophenol (4-NP) in concentration 1 mM, as cometabolites, were used. Stock solutions of benzoate (20 mM), 4-hydroxybenzoate (20 mM), 3,4-dihydroxybenzoate (20 mM), 2-nitrophenol (10 mM), 3-nitrophenol (10 mM) or 4-nitrophenol (10 mM) were prepared in distilled water.

Cells were proliferated in mineral medium with 3 mM of phenol (at 30°C on a rotary shaker at 130 rpm),

harvested by centrifugation (5,000×g at 4°C for 15 min) and washed with fresh sterile medium. Cells prepared in such a way were used as inoculum. Cultures in sterile mineral salt medium supplemented with 3 mM growth substrates and 1 mM of cometabolite were inoculated with previously prepared cells to the final optical density about 0.1 in absorbance scale at  $\lambda = 600$  nm, and incubated at 30°C with shaking at 130 rpm. Suitable growth substrate (benzoate, 4-hydroxybenzoate, 3,4-dihydroxybenzoate) was introduced into the culture every 72 h. All cultures were grown in triplicates.

### Bacterial growth determination

In cultures enriched with nitrophenols culture density was determined indirectly using modified method of Bradford (1976). 0.5 ml of culture was mixed with 0.5 ml KOH and incubated at 100°C for 10 min. After cooling, 0.2 ml of solution were taken, mixed with 1.0 ml of Bradford reagent and incubated for 10 min. Absorbance of sample was measured at 595 nm. Bovine serum albumin was used as a standard.

### Substrate determination

Culture supernatant was prepared by centrifugation (10,000×g at 4°C for 20 min) Concentration of aromatic compounds in the culture supernatant was determined by HPLC (Merck HITACHI) equipped with a LiChromospher<sup>®</sup> RP-18 column (4 × 250 mm) and a DAD detector (Merck HITACHI). The wavelength for detection of substrates, composition of eluent and solvent, as well as the flow rate, were developed separately for each aromatic compound. The mobile phase, in mononitrophenols and benzoic acid determination, was acetonitril and 1% acetic acid (40:60 v/v) at the flow rate of 1 ml min<sup>-1</sup>. 4-HB was separated in methanol:1% acetic acid (60:40 v/v) as a mobile phase, at the flow rate of 0.8 ml min<sup>-1</sup>. For separation of 3,4-DHB methanol:1% acetic acid (30:70 v/v) at the flow rate of 0.8 ml min<sup>-1</sup> was used. The detection wavelength was set at 260 nm. Chemical compounds in the supernatant were identified and quantified by comparing HPLC retention times and UV-visible spectra with those of external standards. Concentration of mononitrophenols as well as benzoic acids was expressed in mM and determined by peak area calculations. Data were corrected by subtraction of abiotic degradation of aromatic compounds.

### Preparation of cell extracts

Cells of *Stenotrophomonas maltophilia* strain KB2 were harvested after 7 days of culture by centrifugation (4,500

for 15 min at 4°C) and the pellet was washed with 50 mM phosphate buffer, pH 7.0, and resuspended in the same buffer. Cell-free extracts were prepared by sonication of the whole cell suspension (6 times for 15 s) and centrifugation at 9,000g for 30 min at 4°C. Clear supernatant was used as a crude cell extract for enzyme assays.

### Enzyme assays

Activity of catechol 1,2-dioxygenase [EC 1.13.11.1] was measured spectrophotometrically by formation of *cis,cis*-muconic acid at 260 nm ( $\lambda_{260} = 16,800 \text{ M}^{-1} \text{ cm}^{-1}$ ). The reaction mixture contained 20  $\mu\text{l}$  of catechol (50 mM), 67  $\mu\text{l}$   $\text{Na}_2\text{EDTA}$  (20 mM), 893  $\mu\text{l}$  of phosphoric buffer pH 7.4 (50 mM) and 20  $\mu\text{l}$  of crude enzyme extracts in a total volume of 1 ml. When activity of catechol 2,3-dioxygenase was detected, crude enzyme extract was incubated with 5%  $\text{H}_2\text{O}_2$  prior to determination of catechol 1,2-dioxygenase. In order to determine catechol 2,3-dioxygenase [EC 1.13.11.2] activity, formation of 2-hydroxymuconic semialdehyde was measured at 375 nm ( $\lambda_{375} = 36,000 \text{ M}^{-1} \text{ cm}^{-1}$ ). The reaction mixture contained 20  $\mu\text{l}$  of catechol (50 mM), 960  $\mu\text{l}$  of phosphoric buffer pH 7.4 (50 mM) and 20  $\mu\text{l}$  of crude extract in a total volume of 1 ml (Hegeman 1966). Activity of protocatechuate 3,4-dioxygenase [EC 1.13.11.3] was measured by protocatechuate depletion at 290 nm in a reaction mixture containing 20  $\mu\text{l}$  of protocatechuate (50 mM), 960  $\mu\text{l}$  of Tris-HCl buffer pH 7.4 (50 mM) and 20  $\mu\text{l}$  of crude extract in a total volume of 1 ml. A molar extinction coefficient of  $2,300 \text{ M}^{-1} \text{ cm}^{-1}$  was used, which is the difference between  $\lambda_{290}$  of protocatechuate ( $3,890 \text{ M}^{-1} \text{ cm}^{-1}$ ) and  $\lambda_{290}$  of the product 3-carboxy-*cis,cis*-muconate ( $1,590 \text{ M}^{-1} \text{ cm}^{-1}$ ). Protocatechuate 4,5-dioxygenase [EC 1.13.11.8] activity was measured spectrophotometrically by formation of 2-hydroxy-4-carboxymuconic semialdehyde at 410 nm ( $\lambda_{410} = 9,700 \text{ M}^{-1} \text{ cm}^{-1}$ ). The reaction mixture contained 20  $\mu\text{l}$  of protocatechuate (50 mM), 960  $\mu\text{l}$  of Tris-HCl buffer pH 7.4 (50 mM) and 20  $\mu\text{l}$  of crude extract in a total volume of 1 ml (Stanier and Ingraham 1954). In order to determine hydroxyquinol 1,2-dioxygenase [EC 1.13.11.37] activity, the formation of 4-hydroxymuconic semialdehyde was measured at 320 nm ( $\lambda_{320} = 11 \text{ mM}^{-1} \text{ cm}^{-1}$ ). The reaction mixture contained 200  $\mu\text{l}$  of hydroquinone (50 mM), 700  $\mu\text{l}$  of K-Na phosphate buffer pH 6.6 (100 mM) and 100  $\mu\text{l}$  of crude extract in a total volume of 1 ml (Zaborina et al. 1995). One unit of enzyme activity was defined as the amount of enzyme required to generate 1  $\mu\text{mol}$  of product per minute. Protein concentrations of the crude extract from different inducer-cultured bacteria were determined by the Bradford method using bovine serum albumin (Bradford 1976).

### Chemicals

2-nitrophenol, 3-nitrophenol, 4-nitrophenol were obtained from Merck AG, Darmstadt, Germany. 4-Hydroxybenzoic acid and protocatechuic acid were obtained from FLUKA, Fluka AG, Buchs, Switzerland. Benzoic acid and other chemicals were of analytical quality and were obtained from local suppliers.

## Results and discussion

### Cometabolic degradation of nitrophenols by *Stenotrophomonas maltophilia* KB2


Within aromatic substrates produced in plants, terpenoids and phenolic compounds, as benzoic acids, are commonly known. These compounds, as secondary metabolites, are produced on the determined growth stage by specific plant tissues or organs (Ornston and Stanier 1966; Singer et al. 2003). They are usually located in leaves by which they enter the soil or are secreted directly into the soil by the root system (Singer et al. 2003). The most important source of these metabolites in soil is decay of plants tissues. Aromatic compounds of plant origin are easily transformed and, as it is known, stimulate bacteria to xenobiotics degradation.

In our studies on mononitrophenols degradation 3 mM benzoate, 4-hydroxybenzoate, 3,4-dihydroxybenzoate as growth substrates were used. Because the ability of strain to metabolise high concentration of xenobiotic is desired quality, in our experiment 1 mM of nitrophenols was used despite the fact that degradation of these substrates at lower concentration (0.05–0.74 mM) was observed (Chauhan et al. 2000; Qui et al. 2007; Walia et al. 2002).

Figure 1 shows that depending on cometabolic system KB2 strain was able to degrade from 0.15 to 0.6 mM of mononitrophenol used after 14 days of cultivation. Moreover, presence of nitrophenols didn't inhibit degradation of carbon source. Increase of microbial biomass suggests resistance of KB2 strain to these substrates (Fig. 1). During 14 days of experiment only partial cometabolite degradation was observed. Nevertheless possible application of *Stenotrophomonas maltophilia* KB2 in natural environment could contribute to faster removal of nitroaromatics from degraded area as the bioremediation process takes a long time and the environment undergoes dynamic changes, accompanied by continuous influx of growth substrates.

### Activity of ring cleavage dioxygenases after induction by various aromatic substrates

KB2 strain is known to synthesize various types of dioxygenases, both catechol and protocatechuate, depending on



**Fig. 1** Cometabolic transformation of nitrophenols in the presence of aromatic compounds of plant origin (arrows indicate introduction of growth substrate into culture; BA-benzoic acid, 4-HB-4-

hydroxybenzoic acid, 3,4-DHB-3,4-dihydroxybenzoic acid, 2-NP-2-nitrophenol, 3-NP-3-nitrophenol, 4-NP-4-nitrophenol)

inductor used. In the previous work on degradation of different phenolic substrates by KB2, synthesis of catechol 1,2- and 2,3- dioxygenase was observed. Activity of catechol 2,3- dioxygenase significantly exceeded catechol 1,2- dioxygenase (Guzik et al. 2009). In the present work activity of catechol 2,3- dioxygenase (4.92 U/mg) was observed only after growing of strain in the presence of phenol, while aromatic hydroxyacids induced synthesis of protocatechuate 3,4- dioxygenase (Table 1). Interesting observation was the lack of protocatechuate dioxygenase activity after induction of KB2 by benzoate (Table 1) that may suggest degradation of this substrate *via* catechol and induction of additional enzyme—decarboxylase (Peng et al. 2003; Sparnins and Dagleys 1975). Isolation of enzymatic fraction after incubation of KB2 in the presence of mononitrophenols has shown no activity of catechol 2,3- dioxygenase but activity of catechol 1,2- dioxygenase (Table 1). According to the obtained results we assumed that catechol 2,3- dioxygenase was especially sensitive to nitroaromatics and lack of its' activity might force bacteria to synthesis of hydroquinone 1,2- dioxygenase, enzyme, which is most often induced by nitrophenols (Moonen et al.

2008; Pakala et al. 2007). However activity of hydroquinone 1,2- dioxygenase in the crude enzyme extract of KB2 strain was not observed (Table 1). Activity of protocatechuate 4,5- dioxygenase observed after culturing of KB2 in the presence of 4-NP (Table 1), suggests lower sensitivity of protocatechuate dioxygenases to inhibitory effect of mononitrophenols.

Presence of aromatic ring in the structure of natural phenolic compounds produced by plants (among them benzoic acids) induces synthesis of similar degradation enzymes as xenobiotics do. Such substrates usually are easily degraded by microorganisms according to the variety of degradation systems formed during evolution processes (Mendonca et al. 2004; Münzenberger et al. 2003; Sutherland et al. 1981). Because mononitrophenols were not degraded by KB2 in monosubstrate systems (Greń et al. 2010) we decided to introduce into cometabolic systems benzoic acid, 4-hydroxybenzoic acid and 3,4-dihydroxybenzoic acid. In the presence of 2-NP or 3-NP and 3,4-DHB or 4-HB as a growth substrate activity of protocatechuate 3,4- dioxygenase was observed (Table 2), while in the culture with 4-NP activity of this enzyme was observed


**Table 1** Dioxygenases activity after induction by aromatic substrates (4-HB-4-hydroxybenzoic acid; 3,4-DHB-3,4-dihydroxybenzoic acid; HQ-hydroquinone; 2-NP-2-nitrophenol; 3-NP-3-nitrophenol; 4-NP-4-nitrophenol; the plus/minus values represent standard deviation)

Enzyme	Catechol 1, 2-dioxygenase, U/mg protein	Catechol 2, 3-dioxygenase, U/mg protein	Protocatechuete 3, 4 -dioxygenase U/mg protein	Protocatechuete 4, 5 -dioxygenase U/mg protein	Hydroquinone 1, 2 -dioxygenase, U/mg protein
Inductor	Substrate in reaction mixture				
	Catechol	Catechol	3,4-DHB	3,4-DHB	HQ
Phenol	0.06 ± 0.02	4.92 ± 0.34	0.00 ± 0.00	0.00 ± 0.00	ND
4-HB	0.00 ± 0.00	0.00 ± 0.00	0.13 ± 0.05	0.00 ± 0.00	ND
3,4-DHB	0.00 ± 0.00	0.00 ± 0.00	0.46 ± 0.13	0.00 ± 0.00	ND
Benzoate	11.97 ± 0.34	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	ND
2-NP	0.29 ± 0.13	0.00 ± 0.00	0.00 ± 0.00	0.11 ± 0.00	0.00 ± 0.00
3-NP	0.12 ± 0.01	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00
4-NP	0.14 ± 0.08	0.01 ± 0.00	0.00 ± 0.00	0.73 ± 0.00	0.00 ± 0.00

**Table 2** Dioxygenases activity in cometabolic cultures (4-HB-4-hydroxybenzoic acid; 3,4-DHB-3,4-dihydroxybenzoic acid; 2-NP-2-nitrophenol; 3-NP-3-nitrophenol; 4-NP-4-nitrophenol; the plus/minus values represent standard deviation)

Enzyme		Catechol 1, 2 -dioxygenase, U/mg protein	Catechol 2, 3 -dioxygenase, U/mg protein	Protocatechuete 3, 4 -dioxygenase U/mg protein	Protocatechuete 4, 5 -dioxygenase U/mg protein
Inductor		Substrate in reaction mixture			
		Catechol	Catechol	3,4-DHB	3,4-DHB
2-NP	3,4-DHB	0.00 ± 0.00	0.06 ± 0.00	1.06 ± 0.12	0.00 ± 0.00
	4-HB	0.01 ± 0.00	0.00 ± 0.00	0.09 ± 0.00	0.00 ± 0.00
	Benzoate	0.01 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00
3-NP	3,4-DHB	0.00 ± 0.00	0.00 ± 0.00	0.26 ± 0.04	0.00 ± 0.00
	4-HB	0.00 ± 0.00	0.00 ± 0.00	0.81 ± 0.46	0.00 ± 0.00
	Benzoate	0.01 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00
4-NP	3,4-DHB	0.00 ± 0.00	0.23 ± 0.00	0.52 ± 0.12	0.00 ± 0.00
	4-HB	0.00 ± 0.00	0.00 ± 0.00	0.71 ± 0.65	0.00 ± 0.00
	Benzoate	0.00 ± 0.00	0.33 ± 0.00	0.00 ± 0.00	0.00 ± 0.00

only in the presence of 3,4-DHB (Table 2). Even though catechol 1,2- dioxygenase was not active and catechol 2,3- dioxygenase was active only in the presence of 4-NP and 3,4-DHB or benzoate (Table 2), partial degradation of nitrophenols was observed in all cometabolic systems (Fig. 1). As isolation and determination of enzymes activity showed synthesis of protocatechuete dioxygenases by KB2 strain in the presence of nitroaromatics (Table 1) it seems very interesting to determine if such enzymes might take part in nitrophenols degradation, even though degradation of these compounds *via* protocatechuete pathway was not observed. As it is known aerobic degradation of mononitrophenols proceeds through benzenetriol (Cassidy et al. 1999; Pakala et al. 2007), substrate which has very similar structure to protocatechuete, except the presence of hydroxyl group at C1, instead of carboxyl one (Fig. 2). Additionally it is known that attack of protocatechuete dioxygenases on aromatic ring take place between C3 and

**Fig. 2** Proposed region of attack of protocatechuete dioxygenases of KB2 strain (a the site of intradiol ring cleavage; b the site of extradiol ring cleavage)

C4 or C4 and C5 of catechol ring characteristic for both protocatechuete and benzenetriol (Vaillancourt et al. 2006). According to this knowledge we suggest that in cometabolite systems enzymes of protocatechuete acid degradation might be involved in degradation of benzenetriol formed in the consequence of mononitrophenols

degradation. However presented assumptions need to be confirmed by appropriate kinetic parameters determination, especially determination of inhibition constant for aromatic acid/mononitrophenol system as similarity of above-mentioned substrates suggest possibility of their competition for enzyme active site.

## Conclusion

Chemical diversity of contaminations introduced daily into natural environment is so huge that it is impossible to rule out the influence of simple as well as composite organic and inorganic compounds on dioxygenases activity. Because of that intensity of bioremediation processes proceed with the use of *Stenotrophomonas maltophilia* strain KB2 depends on the presence, availability and concentration of different aromatic substrates among them phenolic compounds of plant origin. Such substrates are not only very common in the environment but, as we have shown, induce different protocatechuate dioxygenases, enzymes that may be used in degradation of aromatic acids as well as other compounds with catechol structure. Although further research would be necessary to confirm our theory, wide substrate specificity of examined enzymes allows the use of *Stenotrophomonas maltophilia* strain KB2 for bioremediation of environments contaminated with different aromatic compounds.

**Acknowledgments** This work was funded by Polish Ministry of Science and Higher Education funded as a research project from grant-in-aid for scientific research in years 2007–2010. Authors would like to thank Agnieszka Materna for excellent technical assistance.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## References

- Blasco R, Moore E, Wray V, Pieper D, Timmis K, Castillo F (1999) 3-Nitroadipeate, a metabolic intermediate for mineralization of 2, 4-dinitrophenol by a new strain of a *Rhodococcus* species. *J Bacteriol* 181:149–152
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–258
- Cassidy MB, Lee H, Trevors JT, Zablutowicz RB (1999) Chlorophenol and nitrophenol metabolism by *Sphingomonas* sp UG30. *J Industr Microbiol Biotechnol* 23:232–241
- Chauhan A, Samanta SK, Jai K (2000) Degradation of 4-nitrocatechol by *Burkholderia cepacia*: a plasmid-encoded novel pathway. *J Appl Microbiol* 88:764–772
- Cho Y-G, Rhee S-K, Lee S-T (2000) Influence of phenol on biodegradation of *p*-nitrophenol by freely suspended and immobilized *Nocardioidea* sp. NSP41. *Biodegradation* 11:21–28
- Ebert S, Rieger PG, Knackmuss H-J (1999) Function of coenzyme F<sub>420</sub> in aerobic catabolism of 2, 4, 6-trinitrophenol and 2, 4-dinitrophenol by *Nocardioidea simplex* FJ2-1A. *J Bacteriol* 181:2669–2674
- Greń I, Wojcieszynska D, Guzik U, Perkosz M, Hupert-Kocurek K (2010) Enhanced biotransformation of mononitrophenols by *Stenotrophomonas maltophilia* KB2 in the presence of aromatic compounds of plant origin. *World J Microbiol Biotechnol* 26:289–295
- Guzik U, Greń I, Wojcieszynska D, Łabużek S (2009) Isolation and characterization of a novel strain of *Stenotrophomonas maltophilia* possessing various dioxygenases for monocyclic hydrocarbons degradation. *Braz J Microbiol* 40:285–291
- Hegeman GD (1966) Synthesis of enzymes of the mandelate pathways by *Pseudomonas putida*. Synthesis of enzyme by the wild type. *J Bacteriol* 91:1140–1154
- Kieboom J, van den Brink H, Frankena J, de Bont JAM (2001) Production of 3-nitrocatechol by oxygenase-containing bacteria: optimization of the nitrobenzene biotransformation by *Nocardia* S3. *Appl Microbiol Biotechnol* 55:290–295
- Leung KT, Tresse O, Errampalli D, Lee H, Trevors JT (1997) Mineralization of *p*-nitrophenol by pentachlorophenol-degrading *Sphingomonas* ssp. *FEMS Microbiol Lett* 155:107–114
- Leung KT, Campbel S, Gan Y, White DC (1999) The role of the *Sphingomonas* species UG30 pentachlorophenol-4-monooxygenase in *p*-nitrophenol degradation. *FEMS Microbiol Lett* 173:247–253
- Mendonca E, Martins A, Anselmo A (2004) Biodegradation of natural phenolic compounds as single and mixed substrates by *Fusarium flocciferum*. *J Biotechnol* 7:30–37
- Moonen MJH, Synowsky SA, van den Berg WAM, Westphal AH, Heck AJR, van den Heuvel RHH, Fraaije MW, van Berkel WJH (2008) Hydroquinone dioxygenase from *Pseudomonas fluorescens* ACB: a novel member of the family of nonheme(II)-dependent dioxygenases. *J Bacteriol* 190:5199–5209
- Münzenberger B, Hammer E, Wray V, Schauer F, Schmidt J, Strack D (2003) Detoxification of ferulic acid by ectomycorrhizal fungi. *Micorrhiza* 13:117–121
- Ornston LN, Stanier RY (1966) The conversion of catechol and protocatechuate and  $\beta$ -ketoadipeate by *Pseudomonas putida*. *J Biol Chem* 241:3776–3780
- Pakala SB, Gorla P, Pinjari AB, Krovidi RK, Baru R, Yanamandra M, Merrick M, Siddavattam D (2007) Biodegradation of methyl parathion and *p*-nitrophenol: evidence for the presence of a *p*-nitrophenol 2-hydroxylase in a gram-negative *Serratia* sp. strain DS001. *Appl Microbiol Biotechnol* 73:1452–1462
- Peng X, Misawa N, Harayama S (2003) Isolation and characterization of thermophilic *Bacilli* degrading cinnamic, 4-coumaric and ferulic acid. *Appl Environm Microbiol* 69:1417–1427
- Pitter P (1985) Correlation of microbial degradation rates with the chemical structure. *Acta Hydrochim Hydrobiol* 13:453–460
- Qui X, Zhong Q, Li M, Bai W, Li B (2007) Biodegradation of *p*-nitrophenol by methyl parathion-degrading *Ochrobactrum* sp. B2. *Int Biodet Biodegrad* 59:297–301
- Schenzle A, Lenke H, Fischer P, Williams PA, Knackmuss H-J (1997) Catabolism of 3-nitrophenol by *Ralstonia eutropha* JMP134. *Appl Environ Microbiol* 63:1421–1427
- Singer AC, Crowley DE, Thompson IP (2003) Secondary plant metabolites in phytoremediation and biotransformation. *Trends Biotechnol* 21:123–130
- Sparmins VL, Dagleys A (1975) Alternative routes of aromatic catabolism in *Pseudomonas acidovorans* and *Pseudomonas putida*: gallic acid as a substrate and inhibitor of dioxygenases. *J Bacteriol* 124:1374–1385
- Stanier RY, Ingraham JL (1954) Protocatechuic acid oxidase. *J Biol Chem* 210:799–808

- Sutherland JB, Crawford DL, Pometto AL (1981) Catabolism of substituted benzoic acids by *Streptomyces*. *Appl Environ Microbiol* 41:442–448
- Vaillancourt FH, Bolin JT, Eltis LD (2006) The ins and outs of ring-cleaving dioxygenases. *Crit Rev Biochem Mol Biol* 41:241–267
- Walia S, Shi L, Khan AA, Joshi B, Chaudry R (2002) Sequence analysis and molecular characterization of a nitrocatechol dioxygenase gene from *Pseudomonas putida*. *J Environ Sci Health B37*:379–391
- Wan N, Gu J-D, Yan Y (2007) Degradation of *p*-nitrophenol by *Achromobacter xylosoxidans* Ns isolated from wetland sediment. *Int Biodeter Biodegrad* 59:90–96
- Zaborina O, Latus M, Eberspacher J, Golovleva LA, Lingens F (1995) Purification and characterization of 6-chlorohydroxyquinol 1, 2-dioxygenase from *Streptomyces rochei* 303: comparison with an analogous enzyme from *Azotobacter* sp. Strain GP1. *J Bacteriol* 177:229–234