



You have downloaded a document from
RE-BUS
repository of the **University of Silesia in Katowice**


Title: Scalar 1-loop Feynman integrals as meromorphic functions in space-time dimension d

Author: Khiem Hong Phan, Tord Riemann

Citation style: Phan Khiem Hong, Riemann Tord.(2019). Scalar 1-loop Feynman integrals as meromorphic functions in space-time dimension d . "Physics Letters B" (Vol. 791, (2019), s. 257-264), doi 10.1016/j.physletb.2019.02.044



Uznanie autorstwa - Licencja ta pozwala na kopiowanie, zmienianie, rozprowadzanie, przedstawianie i wykonywanie utworu jedynie pod warunkiem oznaczenia autorstwa.



Scalar 1-loop Feynman integrals as meromorphic functions in space-time dimension d



Khiem Hong Phan^a, Tord Riemann^{b,c,*}

^a University of Science, Vietnam National University, Ho Chi Minh City, Vietnam

^b DESY Deutsches Elektronen-Synchrotron, 15738 Zeuthen, Germany

^c Institute of Physics, University of Silesia, 40-007 Katowice, Poland

ARTICLE INFO

Article history:

Received 27 January 2019

Received in revised form 21 February 2019

Accepted 26 February 2019

Available online 28 February 2019

Editor: A. Ringwald

Keywords:

Massive one-loop Feynman integrals

Generalized hypergeometric functions

Tensor integral reduction

ABSTRACT

The long-standing problem of representing the general massive one-loop Feynman integral as a meromorphic function of the space-time dimension d has been solved for the basis of scalar one- to four-point functions with indices one. In 2003 the solution of difference equations in the space-time dimension allowed to determine the necessary classes of special functions: self-energies need ordinary logarithms and Gauss hypergeometric functions ${}_2F_1$, vertices need additionally Kampé de Fériet-Appell functions F_1 , and box integrals also Lauricella-Saran functions F_S . In this study, alternative recursive Mellin-Barnes representations are used for the representation of n -point functions in terms of $(n-1)$ -point functions. The approach enabled the first derivation of explicit solutions for the Feynman integrals at arbitrary kinematics. In this article, we sketch our new representations for the general massive vertex and box Feynman integrals and derive a numerical approach for the necessary Appell functions F_1 and Saran functions F_S at arbitrary kinematical arguments.

© 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

We are studying scalar one-loop Feynman integrals,

$$J_n(d) = \int \frac{d^d k}{i\pi^{d/2}} \frac{1}{D_1^{v_1} D_2^{v_2} \dots D_n^{v_n}}, \quad (1)$$

with inverse propagators $D_i = (k + q_i)^2 - m_i^2 + i\epsilon$. We assume $v_i = 1$ as well as momentum conservation and all external momenta to be incoming, $\sum_{e=1}^n p_e = 0$. The q_i are loop momenta shifts and will be expressed for applications by the external momenta p_e . Dimensions $d = 4 + 2m - 2\epsilon$ with $m \geq 0$ are of physical interest because tensor one-loop Feynman integrals of rank r in $4 - 2\epsilon$ dimensions may be expressed by scalar integrals taken in higher dimensions up to $d = 4 + 2r - 2\epsilon$ [1]. Higher indices v_i will also appear in the reductions, but may be eliminated by integration by parts identities, so that a complete reduction basis of higher-dimensional scalar one- to four point integrals with indices one may be derived. One-loop integrals with variable indices are also needed in the context of the loop-by-loop Mellin-Barnes ap-

proach to multi-loop integrals of the Mathematica package AMBRE [3–6].

The first terms of the ϵ -expansion of one- to four-point scalar functions for $d = 4 - 2\epsilon$, until including the constant term, were given by G. 'tHooft and M. Veltman in 1978 [7]. A systematic numerical treatment of the next terms of order ϵ -terms was performed in 1992 [8], and a systematic numerical approach was worked out in 2001 [9]. It has been shown in 2003 [10,11] that representations in general dimension d , including $d = 4 - 2\epsilon$, will rely on certain multiple hypergeometric functions of the type ${}_2F_1, F_1, F_S$. Though, the explicit solutions for arbitrary kinematics could not be found.

A sketch of the Feynman integrals at arbitrary kinematics in terms of ${}_2F_1, F_1, F_S$ and their explicit numerical determination are the subject of this letter. The dependence on the external momenta p_e will be contained exclusively in the functions R_n :

$$R_n \equiv R_{12\dots n} = -\frac{\lambda_n}{G_n} - i\epsilon. \quad (2)$$

The R_n carry the causal regulator $-i\epsilon$. The Cayley matrix $\lambda_{12\dots n}$ was introduced in [12]. It is composed of the variables Y_{ij} , and its determinant λ_n is:

* Corresponding author at: DESY Deutsches Elektronen-Synchrotron, 15738 Zeuthen, Germany.

E-mail address: tord.riemann@desy.de (T. Riemann).

$$\lambda_n \equiv \det(\lambda_{12\dots n}) = \begin{vmatrix} Y_{11} & Y_{12} & \dots & Y_{1n} \\ Y_{12} & Y_{22} & \dots & Y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ Y_{1n} & Y_{2n} & \dots & Y_{nn} \end{vmatrix}, \quad (3)$$

with

$$Y_{ij} = Y_{ji} = m_i^2 + m_j^2 - (q_i - q_j)^2. \quad (4)$$

Further, we use the $(n - 1) \times (n - 1)$ dimensional Gram determinant G_n ,

$$G_n \equiv -2^n \det(G_{12\dots n}), \quad (5)$$

and

$$\det(G_{12\dots n}) = \begin{vmatrix} (q_1 - q_n)^2 & \dots & (q_1 - q_n)(q_{n-1} - q_n) \\ (q_1 - q_n)(q_2 - q_n) & \dots & (q_2 - q_n)(q_{n-1} - q_n) \\ \vdots & \ddots & \vdots \\ (q_1 - q_n)(q_{n-1} - q_n) & \dots & (q_{n-1} - q_n)^2 \end{vmatrix}. \quad (6)$$

We use the special assignment for tadpoles:

$$G_1 = -2. \quad (7)$$

Both determinants λ_n and G_n are independent of a common shift of the internal momenta q_i . Further, we introduce the notion $R(i)$,

$$R(i) \equiv r(i) - i\varepsilon \equiv -\det(\lambda_i)/G_1 - i\varepsilon = m_i^2 - i\varepsilon, \quad (8)$$

and use, wherever it is unique from the context,

$$R_1 \equiv R(i). \quad (9)$$

We derived in [13] a new ansatz, a recursion relation for the Feynman integrals defined in (1),

$$J_n(d) = \frac{-1}{2\pi i} \int_{c_0 - i\infty}^{c_0 + i\infty} ds \frac{\Gamma(-s)\Gamma(\frac{d-n+1}{2} + s)}{2\Gamma(\frac{d-n+1}{2})} \times \Gamma(s+1)R_n^{-s-1} \sum_{k=1}^n \partial_k R_n \mathbf{k}^- J_n(d+2s), \quad (10)$$

and its solution by a sequence of Mellin-Barnes representations. We use the representation $\partial_k R_n$ for the co-factor of the Cayley matrix, also called signed minors in e.g. [12]:

$$\partial_k R_n = \frac{\partial R_n}{\partial m_k^2} = \begin{pmatrix} 0 \\ k \end{pmatrix}_n. \quad (11)$$

The operator \mathbf{k}^- reduces an n -point Feynman integral $J_n(d)$ to $(n - 1)$ -point integrals $J_{n-1}(d)$ by shrinking the k^{th} propagator, $1/D_k$:

$$\mathbf{k}^- J_n(d) = \int \frac{d^d k}{i\pi^{d/2}} \frac{1}{\prod_{j \neq k, j=1}^n D_j}. \quad (12)$$

The recurrence relation (10) is the master integral for one-loop n -point functions in space-time dimension d , representing them by n integrals over $(n - 1)$ -point functions with a shifted, continuous dimension $d + 2s$. The recurrence starts at $n = 2$ with the tadpole $J_1(d)$ in the integrand:

$$J_1(d; m_i^2) = \int \frac{d^d k}{i\pi^{d/2}} \frac{1}{k^2 - m_i^2 + i\varepsilon} = -\frac{\Gamma(1 - d/2)}{(m_i^2 - i\varepsilon)^{1-d/2}} \equiv -\frac{\Gamma(1 - d/2)}{R_1^{1-d/2}}. \quad (13)$$

Eqn. (10) contains for $n = 2$ the term $\int ds (\frac{R_1}{R_2})^s$, multiplied by Γ -matrices with arguments depending on s , and is formally a Mellin-Barnes integral. Our representation is an alternative to Eq. (19) of [11]. There, an infinite sum over a discrete dimensional parameter s was derived in order to represent an n -point function $J_n(d)$ by integrals $J_{n-1}(d + 2s)$.

The further evaluations will depend, concerning the kinematics, exclusively on the R_1, R_2 , etc. introduced in (2). Although, there will arise exceptional cases when the specific choice of the external scalars $(p_{e_i} p_{e_j})$ or of internal mass squares m_i^2 will lead to vanishing or divergent determinants λ_n or G_n . In such cases, one has to go back to intermediate definitions and look for specific solutions.¹ See also the remarks in [14].

2. Massive vertex and box functions

Representations of the massive self-energy, vertex and box integrals can be derived iteratively from (10) by closing the integration contours of the Mellin-Barnes integrals e.g. to the right and taking the two series of residues of the corresponding Γ -functions with arguments $(-s + \dots)$. One Cauchy sum constitutes the analogue of the so-called boundary or b -terms of [11], the other one has a genuine d -dependence. Both sums together represent the Feynman integrals. In our approach, closed analytical expressions could be determined for arbitrary kinematics.

The general massive vertex and box integrals $J_3(d), J_4(d)$ have first been presented at the conference “Loops and Legs 2018 (LL2018)”. The vertex is

$$J_3(d) = J_{123} + J_{231} + J_{312}, \quad (14)$$

with short notations $R_3 = R_{123}, R_2 = R_{12}$ etc., and:

$$J_{123} = \Gamma\left(2 - \frac{d}{2}\right) \frac{\partial_3 R_3}{R_3} \frac{\partial_2 R_2}{R_2} \frac{R_2}{2\sqrt{1 - R_1/R_2}} \quad (15)$$

$$\left[-R_2^{\frac{d}{2}-2} \frac{\sqrt{\pi}}{2} \frac{\Gamma(\frac{d}{2}-1)}{\Gamma(\frac{d}{2}-\frac{1}{2})} {}_2F_1\left(\frac{d-2}{2}, 1; \frac{d-1}{2}; \frac{R_2}{R_3}\right) \right.$$

$$\left. + R_3^{\frac{d}{2}-2} {}_2F_1\left(1, 1; \frac{3}{2}; \frac{R_2}{R_3}\right) \right]$$

$$+ \Gamma\left(2 - \frac{d}{2}\right) \frac{\partial_3 R_3}{R_3} \frac{\partial_2 R_2}{R_2} \frac{R_1}{4\sqrt{1 - R_1/R_2}}$$

$$\left[+ \frac{2R_1^{\frac{d}{2}-2}}{d-2} F_1\left(\frac{d-2}{2}; 1, \frac{1}{2}; \frac{d}{2}; \frac{R_1}{R_3}, \frac{R_1}{R_2}\right) \right.$$

$$\left. - R_3^{\frac{d}{2}-2} F_1\left(1; 1, \frac{1}{2}; 2; \frac{R_1}{R_3}, \frac{R_1}{R_2}\right) \right]$$

$$+ (R_1(1) \leftrightarrow R_1(2)).$$

¹ A complete analysis of the exceptional kinematical cases has been performed by K.H.P.; to be published elsewhere.

We use the abbreviation (11). For $d \rightarrow 4$, both the sums of expressions with ${}_2F_1$ and F_1 in square brackets in (15) approach zero, thus compensating the pole factor $\Gamma(2 - d/2)$ in this limit. The J_3 stays finite at $d = 4$, as it should be for any massive 3-point function. And the ϵ expansion for J_{123} to order n needs, in this case, the evaluation of the components to order $(n + 1)$.

The corresponding massive four-point function is:

$$J_4(d) = J_{1234} + J_{2341} + J_{3412} + J_{4123}, \tag{16}$$

with $R_4 = R_{1234}, R_3 = R_{123}, R_2 = R_{12}$ etc., and:

$$\begin{aligned} J_{1234} = & \Gamma\left(2 - \frac{d}{2}\right) \frac{\partial_4 R_4}{R_4} \left\{ \right. \\ & \left[\frac{b_{123}}{2} \left(-R_3^{\frac{d}{2}-2} {}_2F_1\left(\frac{d-3}{2}, 1; \frac{d-2}{2}; \frac{R_2}{R_3}\right) \right. \right. \\ & \left. \left. + R_4^{\frac{d}{2}-2} \sqrt{\pi} \frac{\Gamma\left(\frac{d-2}{2}\right)}{\Gamma\left(\frac{d-3}{2}\right)} {}_2F_1\left(\frac{1}{2}, 1; 1; \frac{R_2}{R_3}\right) \right] \right\} \\ & + \frac{\Gamma\left(\frac{d-2}{2}\right)}{\Gamma\left(\frac{d-3}{2}\right)} \frac{\sqrt{\pi}}{4} \frac{\partial_3 R_3}{R_3} \frac{\partial_2 R_2}{\sqrt{1 - R_1/R_2}} \\ & \times {}_2F_1\left(\frac{1}{2}, 1; 1; \frac{R_2}{R_3}\right) \\ & \left[+ \frac{R_2^{\frac{d}{2}-2}}{d-3} F_1\left(\frac{d-3}{2}, 1, \frac{1}{2}; \frac{d-1}{2}; \frac{R_2}{R_4}, \frac{R_2}{R_3}\right) \right. \\ & \left. - R_4^{\frac{d}{2}-2} F_1\left(\frac{1}{2}, 1, \frac{1}{2}; \frac{3}{2}; \frac{R_2}{R_4}, \frac{R_2}{R_3}\right) \right] \\ & + \frac{R_1}{8} \frac{\Gamma\left(\frac{d-2}{2}\right)}{\Gamma\left(\frac{d-3}{2}\right)} \frac{\partial_3 R_3}{R_3} \frac{\partial_2 R_2}{R_2} \frac{1}{1 - R_1/R_3} \frac{1}{1 - R_1/R_2} \\ & \left[- R_1^{\frac{d}{2}-2} \frac{\Gamma\left(\frac{d-3}{2}\right)}{\Gamma\left(\frac{d}{2}\right)} \right. \\ & \left. \times F_5\left(\frac{d-3}{2}, 1, 1; 1, 1, \frac{1}{2}; \frac{d}{2}, \frac{d}{2}, \frac{d}{2}; \frac{R_1}{R_4}, \dots \right. \right. \\ & \left. \left. \frac{R_1}{R_1 - R_3}, \frac{R_1}{R_1 - R_2} \right) \right. \\ & \left. + R_4^{\frac{d}{2}-2} \sqrt{\pi} \right. \\ & \left. \times F_5\left(\frac{1}{2}, 1, 1; 1, 1, \frac{1}{2}; 2, 2, 2, \frac{R_1}{R_4}, \frac{R_1}{R_1 - R_3}, \frac{R_1}{R_1 - R_2}\right) \right. \\ & \left. + (R_1(1) \leftrightarrow R_1(2)) \right\} \\ & + (2, 3, 1) + (3, 1, 2), \tag{17} \end{aligned}$$

where the function b_{123} is independent of d ,

$$\begin{aligned} b_{123} = & \frac{1}{2} \frac{\partial_3 R_3}{R_3} \frac{\partial_2 R_2}{R_2} \left[\frac{R_2}{\sqrt{1 - \frac{R_1}{R_2}}} {}_2F_1\left(1, 1; \frac{3}{2}; \frac{R_2}{R_3}\right) \right. \\ & \left. - \frac{1}{2} \frac{R_1}{\sqrt{1 - \frac{R_1}{R_2}}} F_1\left(1; 1, \frac{1}{2}; 2; \frac{R_1}{R_3}, \frac{R_1}{R_2}\right) \right] + (1 \leftrightarrow 2). \tag{18} \end{aligned}$$

Here, it is $R_1 = R_1(1)$ and (11) defines derivatives like $\partial_2 r_2$. The term b_{123} , when multiplied with $\Gamma(-\frac{d-4}{2})R_3^{\frac{d}{2}-2}$, equals the term of J_{123} in (15) with d -independent F_1 and F_5 . It replaces the so-called b_3 -term of the vertex integral in [11] for arbitrary kinematics, while the d -dimensional parts of J_{1234} agree.

For $d \rightarrow 4$, all the expressions in square brackets in (17) approach zero, thus compensating the pole of $\Gamma(2 - d/2)$ in this limit. As a result, the J_4 stays finite at $d = 4$, as it should be for any massive 4-point function. And the ϵ expansion for J_{1234} to order n needs, in this case, the evaluation of the components to order $(n + 1)$.

The derivations of J_{123} and J_{1234} were done under the assumption that the kinematical arguments x, y, z of the ${}_2F_1, F_1, F_5$ fulfill $|x|, |y|, |z| < 1$. Nevertheless, the above formulae are valid at arbitrary kinematical arguments, for massive vertices at $\Re(d) > 2$ and for box integrals at $\Re(d) > 3$. In Appendix A to Appendix C we will show how to calculate the various F_1 and F_5 for arbitrary complex arguments; for ${}_2F_1$ we assume that such calculations are well-known.

3. Numerical results

The scalar one-loop basis consists of one- to four-point functions. Our two-point function $J_2(d)$ is in complete agreement with [11], while for $J_3(d)$ and $J_4(d)$ our results are novel. Concerning numerical results for the 3-point functions we refer to several tables in [13]. The kinematics was chosen such that the results of [11] could be compared.² Another numerical comparison, for a box integral $J_4(d)$ with vanishing Gram determinant, may be found in [15].

In Table 1 we show few examples of four-point functions in comparison to other packages. We did not aim at maximal accuracy and claim essentially six to eight safe digits (absolute values). Further, one propagator is massive and $d = 4$ or $d = 5$, and we can also allow for complex masses at the internal lines. A sample ϵ -expansion is reproduced for the generalized hypergeometric function F_1 in Table B.2.

For the safe numerical calculation of massive vertices J_3 and massive box integrals J_4 we collect stable numerical representations for the generalized hypergeometric functions F_1 and F_5 in the Appendices.

4. Discussion

The massive one-loop Feynman integrals have been represented as meromorphic functions of space-time d in terms of generalized hypergeometric functions. Many details left out here will be published elsewhere. The Feynman integrals can be calculated numerically at arbitrary kinematics and arbitrary dimension d , including potential pole locations at $d = 4 + 2m$. For phenomenological or multi-loop applications, it is wishful to have the pole expansions in closed analytical form. Their derivation is subject of a subsequent study.

The new recursion (10) has a unique feature. It allows to derive n -dimensional Mellin-Barnes integrals for n -point Feynman integrals. Generally, n -dimensional integrals are obtained by sector decomposition methods, while in the Mellin-Barnes approach, as it is advocated in numerical loop calculations, the number of dimension grows faster. Within the MBSuite, AMBRE generates for the most general massive n -point one-loop function an $N_n = \frac{1}{2}n(n + 1)$ -dimensional MB-integral; according to the number of entries

² We would like to thank Oleg Tarasov for a helpful discussion concerning this issue.

Table 1

Comparison of the box integral J_4 defined in (17) with the LoopTools function $\text{D0}(p_1^2, p_2^2, p_3^2, p_4^2, (p_1 + p_2)^2, (p_2 + p_3)^2, m_1^2, m_2^2, m_3^2, m_4^2)$ [16,17] at $m_2^2 = m_3^2 = m_4^2 = 0$. Further numerical references are the packages K.H.P_DO (PHK, unpublished) and MBOneLoop [15]. External invariants: $(p_1^2 = \pm 1, p_2^2 = \pm 5, p_3^2 = \pm 2, p_4^2 = \pm 7, s = \pm 20, t = \pm 1)$.

$(p_1^2, p_2^2, p_3^2, p_4^2, s, t)$	4-point integral
$(-, -, -, -, -, -)$	$d = 4, m_1^2 = 100$
J_4	0.00917867
LoopTools	0.0091786707
MBOneLoop	0.0091786707
$(+, +, +, +, +, +)$	$d = 4, m_1^2 = 100$
J_4	$-0.0115927 - 0.00040603 i$
LoopTools	$-0.0115917 - 0.00040602 i$
MBOneLoop	$-0.0115917369 - 0.0004060243 i$
$(-, -, -, -, -, -)$	$d = 5, m_1^2 = 100$
J_4	0.00926895
K.H.P_DO	0.00926888
MBOneLoop	0.0092689488
$(+, +, +, +, +, +)$	$d = 5, m_1^2 = 100$
J_4	$-0.00272889 + 0.0126488 i$
K.H.P_DO	(-)
MBOneLoop	$-0.0027284242 + 0.0126488134 i$
$(-, -, -, -, -, -)$	$d = 5, m_1^2 = 100 - 10 i$
J_4	$0.00920065 + 0.000782308 i$
K.H.P_DO	$0.0092006 + 0.000782301 i$
MBOneLoop	$0.0092006481 + 0.0007823090 i$
$(+, +, +, +, +, +)$	$d = 5, m_1^2 = 100 - 10 i$
J_4	$-0.00398725 + 0.012067 i$
K.H.P_DO	$-0.00398723 + 0.012069 i$
MBOneLoop	$-0.0039867702 + 0.0120670388 i$

Y_{ij} in the second Symanzik polynomial, $F(x) = \frac{1}{2}x_i Y_{ij} x_j - i\varepsilon$. For a vertex or box, $N_3 = 6, N_4 = 10$. In the present approach, it is only $N'_3 = 3, N'_4 = 4$. Evidently, a replacement of the original kinematical invariants $m_i^2, (p_{e,i} p_{e,j})$ or Y_{ij} by the alternatives $R_n = -\lambda_n / G_n$ is an essential building block and it might well be possible to find similar lower-dimensional MB-representations also for more involved multi-loop integrals.

Basic numerical features of the new n -dimensional MB-representation (10) have been studied in [15] in comparison with [2], with the package MBOneLoop, including cases of small or vanishing Gram determinant.

It is interesting to compare our results for $J_3(d)$ and $J_4(d)$ with the earlier study [11]. The d -dependent part of $J_3(d)$ as well as much of the d -dependent part of $J_4(d)$ agree with our results. Further, the expressions for the b -terms in [11] differ from our d -independent parts, although in certain kinematical regions they do agree numerically for $J_3(d)$. We find no agreement for $J_4(d)$, due to the various contributing b -terms.

Acknowledgements

T.R. would like to thank J. Fleischer, J. Gluza, M. Kalmykov and O. Tarasov for helpful discussions and J. Usovitsch for assistance in numerical comparisons. P.H.K.'s work is funded by the Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 103.01-2016.33. He would like to thank J. Blümlein and DESY for the opportunity to work in 2015 and 2016 as a guest scientist at Zeuthen. The work of T.R. is supported in part by a 2015 Alexander von Humboldt Honorary Research Scholarship of the Foundation for Polish Sciences (FNP) and by the Polish National Science Centre (NCN) under the Grant Agreement 2017/25/B/ST2/01987.

Appendix A. The Appell function F_1 and Lauricella-Saran function F_5

Numerical calculations of specific Gauss hypergeometric functions ${}_2F_1$, Appell functions F_1 (Eqn. (1) of [18]), and Lauricella-Saran functions F_5 (Eqn. (2.9) of [19]) are needed for the scalar one-loop Feynman integrals:

$${}_2F_1(a, b, c; x) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{k! (c)_k} x^k, \tag{A.1}$$

$$F_1(a; b, b'; c; y, z) = \sum_{m,n=0}^{\infty} \frac{(a)_{m+n} (b)_m (b')_n}{m! n! (c)_{m+n}} y^m z^n, \tag{A.2}$$

$$F_5(a_1, a_2, a_2; b_1, b_2, b_3; c, c, c; x, y, z) = \sum_{m,n,p=0}^{\infty} \frac{(a_1)_m (a_2)_{n+p} (b_1)_m (b_2)_n (b_3)_p}{m! n! p! (c)_{m+n+p}} x^m y^n z^p. \tag{A.3}$$

The $(a)_k$ is the Pochhammer symbol. The series converge for $|x|, |y|, |z| < 1$, but the functions are needed for arbitrary arguments. All the ${}_2F_1, F_1, F_5$ are finite and have no pole terms in ϵ . Practically all aspects of ${}_2F_1$ are well-known and implemented in computer algebra systems, in Mathematica as built-in symbol `Hypergeometric2F1[a, b, c, z]`. There is no public F_5 -package, while the Appell function $F_1(a; b_1, b_2; c; x, y)$ [18] is implemented in Mathematica as built-in symbol `AppellF1[a, b1, b2, c, x, y]` and in few other public packages. All the implementations have systematic limitations.

One approach to the numerics of F_1 and F_5 may be based on Mellin-Barnes representations. For the Gauss function ${}_2F_1$ and the Appell function F_1 , Mellin-Barnes representations are known. See Eqn. (1.6.1.6) in [20],

$${}_2F_1(a, b, c; z) = \frac{1}{2\pi i} \frac{\Gamma(c)}{\Gamma(a)\Gamma(b)} \times \int_{-i\infty}^{+i\infty} ds (-z)^s \frac{\Gamma(a+s)\Gamma(b+s)\Gamma(-s)}{\Gamma(c+s)}, \tag{A.4}$$

and Eqn. (10) in [18], which is a two-dimensional MB-integral:

$$F_1(a; b, b'; c; x, y) = \frac{1}{2\pi i} \frac{\Gamma(c)}{\Gamma(a)\Gamma(b')} \times \int_{-i\infty}^{+i\infty} dt (-y)^t {}_2F_1(a+t, b; c+t, x) \times \frac{\Gamma(a+t)\Gamma(b'+t)\Gamma(-t)}{\Gamma(c+t)}. \tag{A.5}$$

For the Lauricella-Saran function F_5 we derived the following, new, three-dimensional MB-integral:

$$F_5(a_1, a_2, a_2; b_1, b_2, b_3; c, c, c; x, y, z) = \frac{1}{2\pi i} \frac{\Gamma(c)}{\Gamma(a_1)\Gamma(b_1)} \int_{-i\infty}^{+i\infty} dt F_1(a_2; b_2, b_3; c+t, y, z) \times (-x)^t \frac{\Gamma(a_1+t)\Gamma(b_1+t)\Gamma(-t)}{\Gamma(c+t)}. \tag{A.6}$$

A general numerical evaluation of these representations deserves some sophistication. Let us mention the simple one-loop massive

QED vertex for which no trivial MB method exists when the kinematics is Minkowskian, a problem discussed e.g. in [21] and solved in [4]. It was demonstrated in [22] that MBOneLoop, a fork of the package MBnumerics [14,15,23,24] may be used to solve (A.4) to (A.6) at arbitrary kinematics with high precision.

One might also try to approach the generalized hypergeometric functions using Pochhammer's double loop contours [25,26], or study the defining differential equations [27–29], etc. After several trials, we decided to base our numerics on the integral representations of F_1 proposed in [30] and F_5 proposed in [19]; see Appendix B and Appendix C.

Appendix B. The Appell function F_1

A one-dimensional integral representation for F_1 [30] is quoted in Eqn. (9) of [18]:

$$F_1(a; b, b'; c; x, y) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \times \int_0^1 du \frac{u^{a-1}(1-u)^{c-a-1}}{(1-xu)^b(1-yu)^{b'}}. \tag{B.1}$$

We need three specific cases, taken at $d \geq 4$. Namely for vertices:

$$F_1^v(d) \equiv F_1\left(\frac{d-2}{2}; 1, \frac{1}{2}; \frac{d}{2}; x_c, y_c\right) = \frac{1}{2}(d-2) \int_0^1 \frac{du u^{\frac{d}{2}-2}}{(1-x_c u)\sqrt{1-y_c u}}. \tag{B.2}$$

Integrability is violated at $u = 0$ if not $\Re(d) > 2$. Similarly, for box integrals:

$$F_1^b(d) \equiv F_1\left(\frac{d-3}{2}; 1, \frac{1}{2}; \frac{d-1}{2}; x_c, y_c\right) = \frac{1}{2}(d-3) \int_0^1 \frac{du u^{d/2-5/2}}{(1-x_c u)\sqrt{1-y_c u}} = F_1^v(d-1). \tag{B.3}$$

Integrability is violated at $u = 0$ if not $\Re(d) > 3$. Finally for the definition of the box Saran function F_5 (C.1):

$$F_1^S(y_c, z_c) \equiv F_1\left(1; 1, \frac{3}{2}; \frac{3}{2}; y_c, z_c\right) = \frac{1}{2} \int_0^1 \frac{du}{\sqrt{1-u}(1-y_c u)\sqrt{1-z_c u}}. \tag{B.4}$$

The singularity at $u = 1$ is integrable.

Numerical checks may be performed using transformations of F_1 functions with different values of x, y [31], e.g.

$$F_1(a; b, b'; c; x, y) = (1-x)^{-b}(1-y)^{-b'} \times F_1\left(c-a; b, b'; c; \frac{x}{x-1}; \frac{y}{y-1}\right). \tag{B.5}$$

However, the relations do not allow to transform the real parts of both x, y to values smaller than one.

Appendix B.1. Specific values of ${}_2F_1$ and F_1 at $d = 4$

The vertex function (15) contains ${}_2F_1$ and F_1 with specific values at $d = 4$:

$${}_2F_1\left(1, 1; \frac{3}{2}; x_c\right) = \frac{\text{ArcSin}(\sqrt{x_c})}{\sqrt{1-x_c}\sqrt{x_c}} \tag{B.6}$$

and

$$F_1\left(1; 1, \frac{1}{2}; 2; x_c, y_c\right) = 2 \frac{\text{ArcTanh}\left[\frac{\sqrt{x_c}\sqrt{1-y_c}}{\sqrt{x_c-y_c}}\right]}{\sqrt{x_c}\sqrt{x_c-y_c}} - 2 \frac{\text{ArcTanh}\left[\frac{\sqrt{x_c}}{\sqrt{x_c-y_c}}\right]}{\sqrt{x_c}\sqrt{x_c-y_c}}. \tag{B.7}$$

Using logarithms only, $\text{ArcSin}(z) = -i \ln(iz + \sqrt{1-z^2})$ and $\text{ArcTanh}(z) = \frac{1}{2}[\ln(1+z) - \ln(1-z)]$. Eqn. (B.7) is only valid if $(x_c - y_c)$ has a well-defined imaginary part. For $x_c = x - i\epsilon_x$ and $y_c = y - i\epsilon_y$ this is not necessarily the case if ϵ_x and ϵ_y are independent and both infinitesimal. So (B.7) has to be used with a grain of care.

The box function (17) contains additional ${}_2F_1$ and F_1 with specific values at $d = 4$:

$${}_2F_1\left(\frac{1}{2}, 1; 1; x_c\right) = \frac{1}{\sqrt{1-x_c}} \tag{B.8}$$

and

$$F_1\left(\frac{1}{2}; 1, \frac{1}{2}; \frac{3}{2}; x_c, y_c\right) = \frac{1}{\sqrt{1-y_c}} {}_2F_1\left(\frac{1}{2}; 1, \frac{3}{2}; \frac{x_c-y_c}{1-y_c}\right) = \frac{\text{ArcTanh}\left(\sqrt{\frac{x_c-y_c}{1-y_c}}\right)}{\sqrt{x_c-y_c}}. \tag{B.9}$$

Eqn. (B.9) is only valid if $(x_c - y_c)$ has a well-defined imaginary part. Finally, we like to mention that we have no analogue to (B.8) and (B.9) for F_5 at $d = 4$, namely $F_5(\frac{1}{2}, 1, 1; 1, 1, \frac{1}{2}; 2, 2, 2; x_c, y_c, z_c)$.

The Appell function $F_1^S = F_1(1; 1, \frac{1}{2}; \frac{3}{2}; y_c, z_c)$ used in the integrand of the definition of the Saran function (C.1) can also be simplified:

$$F_1\left(1; 1, \frac{1}{2}; \frac{3}{2}; y_c, z_c\right) = \frac{1}{1-z_c} {}_2F_1\left(1, 1; \frac{3}{2}; \frac{y_c-z_c}{1-z_c}\right) = \frac{\text{ArcSin}\sqrt{\frac{y_c-z_c}{1-z_c}}}{\sqrt{(y_c-z_c)(1-z_c)}}. \tag{B.10}$$

Both representations in (B.10) are only valid when the imaginary part of the difference $(y_c - z_c)$ is well-defined.

For the Feynman integrals studied here, we have to take into account that x_c, y_c and z_c may have, in general, *uncorrelated infinitesimal* imaginary parts, and so their difference may be *not* well-defined. Let us remind that $x_c = R_1/R_4$, and $y_c = R_1/(R_1 - R_3)$, and $z_c = R_1/(R_1 - R_2)$. Here, all the R_n have, according to (2), identical imaginary parts $-i\epsilon$. This leads to different infinitesimal imaginary parts $-\epsilon_x, -\epsilon_y, -\epsilon_z$, with potentially different signs. So, one has basically two equivalent options. Either one treats ϵ_x, ϵ_y and ϵ_z as independent quantities and avoids the appearance of terms like $(x_c - y_c)$ and $(y_c - z_c)$. Or one uses the exact knowledge of the imaginary parts of the R_n from their definitions and arrives at well-defined imaginary parts of these $(x_c - y_c)$ and $(y_c - z_c)$.

Appendix B.2. Numerical calculation of $F_1^y(d)$

For $x_c = x - iX$ and $y_c = y - iY$, Eqn. (B.2) may be used for numerics if $(X, Y) \geq \text{const.} > 0$ or if $(x, y) < 1$. The remaining cases $(X = -\epsilon_x, Y = -\epsilon_y) \rightarrow +0$ deserve a closer inspection. They appear from Feynman integrals. We exemplify here the first one of the two more involved cases: $1 < x < y$ and $1 < y < x$ and introduce an auxiliary split parameter

$$u_m = \frac{1}{2} \left(\frac{1}{y} + \frac{1}{x} \right) \text{ with } 0 < \frac{1}{y} < u_m < \frac{1}{x} < 1. \tag{B.11}$$

In the integrand of F_1 there will a cut begin at $u = \frac{1}{y}$ and a pole arise at $u = \frac{1}{x}$ for infinitesimal ϵ_x, ϵ_y . A split of the integral at u_m ,

$$\int_0^1 du = \int_0^{u_m} du + \int_{u_m}^1 du \equiv i_L + i_R, \tag{B.12}$$

will lead to a separation of the singularities. In both integrals at the right hand side, the integrand is regular with one exclusion. We discuss now several opportunities of calculations, all of them with an accuracy of six to eight safe digits or better.

Our most careful approach pursued the following ansatz with additional splittings:

$$\begin{aligned} F_1^y(d) &= I_A + I_0 + I_C + I_D + I_B + I_E \\ &= \lim_{R \rightarrow +0} \left[\int_0^{\frac{1}{y}-R} + \int_{\frac{1}{y}+R}^{\frac{1}{y}+R} + \int_{\frac{1}{y}+R}^{u_m} + \int_{u_m}^{\frac{1}{x}-R} + \int_{\frac{1}{x}-R}^{\frac{1}{x}+R} + \int_{\frac{1}{x}+R}^1 \right] \end{aligned} \tag{B.13}$$

After performing the limit $R \rightarrow 0$ wherever possible, the integrals A and B will give real contributions, and the others are purely imaginary:

$$\begin{aligned} F_1^y(d) &= [\Re F_1^y(d)] + i [\Im F_1^y(d)] \\ &= [A + \text{sign}(\epsilon_x)\text{sign}(\epsilon_y) B] \\ &\quad + i [\text{sign}(\epsilon_y) (-C + D + E)]. \end{aligned} \tag{B.14}$$

It is

$$I_0 = 0, \tag{B.15}$$

$$A = \frac{d-2}{2} \int_0^{\frac{1}{y}} \frac{du u^{d/2-2}}{(1-xu)\sqrt{1-yu}}, \tag{B.16}$$

$$B = \frac{d-2}{2} \pi \frac{1}{x\sqrt{\frac{y}{x}-1} x^{d/2-2}}, \tag{B.17}$$

$$C = \frac{d-2}{2} \int_{\frac{1}{y}}^{u_m} \frac{du u^{d/2-2}}{(1-xu)\sqrt{yu-1}}, \tag{B.18}$$

$$\begin{aligned} D &= \frac{d-2}{2} \int_{u_m}^{\frac{1}{x}} \frac{du}{1-xu} \left(\frac{u^{d/2-2}}{\sqrt{yu-1}} - \frac{x^{-d/2+2}}{\sqrt{\frac{y}{x}-1}} \right) \\ &\quad + \frac{d-2}{2} \frac{1}{\sqrt{\frac{y}{x}-1} x^{d/2-2}} \left[\ln(R) - \ln\left(\frac{1}{2x} - \frac{1}{2y}\right) \right], \end{aligned} \tag{B.19}$$

$$\begin{aligned} E &= \frac{d-2}{2} \int_{\frac{1}{x}}^1 \frac{du}{1-xu} \left(\frac{u^{d/2-2}}{\sqrt{yu-1}} - \frac{x^{-d/2+2}}{\sqrt{\frac{y}{x}-1}} \right) \\ &\quad + \frac{d-2}{2} \frac{1}{\sqrt{\frac{y}{x}-1} x^{d/2-2}} \left[-\ln(R) + \ln\left(1 - \frac{1}{x}\right) \right]. \end{aligned} \tag{B.20}$$

The remaining R -dependences in (B.19) and (B.20) drop out in the sum of D and E .

Alternatively, with a subtraction in each of the two partial integrals in (B.12), one may regularize the integrand of $F_1^y(d)$ as follows:

$$i_L = \int_0^{u_m} du \frac{g_x(u) - g_x\left(\frac{1}{y}\right)}{\sqrt{1-yu}} + i_L^{ana}, \tag{B.21}$$

$$i_R = \int_{u_m}^1 du \frac{g_y(u) - g_y\left(\frac{1}{x}\right)}{1-xu} + i_R^{ana}, \tag{B.22}$$

with

$$\begin{aligned} i_L^{ana} &= -2 \frac{g_x\left(\frac{1}{y}\right)}{y_c} \left[\sqrt{1-y_c u_m} - 1 \right] \\ &\rightarrow -2 \frac{g_x\left(\frac{1}{y}\right)}{y_c} \left[-1 + i \text{sign}(\epsilon_y) \sqrt{y u_m - 1} \right], \end{aligned} \tag{B.23}$$

$$\begin{aligned} i_R^{ana} &= -\frac{g_y\left(\frac{1}{x}\right)}{x_c} \ln\left(\frac{1-x_c}{1-x_c u_m}\right) \\ &\rightarrow -\frac{g_y\left(\frac{1}{x}\right)}{x} \left[\ln\left(\frac{x-1}{1-xu_m}\right) + i\pi \text{sign}(\epsilon_x) \right]. \end{aligned} \tag{B.24}$$

Finally, a simplest approach will also do a reasonable numerics: Perform mean value integrals, like e.g. the built-in function of Mathematica:

$$F_1^y(d) = \lim_{\epsilon \rightarrow +0} \left[\left(\int_0^{\frac{1}{y}-\epsilon} + \int_{\frac{1}{y}+\epsilon}^{u_m} \right) + \left(\int_{u_m}^{\frac{1}{x}-\epsilon} + \int_{\frac{1}{x}-\epsilon}^1 \right) \right]. \tag{B.25}$$

Of course, a calculation with, say, more than six to eight safe digits, will deserve an explicit control of the algorithmic details.

Numerical examples for $F_1^y(d)$ are collected in Tables B.1 and (B.2).

Appendix B.3. Numerical calculation of the box Appell function $F_1^b(d)$

For the calculation of four-point Feynman integrals, one needs $F_1^b(d)$ as introduced in (B.3), both for $d = 4$ and for $d = 4 + 2m - 2\epsilon$. The box F_1 -function is related to the vertex function $F_1^y(d)$ by (B.3). Consequently, the numerics of the foregoing subsections may be taken over.

Appendix C. The Lauricella-Saran function F_S

For the calculation of the 4-point Feynman integrals, one needs the Lauricella-Saran function F_S [19]. Saran defines F_S as three-fold sum (A.3), see Eqn. (2.9) in [19]. He derives a 3-fold integral representation in Eqn. (2.15) and a 2-fold integral in Eqn. (2.16). We will use the following quite useful representation, derived at p. 304 of [19]:

Table B.1

The Appell function F_1 of the massive vertex integrals as defined in (B.2). As a proof of principle, only the constant term of the expansion in $d = 4 - 2\epsilon$ is shown, $F_1(1; 1, \frac{1}{2}; 2; x, y)$. Upper values: this calculation, (Appendix B.2), lower values: (B.7).

$x - i\epsilon_x$	$y - i\epsilon_y$	$F_1(1; 1, \frac{1}{2}; 2; x, y)$	
$+11.1 - 10^{-12} i$	$+12.1 - 10^{-12} i$	-0.1750442480735	$-0.0542281294732 i$
		$-0.17504424807351877884498289912$	$-0.054228129473304027882097641167 i$
$+11.1 - 10^{-12} i$	$+12.1 + 10^{-12} i$	$+1.7108545293244$	$+0.0542281294732 i$
		$+1.71085452932433557134838204175$	$+0.05422812947148217381589270924 i$
$+11.1 + 10^{-12} i$	$+12.1 - 10^{-12} i$	$+1.7108545304114$	$-0.0542281294732 i$
		$+1.71085452932433557134838204175$	$-0.05422812947148217381589270924 i$
$+11.1 + 10^{-12} i$	$+12.1 + 10^{-12} i$	-0.1750442480735	$+0.0542281294733 i$
		$-0.17504424807351877884498289912$	$+0.054228129473304027882097641167 i$
$+12.1 - 10^{-15} i$	$+11.1 - 10^{-15} i$	-0.1700827166484	$-0.0518684846037 i$
$+12.1 - 10^{-10} i$	$+11.1 - 10^{-15} i$	$-0.17008271664800058101165749279$	$-0.05186848460465674976556525621 i$
$+12.1 - 10^{-15} i$	$+11.1 + 10^{-15} i$	-0.1700827166484	$-1.7544202909955 i$
		$-0.17008271664844025647268817399$	$-1.75442029099557688735842562038 i$
$+12.1 + 10^{-15} i$	$+11.1 - 10^{-15} i$	-0.1700827166484	$+1.7544202909955 i$
		$-0.17008271664844025647268817399$	$+1.75442029099557688735842562038 i$
$+12.1 + 10^{-15} i$	$+11.1 + 10^{-15} i$	-0.1700827166484	$+0.0518684846037 i$
$+12.1 - 10^{-10} i$	$+11.1 - 10^{-15} i$	$-0.17008271664800058101165749279$	$+0.05186848460465674976556525621 i$
$+11.1 - 10^{-15} i$	-12.1	-0.0533705146518	$-0.1957692111557 i$
		$-0.05337051465189944473349401152$	$-0.195769211155733985388920833693 i$
$+11.1 + 10^{-15} i$	-12.1	-0.0533705146518	$+0.1957692111557 i$
		$-0.05337051465189944473349401152$	$+0.195769211155733985388920833693 i$
-11.1	$+12.1 - 10^{-12} i$	$+0.1060864084662$	$-0.1447440700082i$
		$+0.10608640847651064287133527599$	$-0.144744070021333407167349619088 i$
-11.1	$+12.1 + 10^{-12} i$	$+0.1060864084662$	$+0.1447440700082i$
		$+0.10608640847651064287133527599$	$+0.144744070021333407167349619088 i$
-12.1	-11.1	$+0.122456767687224028$	
		$+0.12245676768722402506513395161$	

Table B.2

The Appell function $F_1(1 - \epsilon; 1, \frac{1}{2}; 2 - \epsilon; x_c, y_c)$ as defined in (B.2), needed for $d = 4 - 2\epsilon$ at $x_c = 11.1 - 10^{-12} i$, $y_c = 12.1 - 10^{-12} i$.

$F_1(1 - \epsilon; 1, \frac{1}{2}; 2 - \epsilon; x_c, y_c)$	
$+(-0.1750442480735$	$-0.05422812947328 i)$
$+(-0.0086188585913$	$-0.39051763820462 i)\epsilon$
$+(+0.37518853545319$	$-0.34047477405516 i)\epsilon^2$
$+(+0.49765461883470$	$-0.00717399489427 i)\epsilon^3$
$+(+0.32835724868237$	$+0.23005850008124 i)\epsilon^4$
$+(+0.11199125312340$	$+0.25409725390712 i)\epsilon^5$
$+(-0.00954795237038$	$+0.17050760870656 i)\epsilon^6$
$+(-0.04217861994524$	$+0.08576862780838 i)\epsilon^7$

$$F_S(a_1, a_2, a_2; b_1, b_2, b_3; c, c, c, x, y, z) \tag{C.1}$$

$$= \frac{\Gamma(c)}{\Gamma(a_1)\Gamma(c - a_1)} \int_0^1 dt \frac{t^{c-a_1-1} (1-t)^{a_1-1}}{(1-x+tx)^{b_1}} F_1(a_2; b_2, b_3; c - a_1; ty, tz).$$

In our case, this becomes

$$F_S^b(d) = F_S\left(\frac{d-3}{2}, 1, 1; 1, 1, \frac{1}{2}; \frac{d}{2}, \frac{d}{2}, \frac{d}{2}; x_c, y_c, z_c\right) \tag{C.2}$$

$$= \frac{\Gamma(\frac{d}{2})}{\Gamma(\frac{d-3}{2})\Gamma(\frac{3}{2})} \times \int_0^1 dt \frac{\sqrt{t}(1-t)^{\frac{d-5}{2}}}{(1-x_c+x_ct)} F_1(1; 1, \frac{1}{2}; \frac{3}{2}; y_ct, z_ct)$$

Eqn. (C.2) is valid if $\Re(d) > 3$. With a grain of care one may often use (B.10) for F_1^S . Because the F_1 under the t -integral is finite and smooth, we have to concentrate only on the term $1/(1-x_c+x_ct)$,

which develops a pole in the integration region at $t_x = (1-x)/x$ if $\Re(x_c) = x > 1$ and if $\Im(x_c) = -\epsilon_x$ is infinitesimal.

Appendix C.1. Case (i) $F_S^b(d)$ at $x \leq 1$

For $x = 1$, the integral (C.2) is not well-defined. If $x < 1$, a direct, stable numerical integration of F_S is trivial once F_1 is known.

Appendix C.2. Case (ii) $F_S^b(d)$ at $x > 1$

If $x > 1$, one has to apply a regularization procedure to $F_S^b(d)$, as it is described in (Appendix B.2), and will get a stable result for F_S . The calculation of the F_1 in the integrand in (C.2) is discussed in Appendix B.1.

One now has to study the singularity structure of the t -integral as a function of x_c with regular F_1^S . Introduce

$$F_S^b(d) = \int_0^1 dt \frac{g_S(t) - g_S(t_x)}{1-x+xt} + g_S(t_x) I_S^{reg}(x_c), \tag{C.3}$$

with

$$g_S(t) = \sqrt{t} (1-t)^{(d-5)/2} F_1^S(y_ct, z_ct) \tag{C.4}$$

and

$$t_x = 1 - \frac{1}{x}. \tag{C.5}$$

The first integral in (C.3) is numerically stable, and what remains is to calculate analytically the integral

$$I_S^{reg}(x_c) = +\frac{1}{x_c} \int_0^1 \frac{dt}{t-t_x} = \frac{1}{x_c} \ln\left(1 - \frac{1}{t_x}\right). \tag{C.6}$$

For infinitesimal ϵ_x , we get

$$I_S^{reg}(x_c) \rightarrow \frac{1}{x} [-\ln(x-1) + i\pi \text{sign}(\epsilon_x)]. \tag{C.7}$$

References

- [1] A.I. Davydychev, *Phys. Lett. B* 263 (1991) 107–111.
- [2] J. Fleischer, T. Riemann, *Phys. Rev. D* 83 (2011) 073004.
- [3] J. Gluza, K. Kajda, T. Riemann, *Comput. Phys. Commun.* 177 (2007) 879–893.
- [4] I. Dubovyk, J. Gluza, T. Riemann, J. Usovitsch, in: *PoS LL2016*, 2016, p. 034.
- [5] K. Kajda, I. Dubovyk, *Mathematica package AMBRE* 2.2, 12 Sep 2015, <http://prac.us.edu.pl/~gluza/ambre/>.
- [6] I. Dubovyk, Mellin-Barnes representations for multiloop Feynman integrals with applications to 2-loop electroweak Z boson studies, Universität Hamburg, submitted for publication.
- [7] G. 't Hooft, M. Veltman, *Nucl. Phys. B* 153 (1979) 365–401.
- [8] U. Nierste, D. Müller, M. Böhm, *Z. Phys. C* 57 (1993) 605–614.
- [9] G. Passarino, *Nucl. Phys. B* 619 (2001) 257–312.
- [10] O. Tarasov, *Nucl. Phys. Proc. Suppl.* 89 (2000) 237.
- [11] J. Fleischer, F. Jegerlehner, O. Tarasov, *Nucl. Phys. B* 672 (2003) 303.
- [12] D.B. Melrose, *Nuovo Cimento* 40 (1965) 181–213.
- [13] K.H. Phan, J. Blümlein, T. Riemann, *Acta Phys. Pol. B* 48 (2017) 2313.
- [14] J. Usovitsch, I. Dubovyk, T. Riemann, in: *PoS LL2018*, 2018, p. 046.
- [15] J. Usovitsch, I. Dubovyk, and T. Riemann, The MBnumerics project, section E.2 in [32].
- [16] T. Hahn, M. Perez-Victoria, *Comput. Phys. Commun.* 118 (1999) 153–165.
- [17] G. van Oldenborgh, *Comput. Phys. Commun.* 66 (1991) 1–15.
- [18] P. Appell, *Mémoires des Sciences Mathématiques*, fascicule 3, 1925, p. 82.
- [19] S. Saran, *Acta Math.* 93 (1955) 293.
- [20] L. Slater, *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, 1966.
- [21] M. Czakon, *Comput. Phys. Commun.* 175 (2006) 559.
- [22] J. Usovitsch and T. Riemann, New approach to Mellin-Barnes integrals for massive one-loop Feynman integrals, section E.6 in [32].
- [23] J. Usovitsch, *Mathematica/Fortran package MBnumerics*.
- [24] J. Usovitsch, PhD Thesis, Humboldt-Universität zu Berlin Humboldt-Universität zu Berlin, 29 May 2018, https://edoc.hu-berlin.de/bitstream/handle/18452/20256/dissertation_usovitsch_johann.pdf?sequence=8&isAllowed=y, <https://doi.org/10.18452/19484>.
- [25] L. Pochhammer, *Crelle's J. Reine Angew. Math.* 71 (1870) 316.
- [26] A. Erdelyi, *Hypergeometric functions of two variables*, *Acta Math.* 83 (1950) 131.
- [27] V. Bytev, M. Kalmykov, B.A. Kniehl, *Comput. Phys. Commun.* 184 (2013) 2332.
- [28] V. Bytev, B.A. Kniehl, *Comput. Phys. Commun.* 189 (2015) 128.
- [29] V. Bytev, M. Kalmykov, S. Moch, *Comput. Phys. Commun.* 185 (2014) 3041.
- [30] E. Picard, *Ann. Sci. Éc. Norm. Supér.* 10 (2) (1881) 305.
- [31] M. Schlosser, in: *Proceedings of LHCPHenoNet School*, 2012, <https://doi.org/10.1007/978-3-7091-1616-6>.
- [32] A. Blondel, J. Gluza, S. Jadach, P. Janot, T. Riemann (eds.), *Standard Model Theory for the FCC-ee: The Tera-Z*, subm. as CERN Yellow Report, <http://www.arXiv.org/abs/1809.01830>.