



You have downloaded a document from  
**RE-BUŚ**  
repository of the University of Silesia in Katowice

**Title:** On a boundary value problem

**Author:** Józef Kalinowski

**Citation style:** Kalinowski Józef. (1993). On a boundary value problem.  
"Annales Mathematicae Silesianae" (Nr 7 (1993), s. 73-78).



Uznanie autorstwa - Użycie niekomercyjne - Bez utworów zależnych Polska - Licencja ta zezwala na rozpowszechnianie, przedstawianie i wykonywanie utworu jedynie w celach niekomercyjnych oraz pod warunkiem zachowania go w oryginalnej postaci (nie tworzenia utworów zależnych).



UNIwersYTET ŚLĄSKI  
W KATOWICACH



Biblioteka  
Uniwersytetu Śląskiego



Ministerstwo Nauki  
i Szkolnictwa Wyższego

## ON A BOUNDARY VALUE PROBLEM

JÓZEF KALINOWSKI

**Abstract.** The equation  $x''(t) = f(t, x(\alpha(t)), x'(\beta(t)))$  for  $t \in [a, b]$ , where the functions  $\alpha, \beta$  deviated argument of type  $[a, b] \rightarrow [a, b]$  is considered.

A sufficient condition for existence of the end  $b$  of the interval  $[a, b]$ , such that there exists the solution  $x$  of the above equation on  $[a, b]$  fulfilling the boundary value conditions  $x(a) = A$ ,  $x(b) = B$  and  $\|x'(a)\| = v > 0$ , where the constants  $a, v$  and vectors  $A, B$  are given, is proved.

Let  $D := [a, b]$  be an interval and  $d := b - a$  denote length of this interval. Let the symbol  $\|\cdot\|$  denote a norm in the space  $\mathbb{R}^n$ .

Consider a system of ordinary differential equations of the second order with a deviating argument of the form

$$(1) \quad x''(t) = f(t, x(\alpha(t)), x'(\beta(t))), \quad t \in D.$$

Let us denote  $D_1 := D \times \mathbb{R}^n \times \mathbb{R}^n$ . We assume that the function  $f : D_1 \rightarrow \mathbb{R}^n$  is a continuous real function and fulfils the Lipschitz condition of the form

$$(2) \quad \|f(t, x, y) - f(t, \bar{x}, \bar{y})\| \leq p \|x - \bar{x}\| + q \|y - \bar{y}\| \quad \text{for } t \in D,$$

where  $p, q \geq 0$  are constants. The function  $f$  is bounded on the domain  $D_1$ , i.e.

$$(3) \quad \|f(\cdot, \cdot, \cdot)\| \leq K.$$

---

*Received December 11, 1990.*

AMS (1991) subject classification: Primary 34K10. Secondary 34B15.

$\alpha, \beta : D \rightarrow D$ , functions of deviation of the argument, are continuous and  $a \leq \alpha(t) \leq t, a \leq \beta(t) \leq t$ .

We consider boundary value conditions for the system (1)

$$(4) \quad x(a) = A, \quad x(b) = B$$

and

$$(5) \quad \|x'(a)\| = v,$$

where vectors  $A, B \in \mathbb{R}^n$  and the constant  $v > 0$  are given. The right end  $b$  of the interval  $D$  is unknown.

In this paper the existence of the right end  $b$  of the interval  $D$  and the solution  $x$  of the problem (1), (4), (5) on  $D$  will be proved.

In particular case, for equation

$$x''(t) = g(x(t)), \quad t \in D$$

the similar problem was consider in the paper [3].

We will prove the following theorem:

**THEOREM.** *Let the function  $f$  satisfy assumptions (2) and (3). Let us assume that*

$$v > \begin{cases} h(d_1) & \text{for } d_1 < d_2 \\ h(d_2) & \text{for } d_1 \geq d_2, \end{cases}$$

where

$$h(d) := \frac{1}{d} \|B - A\| + \frac{K}{2} \cdot d, \quad d > 0$$

and

$$d_1 := \sqrt{\frac{2\|B - A\|}{K}},$$

$$d_2 := \frac{\sqrt{q^2 + 2p} - q}{p}$$

and

$$(7) \quad \frac{1}{2}d^2p + dq < 1.$$

If the vectors  $A$  and  $B$  satisfy the relation

$$(8) \quad A \neq B$$

then there exists the interval  $D$  and the solution  $x$  of the problem (1), (4), (5) on  $D$ .

PROOF. From the results of the papers [1], [2] and the assumptions (2) and (7) we obtain existence and uniqueness of the solution of the problem (1), (4) on  $D$ , where  $b > a$  and  $b$  is a parameter. From uniqueness of the solution it follows that the formula of the solution may be presented in the form

$$(9) \quad x(t) = \int_a^t \left[ \int_a^s f(z, x(\alpha(z)), x'(\beta(z))) dz \right] ds + M_b \cdot (t - a) + A,$$

for  $t \in D$ . From (9) it follows for  $t = b$  that the vector  $M_b \in \mathbb{R}^n$  is defined by formula

$$(10) \quad M_b = \frac{1}{d} \left\{ (B - A) - \int_a^b \left[ \int_a^s f(z, x(\alpha(z)), x'(\beta(z))) dz \right] \right\}.$$

Differentiating each side of the equation (9) with respect to the variable  $t$  we obtain

$$(11) \quad x'(t) = \int_a^t f(z, x(\alpha(z)), x'(\beta(z))) dz + M_b, \quad t \in D.$$

Using (10), the triangle inequality, the assumption (3) and properties of integrals we obtain

$$\begin{aligned} \| M_b \| &\leq \frac{1}{d} \left\{ \| B - A \| + \int_a^b \left[ \int_a^s \| f(z, x(\alpha(z)), x'(\beta(z))) \| dz \right] ds \right\} \\ &\leq \frac{1}{d} \left\{ \| B - A \| + \int_a^b \left[ \int_a^s K dz \right] ds \right\} = h(d) \end{aligned}$$

i.e.

$$(12) \quad \|M_b\| \leq h(d) \quad \text{for } d > 0.$$

From the definition (10) there exists

$$(13) \quad \lim_{d \rightarrow 0+} \|M_b\| = +\infty.$$

It follows from the definition of the function  $h$  that

$$\lim_{d \rightarrow 0+} h(d) = +\infty$$

and formulas (12), (13) are not in contradiction with themselves.

From continuity of the function  $M_b$  for  $d > 0$ , from (12) and (13), under (8) the norm  $\|M_b\|$  is greater than  $\min_{d>0} h(d)$ . But for satisfying the inequality (7) the argument  $d$  must fulfill the inequality  $d \leq d_2$ . Let us consider two possible cases:

(1°)  $d_1 < d_2$ .

From the definition of the function  $h$

$$\min_{0 \leq d \leq d_2} h(d) = h(d_1).$$

(2°)  $d_1 \geq d_2$ .

Then

$$\min_{0 \leq d \leq d_2} h(d) = h(d_2).$$

From uniqueness of the solution  $x$  and from (5) and (11) it follows that the equality

$$(14) \quad \|M_b\| = v$$

holds. The existence of  $b$  follows from (13) and continuity of  $\|M_b\|$ . Then for  $v$ , satisfying the inequality (6) there exist  $M_b$  defined by (10) and solution  $x$  of the form (9) of the problem (1), (4), (5).

This is the end of the proof. □

REMARK 1. Theorem is not true without the assumption (8).

PROOF. For the problem

$$x''(t) = 0, \quad x(a) = x(b) = 0, \quad t \in D.$$

there exists the unique constant solution  $x(t) = 0, t \in D$ . For all  $v > 0$  the condition (5) is not fulfilled.  $\square$

REMARK 2. The analogous theorem is true for the equation (1) with more than two deviations of the argument.

#### REFERENCES

- [1] J. Kalinowski, *Two point boundary value problem for a system of ordinary differential equations of second order with deviating argument*, Ann. Polon. Math. **30** (1974), (in Russian) 71-76.
- [2] J. Kalinowski, *On the convergence of an iterative sequence to the solution of a system of ordinary differential equations with deviated arguments*, Demonstratio Math. **9** No 1 (1976), 77-93.
- [3] C. A. Vavilov, *Solvability a class of boundary value problems*, Dokl. Akad. Nauk SSSR **305** No 2 (1989), (in Russian) 268-270.

UNIWERSYTET ŚLĄSKI  
INSTYTUT MATEMATYKI  
UL. BANKOWA 14  
40-007 KATOWICE