



You have downloaded a document from
RE-BUŚ
repository of the University of Silesia in Katowice

Title: A characterization of functions with dense graph in the plane or half-plane

Author: Zbigniew Gajda

Citation style: Gajda Zbigniew. (1986). A characterization of functions with dense graph in the plane or half-plane. "Annales Mathematicae Silesianae" (Nr 2 (1986), s. 37-46).



Uznanie autorstwa - Użycie niekomercyjne - Bez utworów zależnych Polska - Licencja ta zezwala na rozpowszechnianie, przedstawianie i wykonywanie utworu jedynie w celach niekomercyjnych oraz pod warunkiem zachowania go w oryginalnej postaci (nie tworzenia utworów zależnych).



UNIwersYTET ŚLĄSKI
W KATOWICACH



Biblioteka
Uniwersytetu Śląskiego



Ministerstwo Nauki
i Szkolnictwa Wyższego

A CHARACTERIZATION OF FUNCTIONS WITH DENSE GRAPH IN THE PLANE OR HALF-PLANE

Abstract. Let \mathbf{R} be the set of all real numbers. In the present paper we shall characterize functions $f: \mathbf{R} \rightarrow \mathbf{R}$ which are either linear or have graph contained and dense in the plane or half-plane determined by a linear function. For this purpose we consider functions satisfying certain limitary conditions which are related to the additivity equation but considerably weaker than that.

Let us introduce the following

DEFINITION. A function $f: \mathbf{R} \rightarrow \mathbf{R}$ is called *limit-additive* iff the following conditions are fulfilled:

- (1)
$$\bigwedge_{x, y \in \mathbf{R}} \bigvee_{\substack{(z_n)_{n \in \mathbf{N}} \\ z_n \in \mathbf{R}, n \in \mathbf{N}}} [z_n \xrightarrow{n \rightarrow \infty} x + y, f(z_n) \xrightarrow{n \rightarrow \infty} f(x) + f(y)],$$
- (2)
$$\bigwedge_{x, y \in \mathbf{R}} \bigvee_{\substack{(x_n)_{n \in \mathbf{N}}, (y_n)_{n \in \mathbf{N}} \\ x_n, y_n \in \mathbf{R}, n \in \mathbf{N}}} [x_n \xrightarrow{n \rightarrow \infty} x, y_n \xrightarrow{n \rightarrow \infty} y, f(x_n) + f(y_n) \xrightarrow{n \rightarrow \infty} f(x + y)],$$
- (3)
$$\bigwedge_{x \in \mathbf{R}} \bigvee_{\substack{(x_n)_{n \in \mathbf{N}} \\ x_n \in \mathbf{R}, n \in \mathbf{N}}} [x_n \xrightarrow{n \rightarrow \infty} x, 2f(x_n) \xrightarrow{n \rightarrow \infty} f(2x)].$$

Conditions (1) and (2) are, in a sense, mutually symmetric. Condition (3) can not be obtained from (2) by setting $x = y$, since even then sequences $(x_n)_{n \in \mathbf{N}}$ and $(y_n)_{n \in \mathbf{N}}$ occurring in (2) may not coincide. Adding condition (3) we obtain the possibility of the choice of a common sequence in the case where $x = y$.

Clearly, every additive function is limit-additive (it suffices to take constant sequences). There exist, however, limit-additive functions which are not additive. Indeed, one can easily check that an arbitrary function $f: \mathbf{R} \rightarrow \mathbf{R}$ with the graph being dense on the plane \mathbf{R}^2 is limit-additive. Let us note that if a function $f: \mathbf{R} \rightarrow \mathbf{R}$ is limit-additive and continuous then it is additive and consequently has the form

$$f(x) = ax, \quad x \in \mathbf{R},$$

where a is a constant.

Received March 15, 1982.

AMS (MOS) subject classification (1980). Primary 39B40, 26A99.

* Instytut Matematyki Uniwersytetu Śląskiego, Katowice, ul. Bankowa 14, Poland.

Now, we are going to give some necessary and sufficient conditions for a limit-additive function to be continuous.

LEMMA 1. *Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be a limit-additive function. Then, for any $k \in \mathbf{N}$, $k \geq 2$, and each points $x_1, \dots, x_k \in \mathbf{R}$, there exists a sequence $(z_n)_{n \in \mathbf{N}}$ of real numbers such that*

$$z_n \xrightarrow[n \rightarrow \infty]{} x_1 + \dots + x_k \text{ and } f(z_n) \xrightarrow[n \rightarrow \infty]{} f(x_1) + \dots + f(x_k).$$

PROOF. For $k=2$ the assertion of the lemma coincides with condition (1). Suppose that our lemma holds true for some $k \in \mathbf{N}$, $k \geq 2$ and for any system of k points $x_1, \dots, x_k \in \mathbf{R}$. Fix $k+1$ points $x_1, \dots, x_{k+1} \in \mathbf{R}$. On account of our assumption, there exists a sequence $(u_n)_{n \in \mathbf{N}}$ such that

$$u_n \xrightarrow[n \rightarrow \infty]{} x_1 + \dots + x_k, \quad f(u_n) \xrightarrow[n \rightarrow \infty]{} f(x_1) + \dots + f(x_k).$$

In view of (1), for each $n \in \mathbf{N}$ one can find a sequence $(w_{n,m})_{m \in \mathbf{N}}$ such that

$$w_{n,m} \xrightarrow[m \rightarrow \infty]{} u_n + x_{k+1} \text{ and } f(w_{n,m}) \xrightarrow[m \rightarrow \infty]{} f(u_n) + f(x_{k+1}).$$

Hence

$$\bigwedge_{n \in \mathbf{N}} \bigvee_{m'_n \in \mathbf{N}} \bigwedge_{m \geq m'_n} |w_{n,m} - u_n - x_{k+1}| < \frac{1}{n},$$

$$\bigwedge_{n \in \mathbf{N}} \bigvee_{m''_n \in \mathbf{N}} \bigwedge_{m \geq m''_n} |f(w_{n,m}) - f(u_n) - f(x_{k+1})| < \frac{1}{n}.$$

We put $m_n := \max(m'_n, m''_n)$, $n \in \mathbf{N}$ and $z_n := w_{n,m_n}$, $n \in \mathbf{N}$. Then we have

$$\begin{aligned} & |z_n - x_1 - \dots - x_{k+1}| \leq \\ & \leq |w_{n,m_n} - u_n - x_{k+1}| + |u_n - x_1 - \dots - x_k| < \frac{1}{n} + |u_n - x_1 - \dots - x_k| \xrightarrow[n \rightarrow \infty]{} 0, \\ & |f(z_n) - f(x_1) - \dots - f(x_{k+1})| \leq \\ & \leq |f(w_{n,m_n}) - f(u_n) - f(x_{k+1})| + |f(u_n) - f(x_1) - \dots - f(x_k)| < \\ & < \frac{1}{n} + |f(u_n) - f(x_1) - \dots - f(x_k)| \xrightarrow[n \rightarrow \infty]{} 0, \end{aligned}$$

whence

$$z_n \xrightarrow[n \rightarrow \infty]{} x_1 + \dots + x_{k+1} \text{ and } f(z_n) \xrightarrow[n \rightarrow \infty]{} f(x_1) + \dots + f(x_{k+1})$$

which, by induction, completes the proof.

THEOREM 1. *If a limit-additive function $f: \mathbf{R} \rightarrow \mathbf{R}$ is continuous at a point then it is continuous everywhere.*

Proof. Assume that f is continuous at the point $x_0 \in \mathbf{R}$.

(a) Let $(x_n)_{n \in \mathbf{N}}$ be an arbitrary sequence of real numbers convergent to zero.

Since

$$x_0 = (x_0 - x_n) + x_n, \quad n \in \mathbf{N}$$

from (1) it follows that, for each $n \in \mathbf{N}$, there exists a sequence $(z_{n,m})_{m \in \mathbf{N}}$ such that

$$z_{n,m} \xrightarrow{m \rightarrow \infty} x_0, \quad f(z_{n,m}) \xrightarrow{m \rightarrow \infty} f(x_0 - x_n) + f(x_n).$$

Hence

$$\bigwedge_{n \in \mathbf{N}} \bigvee_{m'_n \in \mathbf{N}} \bigwedge_{m \geq m'_n} |z_{n,m} - x_0| < \frac{1}{n},$$

$$\bigwedge_{n \in \mathbf{N}} \bigvee_{m''_n \in \mathbf{N}} \bigwedge_{m \geq m''_n} |f(z_{n,m}) - f(x_0 - x_n) - f(x_n)| < \frac{1}{n}.$$

Put $m_n := \max(m'_n, m''_n)$, $z_n := z_{n,m_n}$, $n \in \mathbf{N}$. Then

$$|z_n - x_0| < \frac{1}{n}, \quad n \in \mathbf{N}$$

and

$$|f(z_n) - f(x_0 - x_n) - f(x_n)| < \frac{1}{n}, \quad n \in \mathbf{N},$$

whence

$$(4) \quad z_n \xrightarrow{n \rightarrow \infty} x_0 \quad \text{and} \quad f(z_n) - f(x_0 - x_n) - f(x_n) \xrightarrow{n \rightarrow \infty} 0.$$

By the continuity of f at x_0 we have

$$f(z_n) \xrightarrow{n \rightarrow \infty} f(x_0) \quad \text{and} \quad f(x_0 - x_n) \xrightarrow{n \rightarrow \infty} f(x_0)$$

which, together with (4), gives

$$(5) \quad f(x_n) \xrightarrow{n \rightarrow \infty} 0, \quad \text{for any sequence } (x_n)_{n \in \mathbf{N}} \text{ such that } x_n \xrightarrow{n \rightarrow \infty} 0.$$

(b) Fix an $x \in \mathbf{R}$ and write $0 = x + (-x)$. Using condition (1) again we find a sequence $(z_n)_{n \in \mathbf{N}}$, $z_n \xrightarrow{n \rightarrow \infty} 0$, for which

$$f(z_n) \xrightarrow{n \rightarrow \infty} f(x) + f(-x).$$

Hence and from (5) it follows that

$$f(-x) = -f(x), \quad x \in \mathbf{R}.$$

(c) Now, choose an arbitrary $x \in \mathbf{R}$ and a sequence $(x_n)_{n \in \mathbf{N}}$, $x_n \xrightarrow{n \rightarrow \infty} x$. On account of Lemma 1, for each $n \in \mathbf{N}$ one can find a sequence $(z_{n,m})_{m \in \mathbf{N}}$ such that

$$(6) \quad z_{n,m} \xrightarrow{m \rightarrow \infty} x_n - x + x_0$$

and

$$(7) \quad f(z_{n,m}) \xrightarrow{m \rightarrow \infty} f(x_n) + f(-x) + f(x_0) = f(x_n) - f(x) + f(x_0).$$

In view of (6) and (7) we have

$$\bigwedge_{n \in \mathbf{N}} \bigvee_{m'_n \in \mathbf{N}} \bigwedge_{m \geq m'_n} |z_{n,m} - (x_n - x + x_0)| < \frac{1}{n},$$

$$\bigwedge_{n \in \mathbf{N}} \bigvee_{m''_n \in \mathbf{N}} \bigwedge_{m \geq m''_n} |f(z_{n,m}) - (f(x_n) - f(x) + f(x_0))| < \frac{1}{n}.$$

We put $m_n := \max(m'_n, m_n)$, $z_n := z_{n, m_n}$, $n \in N$. With the aid of this notion we get

$$|z_n - x_0| \leq |z_{n, m_n} - (x_n - x + x_0)| + |x_n - x| < \frac{1}{n} + |x_n - x| \xrightarrow{n \rightarrow \infty} 0,$$

$$|f(z_n) - (f(x_n) - f(x) + f(x_0))| < \frac{1}{n} \xrightarrow{n \rightarrow \infty} 0.$$

Consequently,

$$z_n \xrightarrow{n \rightarrow \infty} x_0 \text{ and } f(z_n) - f(x_n) + f(x) - f(x_0) \xrightarrow{n \rightarrow \infty} 0.$$

Hence and from the continuity of f at the point x_0 it follows that

$$f(x_n) \xrightarrow{n \rightarrow \infty} f(x),$$

which implies that f is continuous at x .

LEMMA 2. *If a function $f: \mathbf{R} \rightarrow \mathbf{R}$ is limit-additive and bounded in a neighbourhood of a point $x_0 \in \mathbf{R}$ then it is bounded in a neighbourhood of zero.*

Proof. Suppose that there exist $M > 0$ and $\delta > 0$ such that

$$|f(y)| \leq M, \text{ for } y \in (x_0 - \delta, x_0 + \delta).$$

Take an $x \in (-\delta, \delta)$. Then $x + x_0 \in (x_0 - \delta, x_0 + \delta)$ and there exists a sequence $(z_n)_{n \in N}$, $z_n \xrightarrow{n \rightarrow \infty} x + x_0$ such that $f(z_n) \xrightarrow{n \rightarrow \infty} f(x) + f(x_0)$.

For almost every $n \in N$ we have

$$z_n \in (x_0 - \delta, x_0 + \delta) \text{ and } |f(z_n)| \leq M$$

whence

$$|f(x) + f(x_0)| \leq M.$$

Thus

$$|f(x)| \leq M + |f(x_0)|, \text{ for } x \in (-\delta, \delta).$$

THEOREM 2. *If $f: \mathbf{R} \rightarrow \mathbf{R}$ is a limit-additive function bounded (in absolute value) on a set $A \subset \mathbf{R}$ such that $\text{int } A \neq \emptyset$ then f is continuous.*

Proof. Taking Lemma 2 into account, we may suppose that there exist $M > 0$ and $\delta > 0$ such that

$$(8) \quad |f(x)| \leq M \text{ for } x \in (-\delta, \delta).$$

Assume that there exists a sequence of real numbers $(x_n)_{n \in N}$, $x_n \xrightarrow{n \rightarrow \infty} 0$ such that the sequence $(f(x_n))_{n \in N}$ is not convergent to zero. Then there exist an $\varepsilon > 0$ and a subsequence $(x_{n_k})_{k \in N}$ of the sequence $(x_n)_{n \in N}$ with the property $|f(x_{n_k})| \geq \varepsilon$, $k \in N$.

From the sequence $(x_{n_k})_{k \in N}$ one can still choose either a subsequence $(x_{n_{k_p}})_{p \in N}$ such that $f(x_{n_{k_p}}) \geq \varepsilon$, $p \in N$ or a subsequence $(x_{n_{k_s}})_{s \in N}$ such that $f(x_{n_{k_s}}) \leq -\varepsilon$, $s \in N$.

Suppose, for instance, that we have a sequence $(y_n)_{n \in N}$, $y_n \xrightarrow{n \rightarrow \infty} 0$ such that $f(y_n) \geq \varepsilon$, $n \in N$. Let us choose numbers $N \in N$ and $n_0 \in N$ so that $N\varepsilon > M$ and $Ny_{n_0} \in (-\delta, \delta)$. According to Lemma 1, there exists a sequence $(z_m)_{m \in N}$ such that $z_m \xrightarrow{m \rightarrow \infty} N \cdot y_{n_0}$ and $f(z_m) \xrightarrow{m \rightarrow \infty} Nf(y_{n_0}) \geq N\varepsilon > M$.

Hence

$$(9) \quad \bigvee_{m_1 \in \mathbf{N}} \bigwedge_{m \geq m_1} z_m \in (-\delta, \delta),$$

$$(10) \quad \bigvee_{m_2 \in \mathbf{N}} \bigwedge_{m \geq m_2} f(z_m) > M.$$

For $m \geq \max(m_1, m_2)$ conditions (9) and (10) are incompatible with (8). If we have a sequence $(y_n)_{n \in \mathbf{N}}$, $y_n \xrightarrow{n \rightarrow \infty} 0$ such that $f(y_n) \leq -\varepsilon, n \in \mathbf{N}$, we obtain the contradiction in a similar manner, using the boundedness of f from below. So we have

$$(11) \quad f(x_n) \xrightarrow{n \rightarrow \infty} 0, \text{ for any sequence } (x_n)_{n \in \mathbf{N}} \text{ such that } x_n \xrightarrow{n \rightarrow \infty} 0.$$

Putting $x = y = 0$ in (1), we obtain the existence of a sequence $(z_n)_{n \in \mathbf{N}}$, $z_n \xrightarrow{n \rightarrow \infty} 0$, for which $f(z_n) \xrightarrow{n \rightarrow \infty} 2f(0)$. This, jointly with (11), implies $f(0) = 0$. Consequently we obtain the continuity of f at zero. In virtue of Theorem 1, f is continuous everywhere on \mathbf{R} .

Now, we are going to investigate some properties of discontinuous limit-additive functions. It follows from Theorem 2 that such functions can not be bounded in absolute value on any non-degenerate interval. In the sequel, the word "interval" will always mean a bounded non-degenerate interval. The example of an arbitrary function $f: \mathbf{R} \rightarrow \mathbf{R}$ which has the graph contained and dense in one of the half-planes $\{(x, y) \in \mathbf{R}^2 : y \geq 0\}$ or $\{(x, y) \in \mathbf{R}^2 : y \leq 0\}$ shows that a discontinuous limit-additive function may be bounded from one side. In the same way as in the proof of Lemma 2 one can show that any limit-additive function bounded below (above) on some interval is bounded below (above) on every interval.

For any function $f: \mathbf{R} \rightarrow \mathbf{R}$ bounded below on every interval, the function $\varphi_f: \mathbf{R} \rightarrow \mathbf{R}$

$$(12) \quad \varphi_f(x) := \sup_{\delta > 0} \inf \{f(z) : z \in (x - \delta, x + \delta)\}, \quad x \in \mathbf{R}$$

is well defined.

Analogously, for any function $f: \mathbf{R} \rightarrow \mathbf{R}$ bounded above on every interval we define the function $\psi_f: \mathbf{R} \rightarrow \mathbf{R}$ by the formula

$$(13) \quad \psi_f(x) := \inf_{\delta > 0} \sup \{f(z) : z \in (x - \delta, x + \delta)\}, \quad x \in \mathbf{R}.$$

LEMMA 3. *If $f: \mathbf{R} \rightarrow \mathbf{R}$ is bounded below (above) on every interval, then the function φ_f (function ψ_f) is lower (upper) semi-continuous.*

For the proof see e.g. [2] or [3].

Up to now, we have only made use of property (1) from the definition of limit-additive functions. From now on, we shall be applying properties (2) and (3), too.

LEMMA 4. *If $f: \mathbf{R} \rightarrow \mathbf{R}$ is a limit-additive function bounded below (above) on every interval, then the function φ_f (function ψ_f) is additive.*

Proof. We proceed only with the proof for the function φ_f . Fix numbers $x, y \in \mathbf{R}$, $\varepsilon > 0$, $\delta > 0$, $\eta > 0$, arbitrarily. We have

$$(14) \quad \bigvee_{u_0 \in (x - \delta, x + \delta)} f(u_0) < \inf \{f(u) : u \in (x - \delta, x + \delta)\} + \frac{\varepsilon}{3}$$

and

$$(15) \quad \bigvee_{w_0 \in (y-\eta, y+\eta)} f(w_0) < \inf \{f(w) : w \in (y-\eta, y+\eta)\} + \frac{\varepsilon}{3}.$$

Observe that $u_0 + w_0 \in (x+y-\delta-\eta, x+y+\delta+\eta)$. It follows from (1) that there exists a sequence $(z_n)_{n \in \mathbf{N}}$, $z_n \xrightarrow{n \rightarrow \infty} u_0 + w_0$ such that

$$f(z_n) \xrightarrow{n \rightarrow \infty} f(u_0) + f(w_0).$$

Hence

$$(16) \quad \bigvee_{n_0 \in \mathbf{N}} \bigwedge_{n \geq n_0} \left[z_n \in (x+y-\delta-\eta, x+y+\delta+\eta), f(z_n) < f(u_0) + f(w_0) + \frac{\varepsilon}{3} \right].$$

(14), (15) and (16) yield

$$\begin{aligned} \inf \{f(z) : z \in (x+y-\delta-\eta, x+y+\delta+\eta)\} &\leq f(u_0) + f(w_0) + \frac{\varepsilon}{3} \leq \\ &\leq \inf \{f(u) : u \in (x-\delta, x+\delta)\} + \inf \{f(w) : w \in (y-\eta, y+\eta)\} + \varepsilon. \end{aligned}$$

Since $\varepsilon > 0$ has been chosen arbitrarily, we have

$$(17) \quad \inf \{f(z) : z \in (x+y-\delta-\eta, x+y+\delta+\eta)\} \leq \inf \{f(u) : u \in (x-\delta, x+\delta)\} + \inf \{f(w) : w \in (y-\eta, y+\eta)\} \leq \varphi_f(x) + \varphi_f(y).$$

As inequality (17) holds for all $\delta > 0$, $\eta > 0$, we obtain the subadditivity of φ_f :

$$(18) \quad \varphi_f(x+y) \leq \varphi_f(x) + \varphi_f(y), \quad x, y \in \mathbf{R}.$$

Fix again numbers $x, y \in \mathbf{R}$, $\varepsilon > 0$. $\delta > 0$ arbitrarily. We have

$$(19) \quad \bigvee_{z_0 \in (x+y-\delta, x+y+\delta)} f(z_0) < \inf \{f(z) : z \in (x+y-\delta, x+y+\delta)\} + \frac{\varepsilon}{2}.$$

One can choose $u_0 \in \left(x - \frac{\delta}{2}, x + \frac{\delta}{2}\right)$ and $w_0 \in \left(y - \frac{\delta}{2}, y + \frac{\delta}{2}\right)$ so that $z_0 = u_0 + w_0$.

In view of (2) there exist sequences $(u_n)_{n \in \mathbf{N}}$ and $(w_n)_{n \in \mathbf{N}}$, $u_n \xrightarrow{n \rightarrow \infty} u_0$, $w_n \xrightarrow{n \rightarrow \infty} w_0$ such that $f(u_n) + f(w_n) \xrightarrow{n \rightarrow \infty} f(z_0)$. Hence

$$(20) \quad \bigvee_{n_0 \in \mathbf{N}} \bigwedge_{n \geq n_0} \left[u_n \in \left(x - \frac{\delta}{2}, x + \frac{\delta}{2}\right), w_n \in \left(y - \frac{\delta}{2}, y + \frac{\delta}{2}\right), f(u_n) + f(w_n) < f(z_0) + \frac{\varepsilon}{2} \right].$$

From (19) and (20) we obtain

$$\begin{aligned} \inf \left\{ f(u) : u \in \left(x - \frac{\delta}{2}, x + \frac{\delta}{2}\right) \right\} + \inf \left\{ f(w) : w \in \left(y - \frac{\delta}{2}, y + \frac{\delta}{2}\right) \right\} &\leq \\ &\leq f(z_0) + \frac{\varepsilon}{2} < \inf \{f(z) : z \in (x+y-\delta, x+y+\delta)\} + \varepsilon. \end{aligned}$$

Letting ε tend to zero we get

$$(21) \quad \inf \left\{ f(u) : u \in \left(x - \frac{\delta}{2}, x + \frac{\delta}{2} \right) \right\} + \inf \left\{ f(w) : w \in \left(y - \frac{\delta}{2}, y + \frac{\delta}{2} \right) \right\} \leq \\ \leq \inf \{ f(z) : z \in (x + y - \delta, x + y + \delta) \} \leq \varphi_f(x + y).$$

Since inequality (21) holds true for all $\delta > 0$, the function φ_f is superadditive:

$$(22) \quad \varphi_f(x) + \varphi_f(y) \leq \varphi_f(x + y), \quad x, y \in \mathbf{R}.$$

Conjunction of conditions (18) and (22) gives the additivity of φ_f . In the same manner one may prove that condition (1) leads to superadditivity of ψ_f and condition (2) to its subadditivity.

As is well known any lower (upper) semi-continuous function is bounded below (above) on every compact interval. Hence and from Lemmas 3 and 4 as well as from the properties of the additive functions we obtain immediately the following

THEOREM 3. *If $f: \mathbf{R} \rightarrow \mathbf{R}$ is a limit-additive function bounded below (above) on every interval, then the function φ_f (function ψ_f) is additive and continuous.*

LEMMA 5. *Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be a limit-additive function. For any $x \in \mathbf{R}$ and each $k \in \mathbf{N}$ there exists a sequence of real numbers $(x_n)_{n \in \mathbf{N}}$ such that*

$$x_n \xrightarrow{n \rightarrow \infty} x \quad \text{and} \quad 2^k f(x_n) \xrightarrow{n \rightarrow \infty} f(2^k x).$$

Proof. For $k = 1$ the assertion of our lemma coincides with condition (3). Suppose that this assertion holds true for any $x \in \mathbf{R}$ and some $k \in \mathbf{N}$. Therefore, for arbitrarily fixed $x \in \mathbf{R}$ there exists a sequence $(y_n)_{n \in \mathbf{N}}$ such that

$$y_n \xrightarrow{n \rightarrow \infty} 2x \quad \text{and} \quad 2^k f(y_n) \xrightarrow{n \rightarrow \infty} f(2^{k+1}x).$$

From (3) it follows that to each $n \in \mathbf{N}$ there corresponds a sequence $(x_{n,m})_{m \in \mathbf{N}}$ such that

$$x_{n,m} \xrightarrow{m \rightarrow \infty} \frac{y_n}{2} \quad \text{and} \quad 2f(x_{n,m}) \xrightarrow{m \rightarrow \infty} f(y_n).$$

Hence

$$\bigwedge_{n \in \mathbf{N}} \bigvee_{m'_n \in \mathbf{N}} \bigwedge_{m \geq m'_n} \left| x_{n,m} - \frac{y_n}{2} \right| < \frac{1}{n}, \\ \bigwedge_{n \in \mathbf{N}} \bigvee_{m''_n \in \mathbf{N}} \bigwedge_{m \geq m''_n} |2f(x_{n,m}) - f(y_n)| < \frac{1}{n}.$$

Put $m_n := \max(m'_n, m''_n)$, $x_n := x_{n,m_n}$, for $n \in \mathbf{N}$. Then we get

$$|x_n - x| \leq \left| x_{n,m_n} - \frac{y_n}{2} \right| + \left| \frac{y_n}{2} - x \right| \leq \frac{1}{n} + \frac{1}{2} |y_n - 2x| \xrightarrow{n \rightarrow \infty} 0$$

and

$$\begin{aligned} |2^{k+1}f(x_n) - f(2^{k+1}x)| &\leq |2^{k+1}f(x_{n,m_n}) - 2^k f(y_n)| + |2^k f(y_n) - f(2^{k+1}x)| \leq \\ &\leq 2^k \frac{1}{n} + |2^k f(y_n) - f(2^{k+1}x)| \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Consequently

$$x_n \xrightarrow{n \rightarrow \infty} x \text{ and } 2^{k+1}f(x_n) \xrightarrow{n \rightarrow \infty} f(2^{k+1}x).$$

By induction, the assertion of our lemma holds true for any $k \in \mathbf{N}$.

LEMMA 6. *Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be a limit-additive function. For any $x \in \mathbf{R}$, $l, k \in \mathbf{N}$, $r := \frac{l}{2^k}$ there exists a sequence of real numbers $(x_n)_{n \in \mathbf{N}}$ such that $x_n \xrightarrow{n \rightarrow \infty} rx$ and $f(x_n) \xrightarrow{n \rightarrow \infty} rf(x)$.*

Proof. Fix $x \in \mathbf{R}$, $l, k \in \mathbf{N}$, $r := \frac{l}{2^k}$. On account of Lemma 5 there exists a sequence $(y_n)_{n \in \mathbf{N}}$ with the property

$$y_n \xrightarrow{n \rightarrow \infty} \frac{x}{2^k} \text{ and } f(y_n) \xrightarrow{n \rightarrow \infty} \frac{1}{2^k} f(x).$$

Hence

$$ly_n \xrightarrow{n \rightarrow \infty} rx \text{ and } lf(y_n) \xrightarrow{n \rightarrow \infty} rf(x).$$

In view of (1), for each $n \in \mathbf{N}$ one can find a sequence $(x_{n,m})_{m \in \mathbf{N}}$ such that

$$x_{n,m} \xrightarrow{m \rightarrow \infty} ly_n \text{ and } f(x_{n,m}) \xrightarrow{m \rightarrow \infty} lf(y_n)$$

which implies that

$$\begin{aligned} \bigwedge_{n \in \mathbf{N}} \bigvee_{m'_n \in \mathbf{N}} \bigwedge_{m \geq m'_n} |x_{n,m} - ly_n| &< \frac{1}{n}, \\ \bigwedge_{n \in \mathbf{N}} \bigvee_{m''_n \in \mathbf{N}} \bigwedge_{m \geq m''_n} |f(x_{n,m}) - lf(y_n)| &< \frac{1}{n}. \end{aligned}$$

Setting $m_n := \max(m'_n, m''_n)$, $x_n := x_{n,m_n}$, $n \in \mathbf{N}$ we obtain

$$|x_n - rx| \leq |x_{n,m_n} - ly_n| + |ly_n - rx| \leq \frac{1}{n} + |ly_n - rx| \xrightarrow{n \rightarrow \infty} 0,$$

$$|f(x_n) - rf(x)| \leq |f(x_{n,m_n}) - lf(y_n)| + |lf(y_n) - rf(x)| \leq \frac{1}{n} + |lf(y_n) - rf(x)| \xrightarrow{n \rightarrow \infty} 0$$

which ends the proof.

LEMMA 7. *Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be a limit-additive function. For any $x, y \in \mathbf{R}$, $l, k \in \mathbf{N}$, $l < 2^k$, $r := \frac{l}{2^k}$ there exists a sequence $(z_n)_{n \in \mathbf{N}}$ such that*

$$z_n \xrightarrow{n \rightarrow \infty} rx + (1-r)y \text{ and } f(z_n) \xrightarrow{n \rightarrow \infty} rf(x) + (1-r)f(y).$$

Proof. According to Lemma 6 there exist sequences $(x_n)_{n \in \mathbf{N}}$, $(y_n)_{n \in \mathbf{N}}$ such that

$$\begin{aligned} x_n &\xrightarrow{n \rightarrow \infty} rx, & f(x_n) &\xrightarrow{n \rightarrow \infty} rf(x), \\ y_n &\xrightarrow{n \rightarrow \infty} (1-r)y, & f(y_n) &\xrightarrow{n \rightarrow \infty} (1-r)f(y). \end{aligned}$$

From (1) it follows that for each $n \in N$ there exists a sequence $(z_{n,m})_{m \in N}$ such that

$$z_{n,m} \xrightarrow{m \rightarrow \infty} x_n + y_n, \quad f(z_{n,m}) \xrightarrow{m \rightarrow \infty} f(x_n) + f(y_n).$$

Hence

$$\bigwedge_{n \in N} \bigvee_{m'_n \in N} \bigwedge_{m \geq m'_n} |z_{n,m} - x_n - y_n| < \frac{1}{n},$$

$$\bigwedge_{n \in N} \bigvee_{m''_n \in N} \bigwedge_{m \geq m''_n} |f(z_{n,m}) - f(x_n) - f(y_n)| < \frac{1}{n}.$$

Putting $m_n := \max(m'_n, m''_n)$, $z_n := z_{n,m_n}$, for $n \in N$ we get

$$\begin{aligned} |z_n - rx - (1-r)y| &\leq |z_{n,m_n} - x_n - y_n| + |x_n - rx| + |y_n - (1-r)y| \leq \\ &\leq \frac{1}{n} + |x_n - rx| + |y_n - (1-r)y| \xrightarrow{n \rightarrow \infty} 0, \\ |f(z_n) - rf(x) - (1-r)f(y)| &\leq |f(z_{n,m_n}) - f(x_n) - f(y_n)| + |f(x_n) - rf(x)| + \\ &+ |f(y_n) - (1-r)f(y)| \leq \frac{1}{n} + |f(x_n) - rf(x)| + |f(y_n) - (1-r)f(y)| \xrightarrow{n \rightarrow \infty} 0 \end{aligned}$$

which completes the proof.

Recall that by the *graph* of a function $f: \mathbf{R} \rightarrow \mathbf{R}$ we mean the set $\{(x, y) \in \mathbf{R}^2 : y = f(x)\}$. We consider the plane \mathbf{R}^2 with its natural topology.

THEOREM 4. *If $f: \mathbf{R} \rightarrow \mathbf{R}$ is a limit-additive function, then the following four cases are the only possible:*

- (i) f is an additive and continuous function;
- (ii) f is a function with the dense graph in \mathbf{R}^2 ;
- (iii) there exists an additive and continuous function $\varphi_f: \mathbf{R} \rightarrow \mathbf{R}$ such that the graph of f is contained and dense in the half-plane $\{(x, y) \in \mathbf{R}^2 : y \geq \varphi_f(x)\}$;
- (iv) there exists an additive and continuous function $\psi_f: \mathbf{R} \rightarrow \mathbf{R}$ such that graph of f is contained and dense in the half-plane $\{(x, y) \in \mathbf{R}^2 : y \leq \psi_f(x)\}$.

Conversely, every function fulfilling one of the conditions (i)–(iv) is limit-additive.

Proof. Suppose $f: \mathbf{R} \rightarrow \mathbf{R}$ to be limit-additive. In virtue of the previous theorems and lemmas the following cases are the only possible:

- (i) f is an additive and continuous function;
- (ii') the restriction of f to any interval is unbounded from above and from below;
- (iii') f is a function bounded from below and unbounded from above on every interval;
- (iv') f is a function bounded from above and unbounded from below on every interval.

Suppose that (ii') holds and choose an arbitrary rectangle $(a, b) \times (c, d)$. Since the set $A := \left\{ r = \frac{l}{2^k} : l, k \in N, l < 2^k \right\}$ is dense in the interval $(0, 1)$, we deduce that

$$\bigvee_{r \in A} rf(x) + (1-r)f(y) \in (c, d)$$

provided $f(x) < c$, $f(y) > d$; the existence of such a pair $(x, y) \in (a, b)^2$ results from our assumption. Let $(z_n)_{n \in \mathbf{N}}$ be such a sequence that

$$z_n \xrightarrow[n \rightarrow \infty]{} rx + (1-r)y \quad \text{and} \quad f(z_n) \xrightarrow[n \rightarrow \infty]{} rf(x) + (1-r)f(y).$$

Hence, for sufficiently large $n \in \mathbf{N}$, we have $(z_n, f(z_n)) \in (a, b) \times (c, d)$. Now, suppose that (iii') holds and let $\varphi_f: \mathbf{R} \rightarrow \mathbf{R}$ denote the function defined by (12); φ_f is additive and continuous. Moreover, the definition of φ_f yields $f(x) \geq \varphi_f(x)$, for $x \in \mathbf{R}$. Suppose that $(a, b) \times (c, d) \subset \{(x, y) \in \mathbf{R}^2 : y > \varphi_f(x)\}$. Then

$$c > \varphi_f\left(\frac{a+b}{2}\right) \geq \inf\{f(x) : x \in (a, b)\}$$

whence

$$\bigvee_{x \in (a, b)} f(x) < c.$$

Since f is not upper-bounded on (a, b) , one can find a $y \in (a, b)$ such that $f(y) > d$. Proceeding further in the same way as in case (ii') we prove that there exists a $z \in (a, b)$ such that $f(z) \in (c, d)$. Consequently, condition (iii) holds true. Using the properties of the function ψ_f defined by (13) one can show that (iv') implies (iv). It is easy to check the converse: every function $f: \mathbf{R} \rightarrow \mathbf{R}$ fulfilling one of the conditions (i)—(iv) is limit-additive.

Our last theorem gives full description of the class of limit-additive functions.

REFERENCES

- [1] J. ACZÉL, *Lectures on functional equations and their applications*, Academic Press, New York and London, 1966.
- [2] S. ŁOJASIEWICZ, *Wstęp do teorii funkcji rzeczywistych*, PWN, Warszawa, 1976.
- [3] B. SZ-NAGY, *Introduction to real functions and orthogonal expansions*, Akadémiai Kiado, Budapest, 1964.