



You have downloaded a document from
RE-BUŚ
repository of the University of Silesia in Katowice

Title: Remarks connected with the weak limit of iterates of some random-valued functions and iterative functional equations

Author: Karol Baron

Citation style: Baron Karol. (2020). Remarks connected with the weak limit of iterates of some random-valued functions and iterative functional equations. "Annales Mathematicae Silesianae" Vol. 34, No 1 (2020), s. 36-44.
DOI: 10.2478/amsil-2019-0015



Uznanie autorstwa - Licencja ta pozwala na kopiowanie, zmienianie, rozprowadzanie, przedstawianie i wykonywanie utworu jedynie pod warunkiem oznaczenia autorstwa.



UNIwersYTET ŚLĄSKI
W KATOWICACH



Biblioteka
Uniwersytetu Śląskiego



Ministerstwo Nauki
i Szkolnictwa Wyższego

REMARKS CONNECTED WITH THE WEAK LIMIT OF ITERATES OF SOME RANDOM-VALUED FUNCTIONS AND ITERATIVE FUNCTIONAL EQUATIONS

KAROL BARON 

Dedicated to Professor Zygfryd Kominek on his 75th birthday

Abstract. The paper consists of two parts. At first, assuming that (Ω, \mathcal{A}, P) is a probability space and (X, ϱ) is a complete and separable metric space with the σ -algebra \mathcal{B} of all its Borel subsets we consider the set \mathcal{R}_c of all $\mathcal{B} \otimes \mathcal{A}$ -measurable and contractive in mean functions $f: X \times \Omega \rightarrow X$ with finite integral $\int_{\Omega} \varrho(f(x, \omega), x) P(d\omega)$ for $x \in X$, the weak limit π^f of the sequence of iterates of $f \in \mathcal{R}_c$, and investigate continuity-like property of the function $f \mapsto \pi^f$, $f \in \mathcal{R}_c$, and Lipschitz solutions φ that take values in a separable Banach space of the equation

$$\varphi(x) = \int_{\Omega} \varphi(f(x, \omega)) P(d\omega) + F(x).$$

Next, assuming that X is a real separable Hilbert space, $\Lambda: X \rightarrow X$ is linear and continuous with $\|\Lambda\| < 1$, and μ is a probability Borel measure on X with finite first moment we examine continuous at zero solutions $\varphi: X \rightarrow \mathbb{C}$ of the equation

$$\varphi(x) = \hat{\mu}(x)\varphi(\Lambda x)$$

which characterizes the limit distribution π^f for some special $f \in \mathcal{R}_c$.

Received: 21.05.2019. Accepted: 19.12.2019. Published online: 08.05.2020.

(2010) Mathematics Subject Classification: 26A18, 39B12, 39B32, 39B52, 60B12, 60B15.

Key words and phrases: random-valued functions, iterates, convergence in law, continuous dependence on the given function, Fourier transform, iterative functional equations, continuous and Lipschitz solutions.

©2020 The Author(s).

This is an Open Access article distributed under the terms of the Creative Commons Attribution License CC BY (<http://creativecommons.org/licenses/by/4.0/>).

Fix a probability space (Ω, \mathcal{A}, P) and a complete and separable metric space (X, ϱ) . Let \mathcal{B} denote the σ -algebra of all Borel subsets of X .

We say that $f: X \times \Omega \rightarrow X$ is a *random-valued function* (an *rv-function* for short) if it is measurable with respect to the product σ -algebra $\mathcal{B} \otimes \mathcal{A}$. The iterates of such an rv-function are given by

$$f^0(x, \omega_1, \omega_2, \dots) = x, \quad f^n(x, \omega_1, \omega_2, \dots) = f(f^{n-1}(x, \omega_1, \omega_2, \dots), \omega_n)$$

for x from X and $(\omega_1, \omega_2, \dots)$ from Ω^∞ defined as $\Omega^{\mathbb{N}}$. Note that $f^n: X \times \Omega^\infty \rightarrow X$ is an rv-function on the product probability space $(\Omega^\infty, \mathcal{A}^\infty, P^\infty)$. More exactly, the n -th iterate f^n is $\mathcal{B} \otimes \mathcal{A}_n$ -measurable, where \mathcal{A}_n denotes the σ -algebra of all sets of the form

$$\{(\omega_1, \omega_2, \dots) \in \Omega^\infty : (\omega_1, \dots, \omega_n) \in A\}$$

with A from the product σ -algebra \mathcal{A}^n . (See [7, Sec. 1.4], [5].)

A simple criterion for the convergence in law of $(f^n(x, \cdot))_{n \in \mathbb{N}}$ to a random variable independent of $x \in X$ was proved in [1] and applied to the equation

$$(1) \quad \varphi(x) = \int_{\Omega} \varphi(f(x, \omega)) P(d\omega) + F(x)$$

with φ as the unknown function. This criterion reads.

(H) There exists a $\lambda \in (0, 1)$ such that

$$\int_{\Omega} \varrho(f(x, \omega), f(z, \omega)) P(d\omega) \leq \lambda \varrho(x, z) \quad \text{for } x, z \in X$$

and

$$(2) \quad \int_{\Omega} \varrho(f(x, \omega), x) P(d\omega) < \infty \quad \text{for } x \in X.$$

Thus, denoting by $\pi_n^f(x, \cdot)$ the distribution of $f^n(x, \cdot)$, i.e.,

$$\pi_n^f(x, B) = P^\infty(f^n(x, \cdot) \in B) \quad \text{for } n \in \mathbb{N} \cup \{0\}, x \in X \text{ and } B \in \mathcal{B},$$

hypothesis (H) guarantees the existence of a probability Borel measure π^f on X such that

$$\lim_{n \rightarrow \infty} \int_X u(z) \pi_n^f(x, dz) = \int_X u(z) \pi^f(dz)$$

for $x \in X$ and for any continuous and bounded $u: X \rightarrow \mathbb{R}$; moreover, as observed in [3] (see also [6]),

$$(3) \quad \int_X \varrho(x, z) \pi^f(dz) < \infty \quad \text{for } x \in X.$$

In [2] we considered continuity-like property of the function $f \mapsto \pi^f$. In [4] we characterized the limit distribution π^f via a functional equation for its characteristic function for some special rv-functions in Hilbert spaces. In the present paper we are strengthening the result of [2], apply it to equation (1) and consider also the equation used for the above mentioned characterization of the limit distribution.

1. Assuming that (Ω, \mathcal{A}, P) is a probability space and (X, ϱ) is a complete and separable metric space, consider the set \mathcal{R}_c of all rv-functions $f: X \times \Omega \rightarrow X$ such that

$$\int_{\Omega} \varrho(f(x, \omega), f(z, \omega)) P(d\omega) \leq \lambda_f \varrho(x, z) \quad \text{for } x, z \in X$$

with a $\lambda_f \in [0, 1)$ and (2) holds. Put also

$$d(f, g) = \sup \left\{ \int_{\Omega} \varrho(f(x, \omega), g(x, \omega)) P(d\omega) : x \in X \right\} \quad \text{for } f, g \in \mathcal{R}_c.$$

The theorem in [2] says that if $f, g \in \mathcal{R}_c$, then

$$(4) \quad \left| \int_X u d\pi^f - \int_X u d\pi^g \right| \leq \frac{1}{1 - \min\{\lambda_f, \lambda_g\}} d(f, g)$$

for every non-expansive $u: X \rightarrow [-1, 1]$. In fact the above inequality was proved there for every non-expansive and bounded $u: X \rightarrow \mathbb{R}$. But if $f \in \mathcal{R}_c$, then (3) holds and so every Lipschitz function mapping X into a separable Banach space is Bochner integrable with respect to π^f . Therefore we can ask whether (4) holds also for such a function. The theorem reads as follows.

THEOREM 1. *If $f, g \in \mathcal{R}_c$, then*

$$(5) \quad \left\| \int_X u d\pi^f - \int_X u d\pi^g \right\| \leq \frac{1}{1 - \min\{\lambda_f, \lambda_g\}} d(f, g)$$

for every non-expansive u mapping X into a separable Banach space.

PROOF. Let u be a non-expansive mapping of X into a separable Banach space Y . To show that (5) holds we may assume that Y is a real space.

Fix $y^* \in Y^*$ such that $\|y^*\| \leq 1$ and

$$(6) \quad \left\| \int_X u d\pi^f - \int_X u d\pi^g \right\| = y^* \left(\int_X u d\pi^f - \int_X u d\pi^g \right).$$

For every $k \in \mathbb{N}$ the function $\tau_k: \mathbb{R} \rightarrow \mathbb{R}$ given by $\tau_k(t) = -k$ for $t \in (-\infty, -k)$, $\tau_k(t) = t$ for $t \in [-k, k]$, $\tau_k(t) = k$ for $t \in (k, \infty)$ is non-expansive and $|\tau_k(t)| \leq |t|$ for $t \in \mathbb{R}$. Consequently, since (4) holds for every non-expansive and bounded $u: X \rightarrow \mathbb{R}$, for every $k \in \mathbb{N}$ we have

$$(7) \quad \left| \int_X \tau_k \circ y^* \circ u d\pi^f - \int_X \tau_k \circ y^* \circ u d\pi^g \right| \leq \frac{1}{1 - \min\{\lambda_f, \lambda_g\}} d(f, g)$$

and

$$|\tau_k(y^*u(z))| \leq \|u(z)\| \quad \text{for } z \in X \text{ and } k \in \mathbb{N}.$$

Hence, applying the Lebesgue dominated convergence theorem and passing with k to the limit in (7) we get

$$\left| \int_X y^* \circ u d\pi^f - \int_X y^* \circ u d\pi^g \right| \leq \frac{1}{1 - \min\{\lambda_f, \lambda_g\}} d(f, g)$$

and (5) follows now from (6). □

The following example shows that both sides of (5) can be equal and non-zero.

EXAMPLE 1. If $\Omega = \{0, 1\}$, $P(\{\omega\}) = 1/2$ for $\omega \in \{0, 1\}$ and $f_\alpha(x, \omega) = (x + \alpha\omega)/2$ for $x \in \mathbb{R}$, $\omega \in \{0, 1\}$ and $\alpha \in (0, \infty)$, then $f_\alpha \in \mathcal{R}_c$ with $\lambda_{f_\alpha} = \frac{1}{2}$ and (see [4, Example 1])

$$\pi^{f_\alpha}(B) = \frac{1}{\alpha} \lambda_1(B \cap [0, \alpha]) \quad \text{for Borel } B \subset \mathbb{R} \text{ and } \alpha \in (0, \infty),$$

where λ_1 denotes the one-dimensional Lebesgue measure. Hence

$$\int_{\mathbb{R}} u d\pi^{f_\alpha} = \frac{1}{\alpha} \int_0^\alpha u(x) dx$$

for every $\alpha \in (0, \infty)$ and Lipschitz $u: \mathbb{R} \rightarrow \mathbb{R}$. In particular,

$$\left| \int_{\mathbb{R}} x \pi^{f_\alpha}(dx) - \int_{\mathbb{R}} x \pi^{f_\beta}(dx) \right| = \frac{1}{2} |\alpha - \beta| = \frac{1}{1 - \min\{\lambda_{f_\alpha}, \lambda_{f_\beta}\}} d(f_\alpha, f_\beta)$$

for $\alpha, \beta \in (0, \infty)$.

Denoting by $\|F\|_L$ the smallest Lipschitz constant for a Lipschitz function F we have the following corollary concerning Lipschitz solutions φ of (1).

COROLLARY 1. *Assume F is a Lipschitz mapping of X into a separable Banach space Y . If $f, g \in \mathcal{R}_c$ and*

$$(8) \quad \frac{\|F\|_L}{1 - \min\{\lambda_f, \lambda_g\}} d(f, g) < \left\| \int_X F d\pi^g \right\|,$$

then equation (1) has no Lipschitz solution $\varphi: X \rightarrow Y$.

PROOF. It follows from Theorem 1 and (8) that

$$\left\| \int_X F d\pi^f - \int_X F d\pi^g \right\| \leq \frac{\|F\|_L}{1 - \min\{\lambda_f, \lambda_g\}} d(f, g) < \left\| \int_X F d\pi^g \right\|,$$

whence $\int_X F d\pi^f \neq 0$ and according to [4, Theorem 2.1] equation (1) has no Lipschitz solution $\varphi: X \rightarrow Y$. \square

The following example shows that under the assumptions of Corollary 1 equation (1) may have a continuous solution. (Cf. [1, Example 4.2].)

EXAMPLE 2. Assume p_1, p_2 are positive reals, $p_2 < \frac{1}{9}$ and $p_1 + p_2 = 1$, reals L_1, L_2 satisfy

$$p_1 L_1^2 < 1, \quad p_1 |L_1| + 3\sqrt{p_2(1 - p_1 L_1^2)} < 1, \quad |L_2| = \sqrt{(1 - p_1 L_1^2)/p_2},$$

and $a \in \mathbb{R} \setminus \{0\}$. Define $F: \mathbb{R} \rightarrow \mathbb{R}$ by

$$F(x) = -ap_2(2L_2x + a).$$

Putting $\Omega = \{1, 2\}$ and $P(\{k\}) = p_k$ for $k \in \{1, 2\}$ consider the rv -functions $f, g: \mathbb{R} \times \Omega \rightarrow \mathbb{R}$ given by

$$f(x, 1) = L_1x, \quad f(x, 2) = L_2x + a, \quad g(x, k) = L_kx \quad \text{for } k \in \{1, 2\}.$$

Then $f, g \in \mathcal{R}_c$ with

$$\lambda_f = \lambda_g = p_1|L_1| + p_2|L_2| = p_1|L_1| + \sqrt{p_2(1 - p_1L_1^2)} < 1,$$

and equation (1) takes the form

$$\varphi(x) = p_1\varphi(L_1x) + p_2\varphi(L_2x + a) + F(x).$$

Since $p_1L_1^2 + p_2L_2^2 = 1$, the function $x \mapsto x^2$, $x \in \mathbb{R}$, solves it. Moreover, $g(0, \omega) = 0$ for $\omega \in \Omega$, whence also $g^n(0, \omega) = 0$ for $\omega \in \Omega^\infty$ and $n \in \mathbb{N}$. Consequently, π^g is the Dirac measure δ_0 and

$$\int_{\mathbb{R}} F d\pi^g = F(0) = -p_2a^2.$$

Finally, $\|F\|_L = 2p_2|aL_2|$, $d(f, g) = p_2|a|$, and so

$$\frac{\|F\|_L}{1 - \min\{\lambda_f, \lambda_g\}} d(f, g) = \frac{2p_2^2a^2|L_2|}{1 - (p_1|L_1| + \sqrt{p_2(1 - p_1L_1^2)})} < p_2a^2 = \left| \int_{\mathbb{R}} F d\pi^g \right|.$$

Consider also a special case of (1), viz.

$$(9) \quad \varphi(x) = \frac{1}{n} \sum_{k=0}^{n-1} \varphi\left(\frac{1}{n}x + a_k\right) + F(x).$$

COROLLARY 2. Assume $n \geq 2$ is an integer and F is a Lipschitz mapping of \mathbb{R} into a separable Banach space Y with $\int_0^1 F(x)dx \neq 0$. If reals a_0, \dots, a_{n-1} satisfy

$$\|F\|_L \sum_{k=0}^{n-1} \left| \frac{k}{n} - a_k \right| < (n-1) \left\| \int_0^1 F(x)dx \right\|,$$

then equation (9) has no Lipschitz solution $\varphi: \mathbb{R} \rightarrow Y$.

PROOF. Put $\Omega = \{0, 1, \dots, n-1\}$, $P(\{k\}) = \frac{1}{n}$ for $k \in \Omega$ and define $f, g: \mathbb{R} \times \Omega \rightarrow \mathbb{R}$ by

$$f(x, k) = \frac{1}{n}x + a_k, \quad g(x, k) = \frac{1}{n}x + \frac{k}{n}.$$

Clearly $f, g \in \mathcal{R}_c$ with $\lambda_f = \lambda_g = \frac{1}{n}$ and (see [4, Example 1]) $\pi^g(B) = \lambda_1(B \cap [0, 1])$ for Borel $B \subset \mathbb{R}$. Moreover,

$$d(f, g) = \frac{1}{n} \sum_{k=0}^{n-1} \left| \frac{k}{n} - a_k \right|.$$

By Corollary 1 equation (9) has no Lipschitz solution $\varphi: \mathbb{R} \rightarrow Y$. □

2. Assuming now that X is a real separable Hilbert space, $X \neq \{0\}$, $\Lambda: X \rightarrow X$ is linear and continuous with $\|\Lambda\| < 1$, and μ is a probability Borel measure on X , consider the equation

$$(10) \quad \varphi(x) = \hat{\mu}(x)\varphi(\Lambda x),$$

where $\hat{\mu}$ denotes the Fourier transform of μ ,

$$\hat{\mu}(x) = \int_X e^{i(x|z)} \mu(dz) \quad \text{for } x \in X.$$

THEOREM 2. *If μ has a finite first moment, then there exists a probability Borel measure ν on X with a finite first moment such that $\hat{\nu}$ solves (10), and for any continuous at zero solution $\varphi: X \rightarrow \mathbb{C}$ of (10) we have*

$$\varphi = \varphi(0)\hat{\nu};$$

in particular, every continuous at zero solution $\varphi: X \rightarrow \mathbb{C}$ of (10) is of class C^1 and Lipschitz.

Remind that a probability Borel measure ν on X has a finite first moment provided the integral $\int_X \|x\| \nu(dx)$ is finite. We shall prove Theorem 2 later on, together with the next one and with the following remark.

REMARK. If μ has a finite first moment and Λ is injective, then for every $c \in \mathbb{C}$ the set of all discontinuous at zero solutions $\varphi: X \rightarrow \mathbb{C}$ of (10) such that $\varphi(0) = c$ and $\varphi|_{X \setminus \{0\}}$ is of class C^1 and Lipschitz has the cardinality of the continuum.

Theorem 2 implies that for every Borel and integrable with respect to μ function $\xi: X \rightarrow X$ the equation

$$(11) \quad \varphi(x) = \varphi(\Lambda x) \int_X e^{i(x|\xi(z))} \mu(dz)$$

has exactly one continuous at zero solution $\varphi^\xi: X \rightarrow \mathbb{C}$ such that $\varphi^\xi(0) = 1$, and it is of class C^1 and Lipschitz. Consequently, we have the operator $\xi \mapsto \varphi^\xi$, $\xi \in L^1(\mu, X)$, and a kind of its continuity gives the following theorem.

THEOREM 3. *If $\xi, \eta: X \rightarrow X$ are Borel and integrable with respect to μ , then*

$$|\varphi^\xi(x) - \varphi^\eta(x)| \leq \frac{\|x\|}{1 - \|\Lambda\|} \int_X \|\xi(z) - \eta(z)\| \mu(dz) \quad \text{for } x \in X.$$

PROOFS. Consider the probability space (X, \mathcal{B}, μ) and, given Borel $\xi: X \rightarrow X$ integrable with respect to μ , the rv-function f on it defined by

$$f(x, \omega) = \Lambda^* x + \xi(\omega) \quad \text{for } (x, \omega) \in X \times X,$$

as well as the limit distribution π^f . Put $\pi_\xi = \pi^f$. According to [4, Theorem 3.1] $\hat{\pi}_\xi$ solves (11). Since the first moment of π_ξ is finite, $\hat{\pi}_\xi$ is of class C^1 and Lipschitz.

To prove Theorem 2 put $\nu = \pi_{id_X}$ and let $\varphi: X \rightarrow \mathbb{C}$ be a continuous at zero solution of (10). Then

$$\varphi(x) = \varphi(\Lambda^n x) \prod_{k=0}^{n-1} \int_X e^{i(\Lambda^k x|z)} \mu(dz) \quad \text{for } n \in \mathbb{N}, x \in X,$$

and $\lim_{n \rightarrow \infty} \Lambda^n x = 0$ for $x \in X$. Since

$$\left| \int_X e^{i(\Lambda^k x|z)} \mu(dz) \right| \leq \int_X |e^{i(\Lambda^k x|z)}| \mu(dz) = 1$$

for $k \in \mathbb{N} \cup \{0\}$ and $x \in X$, it shows that if $\varphi(0) = 0$, then $\varphi = 0$, and if $\varphi(0) \neq 0$, then

$$\varphi(x) = \varphi(0) \prod_{n=0}^{\infty} \int_X e^{i(\Lambda^n x|z)} \mu(dz) \quad \text{for } x \in X.$$

Consequently, for every $c \in \mathbb{C}$ equation (10) has at most one continuous at zero solution $\varphi: X \rightarrow \mathbb{C}$ satisfying $\varphi(0) = c$ and by the first part of the proof $c\hat{\nu}$ is such a solution.

To get Theorem 3 it is enough to observe that since $\varphi^\xi = \hat{\pi}_\xi$, $\varphi^\eta = \hat{\pi}_\eta$ and

$$\left| e^{i(x|z_1)} - e^{i(x|z_2)} \right| \leq \|x\| \|z_1 - z_2\| \quad \text{for } x, z_1, z_2 \in X,$$

by Theorem 1 for every $x \in X$ we have

$$\begin{aligned} |\varphi^\xi(x) - \varphi^\eta(x)| &= \left| \int_X e^{i(x|z)} \pi_\xi(dz) - \int_X e^{i(x|z)} \pi_\eta(dz) \right| \\ &\leq \frac{\|x\|}{1 - \|\Lambda\|} \int_X \|\xi(z) - \eta(z)\| \mu(dz). \end{aligned}$$

To verify the Remark given $c \in \mathbb{C}$ for every $a \in \mathbb{C} \setminus \{c\}$ define $\varphi_a: X \rightarrow \mathbb{C}$ by

$$\varphi_a(x) = a\hat{\nu}(x) \quad \text{for } x \in X \setminus \{0\}, \quad \varphi_a(0) = c,$$

and note that it solves (10), it is discontinuous at zero and $\varphi_a|_{X \setminus \{0\}}$ is of class C^1 and Lipschitz. \square

Acknowledgement. The research was supported by the Institute of Mathematics of the University of Silesia (Iterative Functional Equations and Real Analysis program).

References

- [1] K. Baron, *On the convergence in law of iterates of random-valued functions*, Aust. J. Math. Anal. Appl. **6** (2009), no. 1, Art. 3, 9 pp.
- [2] K. Baron, *On the continuous dependence in a problem of convergence of iterates of random-valued functions*, Grazer Math. Ber. **363** (2015), 1–6.
- [3] K. Baron, *Weak law of large numbers for iterates of random-valued functions*, Aequationes Math. **93** (2019), 415–423.
- [4] K. Baron, *Weak limit of iterates of some random-valued functions and its application*, Aequationes Math. DOI: 10.1007/s00010-019-00650-z.
- [5] R. Kapica, *Sequences of iterates of random-valued vector functions and solutions of related equations*, Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II **213** (2004), 113–118 (2005).
- [6] R. Kapica, *The geometric rate of convergence of random iteration in the Hutchinson distance*, Aequationes Math. **93** (2019), 149–160.
- [7] M. Kuczma, B. Choczewski and R. Ger, *Iterative functional equations*, Encyclopedia of Mathematics and its Applications, vol. 32, Cambridge University Press, Cambridge, 1990.

INSTITUTE OF MATHEMATICS
 UNIVERSITY OF SILESIA
 BANKOWA 14
 40-007 KATOWICE
 POLAND
 e-mail: baron@us.edu.pl