



You have downloaded a document from  
**RE-BUŚ**  
repository of the University of Silesia in Katowice

**Title:** Faint trace of a particle in a noisy Vaidman three-path interferometer

**Author:** Jerzy Dajka

**Citation style:** Dajka Jerzy. (2021). Faint trace of a particle in a noisy Vaidman three-path interferometer. "Scientific Reports" (Vol. 11 (2021), art. no. 1123), doi 10.1038/s41598-020-80806-z



Uznanie autorstwa - Licencja ta pozwala na kopiowanie, zmienianie, rozprowadzanie, przedstawianie i wykonywanie utworu jedynie pod warunkiem oznaczenia autorstwa.



UNIWERSYTET ŚLĄSKI  
W KATOWICACH



Biblioteka  
Uniwersytetu Śląskiego



Ministerstwo Nauki  
i Szkolnictwa Wyższego

OPEN

# Faint trace of a particle in a noisy Vaidman three-path interferometer

Jerzy Dajka


**We study weak traces of particle passing Vaidman’s nested Mach–Zehnder interferometer. We investigate an effect of decoherence caused by an environment coupled to internal degree of freedom (a spin) of a travelling particle. We consider two models: pure decoherence leading to exact results and weak coupling Davies approximation allowing to include dissipative effects. We show that potentially anomalous discontinuity of particle paths survives an effect of decoherence unless it affects internal part of the nested interferometer.**

Quantum particle can be prepared (preselected) in a given and desired state and may also be postselected in another state with a known at least in principle probability. That what occurs in between, what is the particle’s past, remains problematic due to specific features of quantum measurements unavoidably modifying quantum states of measured objects. It is clearly visible for a quantum particle passing through interferometer: the particle enters the device and leaves it (if its outcome is measured), however, what occurs inside an interferometer is, as it will be shown below, disputable and even controversial. One of the approaches<sup>1,2</sup> is that a particle, staying coherent, it leaves nothing but a faint trace which—if comparable to an order of a trace potentially left by a localized wave packet—can serve as a hallmark of its presence. A faint trace which is a *small* change of an amplitude of a component orthogonal to an undisturbed particle’s state (weakly) measurable only in experiments with an ensemble of particles having the same pre- and postselection.

Past of a quantum particle in a nested Mach–Zehnder interferometer—proposed in Ref.<sup>1</sup> and presented in Fig. 1 and abbreviated here as a Vaidman interferometer—was recently studied in Ref.<sup>1</sup> using quantum weak values<sup>3,4</sup> and the two state vector formalism (TSVF)<sup>5</sup>. The results are confounding: particles seem to follow anomalous *discontinuous* path. Such a seemingly weird conclusion leads to plethora of controversies<sup>6,7</sup> and since that time (almost) all works on that problem have come in triads: a paper, commentary inspired by the paper and a reply to the comment<sup>6,7</sup>. The main reason is that the TSVF<sup>5</sup> applied in Ref.<sup>1</sup> is one of few possible approaches to studies of quantum past. The other non-equivalent alternatives are consistent (decoherent) histories<sup>8,9</sup> standard quantum mechanics<sup>10–12</sup> and many other alternative studies<sup>13–20</sup>. Moreover, even recent experiments and their detailed analyses fail to resolve all the controversial issues<sup>20–26,26–28</sup>. Despite counter-intuitive of a conception of discontinuous path there are analyses<sup>2,29</sup> and claims which support the faint-trace anomalous picture as experimentally confirmed.

Our present aim is to follow Ref.<sup>1,2</sup> and to supplement the analysis of weak trace (based upon weak values) of particle by including an effect of decoherence affecting internal degree of freedom of the particle passing the Vaidman interferometer. Our results allow to identify natural obstructions for an effective verification of theoretical predictions and exclude factors seemingly but not truly responsible for experimental failures and limitations. Studying internal degrees of freedom e.g. spin or polarization of particles in Vaidman interferometer becomes particularly reasonable for most recent experiments and models utilizing neutrons<sup>24,26</sup>. There are various phenomenologic approaches<sup>30</sup> dedicated to particular quantum systems which usefulness and validity was confirmed by many repeatable experiments. However, for fragile quantum systems microscopic models are at least a good starting point to make predictions which are credible in a tailored and well identified conditions. In this paper we consider two well established microscopic models: (i) an exact model of pure decoherence<sup>31,32</sup> of a solely quantum character and (ii) a weak coupling Davies approximation<sup>33</sup> allowing to include dissipation. Pure decoherence (or pure dephasing) model is limited by a choice of system–environment interaction encoded in a Hamiltonian by an integral of motion: in the Caldeira–Leggett<sup>30</sup> “system + bath + interaction” Hamiltonian the interaction is given by an operator commuting with a system. Davies approach<sup>33</sup> allows for an arbitrary system–environment coupling provided that it is small enough for a perturbative treatment to hold true. We show that the anomalous discontinuity<sup>1,2</sup> of the faint traces (given by weak values of suitable projection operators) left by particles are rigid with respect to decoherence affecting external arm of Vaidman interferometer while very fragile if a source of decoherence disturbs balance of the internal, nested arms of Vaidman interferometer.

<sup>1</sup>Institute of Computer Science, University of Silesia in Katowice, 40-007 Katowice, Poland. <sup>2</sup>Institute of Physics, University of Silesia in Katowice, 40-007 Katowice, Poland. <sup>3</sup>Silesian Center for Education and Interdisciplinary Research, University of Silesia in Katowice, 41-500 Chorzów, Poland. <sup>email:</sup> jerzy.dajka@us.edu.pl



**Figure 1.** Vaidman interferometer consisting of two Mach-Zehnder nested devices with four beam splitters  $BS_i, i = 1, \dots, 4$ . Roman numbers I, II, III correspond to vectors in Eq. (1) and A, B, C denote instants where the weak trace is calculated.

Since an experiment is an only way to resolve an (interpretative) ambiguity concerning (un)presence of particles in the Vaidman interferometer the answer if the controversies survive also in the presence of decoherence seems to be crucial.

For a sake of completeness we re-introduce the Vaidman interferometer and review the controversial features of the faint traces of particles passing it. The original Vaidman interferometer presented in Fig. 1 consists of spatial degree of freedom given by three paths denoted by I, II, III and four beam splitters. Vaidman interferometer can effectively be described<sup>10–12</sup> as a three level quantum system with a state space spanned by a basis

$$|I\rangle = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, |II\rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \text{ and } |III\rangle = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad (1)$$

In an ideal setting of a noise-less system, a passage of a particle is described by a unitary transformation composed of four unitaries  $U_4 U_3 U_2 U_1$  corresponding to subsequent beam splitters termed as  $BS_i, i = 1, \dots, 4$  in Fig. 1:

$$U_1 = U_4 = \frac{1}{\sqrt{3}} \begin{pmatrix} \sqrt{3} & 0 & 0 \\ 0 & -1 & \sqrt{2} \\ 0 & \sqrt{2} & 1 \end{pmatrix} \text{ and } U_2 = U_3 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & \sqrt{2} \end{pmatrix} \quad (2)$$

The strategy applied in Ref.<sup>1</sup> to infer the path of a particle entering and leaving Vaidman interferometer in a state  $|III\rangle$  was to investigate *weak trace* of a particle inside Vaidman interferometer at three instants A, B, C indicated in Fig. 1. According to Ref.<sup>1</sup> the weak trace is indicated by a non-vanishing weak value<sup>3,34</sup> of one of the projectors

$$\langle \Pi_i \rangle_w^q = \frac{\langle \psi_{post}^q | \Pi_i | \psi_{pre}^q \rangle}{\langle \psi_{post}^q | \psi_{pre}^q \rangle}, \text{ where } \Pi_i = |i\rangle\langle i|, i = I, II, III \text{ and } q = A, B, C \quad (3)$$

where preselected (directly prior to the measurement of  $\Pi_i$ ) and postselected (immediately after the measurement) states compose a *two-state vector*  $\langle \psi_{post} | | \psi_{pre} \rangle$  being a fundamental object of the TSVF<sup>5</sup>.

Let us emphasise that the physical meaning of vanishing weak values in a current context remains disputable<sup>34–37</sup>. Most of controversies originate, however, from highly counter-intuitive conclusions provided in Ref.<sup>1</sup> indicating possibility of discontinuous trajectories followed by a particle passing through Vaidman interferometer. Let us review: there are three instants A, B, C where the weak trace is measured: A: just after it is injected to the Vaidman interferometer in a state  $|III\rangle$  and passes  $BS_1$ , B: where the weak measurement becomes conducted for all potential paths in Vaidman interferometer and C: after the  $BS_3$  beam splitter as presented in Fig. 1. The corresponding preselected states read as:  $|\psi_{pre}^A\rangle = U_1|III\rangle$ ,  $|\psi_{pre}^B\rangle = U_2 U_1|III\rangle$  and  $|\psi_{pre}^C\rangle = U_3 U_2 U_1|III\rangle$ .

$(\Pi_{I,II,III})_w^{A,B,C}$	I	II	III	$U_{pre}^{A,B,C}$	$U_{post}^{A,B,C}$
A	0	0	1	$U_1$	$U_2^\dagger U_3^\dagger U_4^\dagger$
B	-1	1	1	$U_2 U_1$	$U_3^\dagger U_4^\dagger$
C	0	0	1	$U_3 U_2 U_1$	$U_4^\dagger$

**Table 1.** Weak traces  $\langle \Pi_{I,II,III} \rangle_w^{A,B,C}$  Eq. (3) of a particle in a noise-less Vaidman interferometer at different instants A, B, C indicated in Fig. 1 and corresponding pre-and postselections given by  $|\psi_{pre}^{A,B,C}\rangle = U_{pre}^{A,B,C} |III\rangle$  and  $|\psi_{post}^{A,B,C}\rangle = U_{post}^{A,B,C} |III\rangle$  respectively.

In the time-symmetric TSVF setting the postselected states  $|\psi_{post}^A\rangle = U_2^\dagger U_3^\dagger U_4^\dagger |III\rangle$ ,  $|\psi_{post}^B\rangle = U_3^\dagger U_4^\dagger |III\rangle$  and  $|\psi_{post}^C\rangle = U_4^\dagger |III\rangle$  describe a hypothetical particle detected at III evolving backward in time. Results of the weak measurement are summarized in Table 1. According to Ref.<sup>1</sup> a presence of a particle is defined by its non-vanishing weak trace. The counter-intuitive conclusion of Table 1 is the following: at A and C the particle is present in III, what upon Fig. 1 is intuitively acceptable, but at B it is also present in an internal loop (I, II) of Vaidman interferometer. Such confounding result needs experimental verification. One can safely assume that any potential experiment, as it was so far, will be highly subtle and sophisticated. Moreover, such an experiment will rely on quantum properties and it may be fragile with respect to decoherence. Our objective is to investigate if in a presence of omnipresent noise one can still support claims inferred from Table 1 or if they are nothing but an artifact absent in real noisy systems.

### Pure decoherence

There are circumstances when internal degrees of freedom of particles need to be taken into account and affect an interference<sup>38</sup>. An output pattern of such an interference becomes further significantly modified by external bath coupling to an internal degree of freedom<sup>39</sup>. In the following for simplicity we assume that an internal degree of freedom of a particle in the Vaidman interferometer requires two-dimensional space  $\mathcal{H}_{int}$  spanned by

$$|+\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, |-\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (4)$$

Physically, such a qubit model can correspond to an interference of spin-half particles or any qubits. If, moreover, one assumes the internal degree of freedom is coupled to (affected by) an environment one arrives at a composite quantum system with a state space

$$\mathcal{H} = \mathcal{H}_{path} \otimes \mathcal{H}_{int} \otimes \mathcal{H}_{env} \quad (5)$$

where  $\mathcal{H}_{int} = \mathbf{C}^2$  and  $\mathcal{H}_{path} = \mathbf{C}^3$  with a basis  $|I\rangle, |II\rangle, |III\rangle$  in Eq. (1). As a composite system consisting of a particle and its environment considered *in toto* is closed, its evolving states undergo unitary transformations. Unitaries corresponding to beam splitters Eq. (2) and projectors Eq. (3) required for weak measurement become now tensorized with identity operators  $\mathbb{I}$  acting on remaining parts of a composite space  $\mathcal{H}$  and read:

$$U_i = U_i \otimes \mathbb{I}_2 \otimes \mathbb{I}_{env} : \mathcal{H} \rightarrow \mathcal{H}, \quad i = 1, \dots, 4 \quad (6)$$

$$\tilde{\Pi}_i = \Pi_i \otimes \mathbb{I}_2 \otimes \mathbb{I}_{env} : \mathcal{H} \rightarrow \mathcal{H}, \quad i = I, II, III \quad (7)$$

where  $U_i$  and  $\Pi_i$  are given in Eqs. (2) and (3) respectively.

Let us notice that an effect of a generic interaction between particle and its environment shall result in a modification of  $U_i$  which will essentially be interaction-type-dependent. Pure decoherence (dephasing)<sup>30,31</sup> is probably the simplest class of an interaction between an open quantum system and its environment. It is characterized by a high symmetry preventing energy exchange with a surrounding<sup>31</sup>. Despite such an obvious simplification pure decoherence can effectively be applied to a broad class of problems<sup>40-47</sup> ranging from theoretical quantum information up to experiments in optics. Simplifying our model even further we assume *local* decoherence i.e. the particle remains unaffected by a dephasing environment unless it follows a particular 'noisy' path between two particular beam splitters. The term 'local' is used to distinguish circumstances when a whole Vaidman interferometer, not just a part of one of its arms, is embedded in an either thermal or non-thermal bath. Hamiltonian describing interaction between particle (which is a qubit with a Hamiltonian given by  $\sigma_z$  Pauli matrix) and its environment is then given by a standard Caldeira-Leggett form<sup>30</sup>:

$$H_i = E\mathbb{I}_3 \otimes \sigma_z \otimes \mathbb{I}_{env} + \Pi_i \otimes \sigma_z \otimes \int_0^\infty d\omega (g(\omega)a(\omega) + h.c.) + \mathbb{I}_3 \otimes \mathbb{I}_2 \otimes \int_0^\infty d\omega h(\omega)a^\dagger(\omega)a(\omega), \quad i = I, II, III \quad (8)$$

Time evolution of a total (closed) system is unitary and reads

$$U_i = \exp(-itH_i) = |i\rangle\langle i| \otimes \begin{pmatrix} U_+ & 0 \\ 0 & U_- \end{pmatrix} + (\mathbb{I}_3 - |i\rangle\langle i|) \otimes \mathbb{I}_2 \otimes \mathbb{I}_{env}, \quad i = I, II, III \quad (9)$$

$\langle \tilde{\Pi}_{I,II,III} \rangle_w^{A,B,C}$	I	II	III	$U_{pre}^{A,B,C}$	$U_{post}^{A,B,C}$	$U_{pre}^{A,B,C}$	$U_{post}^{A,B,C}$
A	0	0	1	$U_{II} U_I$	$U_2^\dagger U_3^\dagger U_4^\dagger$	$U_{III} U_I$	$U_2^\dagger U_3^\dagger U_4^\dagger$
B	-q/2	q/2	1	$U_2 U_{II} U_I$	$U_3^\dagger U_4^\dagger$	$U_2 U_{III} U_I$	$U_3^\dagger U_4^\dagger$
C	0	0	1	$U_3 U_2 U_{II} U_I$	$U_4^\dagger$	$U_3 U_2 U_{III} U_I$	$U_4^\dagger$

**Table 2.** Weak traces  $\langle \tilde{\Pi}_{I,II,III} \rangle_w^{A,B,C}$  Eq. (7) of a particle in a noisy Vaidman interferometer at different instants  $A, B, C$  indicated in Fig. 1 and corresponding pre-and postselections given by  $\rho_{pre}^{A,B,C} = U_{pre}^{A,B,C} (|\psi_0\rangle\langle\psi_0| \otimes |\Omega\rangle\langle\Omega|) [U_{pre}^{A,B,C}]^\dagger$  and  $\rho_{post}^{A,B,C} = U_{post}^{A,B,C} (\rho_{out} \otimes |\Omega\rangle\langle\Omega|) [U_{post}^{A,B,C}]^\dagger$  respectively. The quantity  $q = \langle\Omega|U_+\Omega\rangle + \langle\Omega|U_-\Omega\rangle$  is given in Eq. (15) with  $U_\pm$  given in Eq. (10). The effect of pure decoherence indicated by  $U_{II}$  occurs for path II (by  $U_{III}$  for path III) after beam splitter  $BS_1$ .

$\langle \tilde{\Pi}_{I,II,III} \rangle_w^{B,C}$	I	II	III	$U_{pre}^{B,C}$	$U_{post}^{B,C}$	$U_{pre}^{B,C}$	$U_{post}^{B,C}$
B	-(q+2)/4	(q+2)/4	1	$U_I U_2 U_I$	$U_3^\dagger U_4^\dagger$	$U_{II} U_2 U_I$	$U_3^\dagger U_4^\dagger$
C	0	(2-q)/2	q/2	$U_3 U_I U_2 U_I$	$U_4^\dagger$	$U_3 U_{II} U_2 U_I$	$U_4^\dagger$

**Table 3.** Weak traces  $\langle \tilde{\Pi}_{I,II,III} \rangle_w^{B,C}$  Eq. (7) of a particle in a noisy Vaidman interferometer at different instants  $B, C$  indicated in Fig. 1 and corresponding pre-and postselections given by  $\rho_{pre}^{B,C} = U_{pre}^{B,C} (|\psi_0\rangle\langle\psi_0| \otimes |\Omega\rangle\langle\Omega|) [U_{pre}^{B,C}]^\dagger$  and  $\rho_{post}^{B,C} = U_{post}^{B,C} (\rho_{out} \otimes |\Omega\rangle\langle\Omega|) [U_{post}^{B,C}]^\dagger$  respectively. The quantity  $q = \langle\Omega|U_+\Omega\rangle + \langle\Omega|U_-\Omega\rangle$  with  $U_\pm$  given in Eq. (10). The effect of pure decoherence indicated by  $U_I$  occurs for path I after beam splitter  $BS_2$ . The effect of pure decoherence indicated by  $U_{II}$  occurs for path II after beam splitter  $BS_2$ .

$$U_\pm = \exp(-itH_\pm) : \mathcal{H}_{env} \rightarrow \mathcal{H}_{env} \quad (10)$$

where  $t$  denotes duration of particle–bath interaction which is assumed to be smaller than the passage between any pair of beam splitters. A block–diagonal structure of the first term in Eq. (9) with unitary blocks  $U_\pm$  given by Eq. (10) is a hallmark of the assumed pure decoherence model with

$$H_\pm = \int_0^\infty d\omega [h(\omega)a^\dagger(\omega)a(\omega) \pm (g(\omega)a(\omega) + h.c.)] \pm E \quad (11)$$

where  $E$  denotes energy separating qubit levels, the environment is simplified to a one-dimensional bosonic field with bosonic operators  $a(\omega)$ ,  $a^\dagger(\omega)$  and real-valued  $h(\omega)$  and  $g(\omega)$ . To clarify the notation of Eq. (9) let us exemplify: a unitary transformation  $U_4 U_3 U_2 U_{II} U_I : \mathcal{H} \rightarrow \mathcal{H}$  describes Vaidman interferometer with a purely dephasing environment coupled to a path II locally between beam splitters  $BS_1$  and  $BS_2$ .

Now we are ready to analyse an impact of decoherence on particles travelling via Vaidman interferometer. We recognize two classes of effects: the first, quantitative when the anomalous effect of Ref.<sup>1</sup> survives and the second when the pure decoherence spoils unusual features of noise-less system. The first case is presented in Table 2. We consider quantum particle entering Vaidman interferometer in a state

$$|\psi_0\rangle = |III\rangle \otimes [|+\rangle + |-\rangle] / \sqrt{2} \quad (12)$$

where the first, spatial, component denotes path of the particle whereas the second is a state of internal degree of freedom (qubit). The particle leaving the Vaidman interferometer is assumed to be in a state

$$\rho_{out} = |III\rangle\langle III| \otimes \frac{1}{2} [|+\rangle\langle +| + |-\rangle\langle -|] = |III\rangle\langle III| \otimes \frac{1}{2} \mathbb{I}_2 \quad (13)$$

i.e. it is postselected in its spatial (external) degree of freedom but with no information regarding its internal (here maximally mixed) state. Moreover, we assume that an environment both initially and finally is its ground state (vacuum)  $|\Omega\rangle$  and affects only these particles which, according to TSVF, travel forward in time. Working essentially with mixed states requires generalisation of a definition of a weak value of an operators<sup>34</sup>. In particular for a faint trace one calculates

$$\langle \Pi_{I,II,III} \otimes \mathbb{I}_2 \rangle_w = \frac{\text{Tr}(\rho_{post} [\Pi_{I,II,III} \otimes \mathbb{I}_2] \rho_{pre})}{\text{Tr}(\rho_{pre} \rho_{post})} \quad (14)$$

for generically mixed pre-and postselections. Depending on the instants  $A, B, C$ , cf. Fig. 1 of a weak measurement, we consider three different pre-and postselection  $\rho_{pre}^{A,B,C} = U_{pre}^{A,B,C} (|\psi_0\rangle\langle\psi_0| \otimes |\Omega\rangle\langle\Omega|) [U_{pre}^{A,B,C}]^\dagger$  and  $\rho_{post}^{A,B,C} = U_{post}^{A,B,C} (\rho_{out} \otimes |\Omega\rangle\langle\Omega|) [U_{post}^{A,B,C}]^\dagger$  respectively with  $U_{pre,post}^{A,B,C}$  given explicitly in Tables 2, 3. The results of

weak measurement of a particle disturbed by decoherence coupled either to II or III are summarized in Table 2. Let us notice that decoherence affects particle traveling forward in time after it passes the beam splitter  $BS_1$ .

Modifications of weak values due to decoherence reported in Tables 2 and 3 are qualified by

$$q = \langle \Omega | U_+ \Omega \rangle + \langle \Omega | U_- \Omega \rangle \quad (15)$$

which for pure decoherence has an exact solution<sup>30–32</sup>. Since expressed in terms of displacement operators  $D$ <sup>48</sup> time evolution of a vacuum  $|\Omega\rangle \in \mathcal{H}_{env}$ <sup>39</sup> reads

$$U_{\pm}|\Omega\rangle = \exp(\mp iEt)D\left(\pm \frac{g(\omega)}{h(\omega)}(1 - e^{-ih(\omega)t})\right)|\Omega\rangle, \quad D(f) = \exp\left[\int_0^\infty d\omega(f(\omega)a^\dagger(\omega) - h.c.)\right] \quad (16)$$

for a typical choice  $h(\omega) = \omega$  and  $g^2(\omega) = \lambda\omega^{1+\mu} \exp(-\omega/\omega_c)$ <sup>30</sup> one can calculate explicitly

$$q = 2 \cos(Et) \exp\left(-\int_0^\infty d\omega \frac{g^2(\omega)}{h^2(\omega)}(1 - \cos(h(\omega)t))\right) = 2 \cos(Et) \exp\left(-\lambda\Gamma(\mu)\omega_c^\mu \frac{1 - \cos(\mu \arctan(\omega_c t))}{(1 + \omega_c^2 t^2)^{\mu/2}}\right) \quad (17)$$

where the parameter  $\lambda$  denotes strength of a particle–environment coupling and  $0 \leq \mu$  allows to classify environment<sup>30</sup> as Ohmic  $\mu = 0$  or super-Ohmic for  $\mu > 0$ . The sub-Ohmic regime  $\mu < 0$  suffers from known<sup>31</sup> mathematical difficulties and is not considered. Let us notice that for  $t = 0$  the integrand in Eq. (17) vanishes,  $q = 2$  and the results of Ref.<sup>1</sup> are reproduced. It holds also true for  $\lambda \rightarrow 0$  corresponding to a particle *uncoupled* with its environment. We also conclude that even in a long time limit  $t \rightarrow \infty$  the quantity  $q > 0$  i.e. it remains finite. Therefore we infer that discontinuous faint trace changes only quantitatively. However, there is a qualitative change if decoherence is present in an *internal* interferometer of Vaidman interferometer as presented in Table 3 i.e. if an environment is coupled either to I or II after beam splitter  $BS_2$ . In both cases, still assuming that particles travelling forward in time are affected by decoherence, we observe that a faint trace of a particle contributes to II after  $BS_3$ . This observation supports claim of Ref.<sup>10</sup> connecting the faint trace discontinuity with a perfect balance of internal interferometer leading to destructive interference on its output. If decoherence affects internal loop in the Vaidman interferometer it removes anomaly of a faint trace. To summarize, the faint trace, otherwise fragile, remains resistant with respect to decoherence present in an external arm of Vaidman interferometer.

## Dissipation

Rigidity of faint traces with respect to decoherence affecting external interferometer of Vaidman interferometer accompanied by its fragility with respect to decoherence present in an internal loop was analysed in previous paragraph for a very special model of pure decoherence. Here we investigate if the above predictions survive also in a presence of dissipation i.e. particle–environment realistic interaction not excluding an energy exchange. Such a problem, contrary to exactly solvable pure decoherence, demands approximate treatment<sup>30</sup>. We apply Davies approach<sup>33</sup>, one of rigorous approaches to quantum open systems dedicated to weak coupling to an environment<sup>30</sup>. Davies treatment, mathematically rigorous, can effectively be applied in various areas of quantum information<sup>49–52</sup>.

Let us keep previous assumption of decoherence affecting locally an external arm of Vaidman interferometer only and consider particle–environment interaction encoded in one of two following Hamiltonians


$$H_i = E\mathbb{I}_3 \otimes \sigma_z \otimes \mathbb{I}_{env} + \varepsilon\Pi_i \otimes \sigma_x \otimes \int_0^\infty d\omega(g(\omega)a(\omega) + h.c.) + \mathbb{I}_3 \otimes \mathbb{I}_2 \otimes \int_0^\infty d\omega h(\omega)a^\dagger(\omega)a(\omega), \quad i = \text{II, III} \quad (18)$$

Here  $\varepsilon$  is assumed to be small. The choice of coupling (via  $\sigma_x$  Pauli matrix) is complementary to the pure decoherence model studied in the previous Section and allows us to verify if previously predicted stability of faint trace holds also true disturbed by dissipation. None of two Hamiltonians in Eq. (18) allows for an exact treatment similar to pure decoherence. Instead we assume vanishing temperature limit  $T = 0$  and utilize Davies approximation to find time evolution of reduced (with respect to an environment) density matrix of a particle  $\rho(t) \in \mathcal{B}(\mathcal{H}_{path} \otimes \mathcal{H}_{int})$  in terms of a Master equation

$$\frac{d}{dt}\rho(t) = -i \left[ \left( E\mathbb{I}_3 \otimes \sigma_z + \sum_{k,l=1}^6 s(\Omega_{kl})A_{kl}^{(i)\dagger}A_{kl}^{(i)} \right), \rho(t) \right] + \frac{1}{2} \sum_{k,l=1}^6 c(\Omega_{kl})([A_{kl}^{(i)}\rho(t), A_{kl}^{(i)\dagger}] + [A_{kl}, \rho(t)A_{kl}^{(i)\dagger}]) \quad (19)$$

with  $A_{kl}^{(i)} = P_k(\varepsilon\Pi_i \otimes \sigma_x)P_l$ ,  $i = \text{II, III}$ ,  $c(x) = 2\pi x \exp(-x/\omega_c)\Theta(x)$  and numerically  $2\pi s(x) = \text{PV} \int_{-\infty}^\infty dy[c(y)/(y-x)]$ . For any initial density matrix  $\rho_0$  any solution of Eq. (19) is a one-parameter completely positive quantum dynamical semi-group<sup>53</sup>  $\rho(t) = \mathcal{D}[\rho_0]$ .

We investigate weak traces of a particle in Vaidman interferometer assuming that there is dissipative environment affecting particle just after  $BS_1$  and it is attached to the external interferometer in the Vaidman interferometer in Fig. 1. To calculate  $\langle \Pi_{\text{I,II,III}} \otimes \mathbb{I}_2 \rangle_w$  at instants  $A, B, C$ , cf. Fig. 1, for a particle injected to the Vaidman interferometer in a state  $|\psi_0\rangle = |\text{III}\rangle \otimes [ |+\rangle + |-\rangle ]/\sqrt{2} \in \mathcal{H}_{path} \otimes \mathcal{H}_{int}$  Eq. (12) we consider the following pre- and postselections:



**Figure 2.** Weak values of  $(\Pi_{I,II,III} \otimes \mathbb{I}_2)_w$  Eq. (14) *upper panel* at instants A and C, cf. Fig. 1, for a pre- and a postselection given in Eqs. (20) and (22) respectively and, *lower panel*, at B for a pre- and postselection given in Eq. (21) as a function of duration of particle-environment interaction. Time is given in  $1/\omega_c$  and we set  $\varepsilon = 0.05$ .

$$\rho_{pre}^A = \mathcal{D} [U_1 \otimes \mathbb{I}_2 |\psi_0\rangle \langle \psi_0| U_1^\dagger \otimes \mathbb{I}_2], \quad \rho_{post}^A = U_2^\dagger \otimes \mathbb{I}_2 \rho_{post}^B U_2 \otimes \mathbb{I}_2 \quad (20)$$

$$\rho_{pre}^B = U_2 \otimes \mathbb{I}_2 \rho_{pre}^A U_2^\dagger \otimes \mathbb{I}_2, \quad \rho_{post}^B = U_3^\dagger \otimes \mathbb{I}_2 \rho_{post}^C U_3 \otimes \mathbb{I}_2 \quad (21)$$



$$\rho_{pre}^C = U_3 \otimes \mathbb{I}_2 \rho_{pre}^B U_3^\dagger \otimes \mathbb{I}_2, \quad \rho_{post}^C = U_4^\dagger \otimes \mathbb{I}_2 \rho_{out} U_4 \otimes \mathbb{I}_2 \quad (22)$$

with  $\rho_{out}$  given in Eq. (13) and hence we avoid discussing decohering states evolving backward in time needed for TSVF<sup>34</sup>. Weak values Eq. (14) indicating faint trace of a particle in Vaidman interferometer at A, B, C in Fig. 1 are calculated numerically and presented in Fig. 2 as a function of duration (time) of interaction between the particle and its environment which is coupled to lower (III) arm of the external interferometer in the Vaidman interferometer in Fig. 1 after  $BS_1$ . Let us notice that none of the calculated weak values has a non-vanishing imaginary part requiring careful physical interpretation<sup>4</sup>. Weak trace calculated at A, C (B) is presented in the upper (lower) panel of Fig. 2 respectively. The weak trace indicating particle at A and C is present for the path III and absent otherwise. At B the particle leaves weak trace in all the three paths provided that the duration (time) of dissipation is short with respect to a passage time. If it is not the case one obtains oscillations which, for certain value of duration time lead to decreasing of weak traces at I and II to a level which may be undetectable and, effectively but not factually, the particle remains 'visible' in III only. For an environment weakly disturbing path II after  $BS_1$  one obtains qualitatively similar results.

## Discussion

Decoherence is a trespasser of failure of many experiments attempting to predict or confirm quantum properties of Nature. Recent predictions of discontinuous path of particle in a nested Mach–Zehnder interferometer can serve as a particular example of a deeply quantum effect requiring further experimental verification. One could have doubt in precise measurability of the controversial and to some extent exotic properties of Ref.<sup>1</sup> and in particular Ref.<sup>2</sup> due to omnipresent noise blurring results of measurements. To dispel such doubts we investigated how decoherence can affect theoretical predictions of noise-less models and if it can obscure or even definitely indisposed theoretically predicted anomalies. Since recent experiments<sup>24,26</sup> utilized neutrons which are particles with internal, spin degree of freedom we consider interference of qubits and decoherence affecting its spin. From a wide spectrum of different models describing quantum open systems we select two limiting cases: (i) an exact, non-Markov but specific pure dephasing model and (ii) a very general but approximate weak coupling Davies approximation. Pure decoherence, being exact, does not take into account dissipation of energy how the Davies approach does but at a cost of applied approximations. However, the results obtained for this seemingly far models were confluent: qualitative properties of weak traces (and their discontinuity) of a particle in the Vaidman interferometer are rigid with respect to decoherence affecting external interferometer but at the same time extraordinarily fragile if decoherence is present in an internal loop of Vaidman interferometer provided that duration of decoherence remain short with respect to overall time scales of particle motion in the Vaidman interferometer. We hope that our results, although only qualitative, can serve as a guidelines for experiments and support further investigations concerning past of quantum particles.

## Methods

For analytical calculation of pure decoherence model we utilized coherent state techniques. Numerical calculation for dissipative environment in Davies approximation we performed with Python-based toolbox QuTip<sup>54,55</sup> using `mesolve` for Master Equation Eq. (19).

Received: 2 October 2020; Accepted: 28 December 2020

Published online: 13 January 2021

## References

- Vaidman, L. Past of a quantum particle. *Phys. Rev. A* **87**, 052104. <https://doi.org/10.1103/PhysRevA.87.052104> (2013).
- Vaidman, L. Neutrons and photons inside a nested Mach–Zehnder interferometer. *Phys. Rev. A* **101**, 052119. <https://doi.org/10.1103/PhysRevA.101.052119> (2020).
- Aharonov, Y., Albert, D. Z. & Vaidman, L. How the result of a measurement of a component of the spin of a spin-1/2 particle can turn out to be 100. *Phys. Rev. Lett.* **60**, 1351–1354. <https://doi.org/10.1103/PhysRevLett.60.1351> (1988).
- Dressel, J., Malik, M., Miatto, F. M., Jordan, A. N. & Boyd, R. W. Colloquium. *Rev. Mod. Phys.* **86**, 307–316. <https://doi.org/10.1103/RevModPhys.86.307> (2014).
- Aharonov, Y. & Vaidman, L. *The Two-State Vector Formalism: An Updated Review* 399–447 (Springer, Berlin, 2008).
- Li, Z.-H., Al-Amri, M. & Zubairy, M. S. Comment on “past of a quantum particle”. *Phys. Rev. A* **88**, 046102. <https://doi.org/10.1103/PhysRevA.88.046102> (2013).
- Vaidman, L. Reply to “comment on ‘past of a quantum particle’”. *Phys. Rev. A* **88**, 046103. <https://doi.org/10.1103/PhysRevA.88.046103> (2013).
- Griffiths, R. B. Particle path through a nested Mach–Zehnder interferometer. *Phys. Rev. A* **94**, 032115. <https://doi.org/10.1103/PhysRevA.94.032115> (2016).
- Vaidman, L. Comment on “particle path through a nested Mach–Zehnder interferometer”. *Phys. Rev. A* **95**, 066101. <https://doi.org/10.1103/PhysRevA.95.066101> (2017).
- Englert, B.-G., Horia, K., Dai, J., Len, Y. L. & Ng, H. K. Past of a quantum particle revisited. *Phys. Rev. A* **96**, 022126. <https://doi.org/10.1103/PhysRevA.96.022126> (2017).
- Peleg, U. & Vaidman, L. Comment on “past of a quantum particle revisited”. *Phys. Rev. A* **99**, 026103. <https://doi.org/10.1103/PhysRevA.99.026103> (2019).
- Englert, B.-G., Horia, K., Dai, J., Len, Y. L. & Ng, H. K. Reply to “comment on ‘past of a quantum particle revisited’”. *Phys. Rev. A* **99**, 026104. <https://doi.org/10.1103/PhysRevA.99.026104> (2019).
- Bartkiewicz, K. *et al.* One-state vector formalism for the evolution of a quantum state through nested Mach–Zehnder interferometers. *Phys. Rev. A* **91**, 012103. <https://doi.org/10.1103/PhysRevA.91.012103> (2015).
- Vaidman, L. Comment on “one-state vector formalism for the evolution of a quantum state through nested Mach–Zehnder interferometers”. *Phys. Rev. A* **93**, 036103. <https://doi.org/10.1103/PhysRevA.93.036103> (2016).
- Potoček, V. & Ferenczi, G. Which-way information in a nested Mach–Zehnder interferometer. *Phys. Rev. A* **92**, 023829. <https://doi.org/10.1103/PhysRevA.92.023829> (2015).



16. Vaidman, L. Comment on “which-way information in a nested Mach–Zehnder interferometer”. *Phys. Rev. A* **93**, 017801. <https://doi.org/10.1103/PhysRevA.93.017801> (2016).
17. Hashmi, F. A., Li, F., Zhu, S.-Y. & Zubairy, M. S. Two-state vector formalism and quantum interference. *J Phys A Math. Theor.* **49**, 345302. <https://doi.org/10.1088/1751-8113/49/34/345302> (2016).
18. Vaidman, L. Comment on ‘two-state vector formalism and quantum interference’. *J Phys. A Math. Theor.* **51**, 068002. <https://doi.org/10.1088/1751-8121/aa8d24> (2018).
19. Hashmi, F. A., Li, F., Zhu, S.-Y. & Zubairy, M. S. Reply to the comment on ‘two-state vector formalism and quantum interference’. *J Phys. A Math. Theor.* **51**, 068001. <https://doi.org/10.1088/1751-8121/aa8d23> (2018).
20. Aharonov, Y., Cohen, E., Landau, A. & Elitzur, A. C. The case of the disappearing (and re-appearing) particle. *Sci. Rep.* **7**, 531. <https://doi.org/10.1038/s41598-017-00274-w> (2017).
21. Danan, A., Farfurnik, D., Bar-Ad, S. & Vaidman, L. Asking photons where they have been. *Phys. Rev. Lett.* **111**, 240402. <https://doi.org/10.1103/PhysRevLett.111.240402> (2013).
22. Salih, H. Commentary: “asking photons where they have been”—without telling them what to say. *Front. Phys.* **3**, 47. <https://doi.org/10.3389/fphy.2015.00047> (2015).
23. Vaidman, L., Danan, A., Farfurnik, D. & Bar-Ad, S. Response: Commentary: “asking photons where they have been”—without telling them what to say. *Front. Phys.* **3**, 48. <https://doi.org/10.3389/fphy.2015.00048> (2015).
24. Sponar, S., Geppert, H., Denkmayr, T., Lemmel, H. & Hasegawa, Y. Asking neutrons where they have been. *J. Phys. Conf. Ser.* **1316**, 012002. <https://doi.org/10.1088/1742-6596/1316/1/012002> (2019).
25. Zhou, Z.-Q. *et al.* Experimental observation of anomalous trajectories of single photons. *Phys. Rev. A* **95**, 042121. <https://doi.org/10.1103/PhysRevA.95.042121> (2017).
26. Geppert-Kleinrath, H. *et al.* Multifold paths of neutrons in the three-beam interferometer detected by a tiny energy kick. *Phys. Rev. A* **97**, 052111. <https://doi.org/10.1103/PhysRevA.97.052111> (2018).
27. Saldanha, P. L. Interpreting a nested Mach–Zehnder interferometer with classical optics. *Phys. Rev. A* **89**, 033825. <https://doi.org/10.1103/PhysRevA.89.033825> (2014).
28. Wieśniak, M. Spectra in nested Mach–Zehnder interferometer experiments. *Phys. Lett. A* **382**, 2565–2568. <https://doi.org/10.1016/j.physleta.2018.06.032> (2018).
29. Aharonov, Y., Cohen, E., Waegell, M. & Elitzur, A. C. The weak reality that makes quantum phenomena more natural: Novel insights and experiments. *Entropy*. <https://doi.org/10.3390/e20110854> (2018).
30. Breuer, H.-P. & Petruccione, F. *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2003).
31. Alicki, R. Pure decoherence in quantum systems. *Open Syst. Inf. Dyn.* **11**, 53. <https://doi.org/10.1023/B:OPSY.0000024755.58888.ac> (2004).
32. Łuczka, J. Spin in contact with thermostat: Exact reduced dynamics. *Phys. A Stat. Mech. Appl.* **167**, 919–934. [https://doi.org/10.1016/0378-4371\(90\)90299-8](https://doi.org/10.1016/0378-4371(90)90299-8) (1990).
33. Davies, E. Markovian master equations. *Commun. Math. Phys.* **39**, 91–110. <https://doi.org/10.1007/BF01608389> (1974).
34. Vaidman, L. *et al.* Weak value beyond conditional expectation value of the pointer readings. *Phys. Rev. A* **96**, 032114. <https://doi.org/10.1103/PhysRevA.96.032114> (2017).
35. Duprey, Q. & Matzkin, A. Null weak values and the past of a quantum particle. *Phys. Rev. A* **95**, 032110. <https://doi.org/10.1103/PhysRevA.95.032110> (2017).
36. Sokolovski, D. Comment on “null weak values and the past of a quantum particle”. *Phys. Rev. A* **97**, 046102. <https://doi.org/10.1103/PhysRevA.97.046102> (2018).
37. Duprey, Q. & Matzkin, A. Reply to “comment on ‘null weak values and the past of a quantum particle’”. *Phys. Rev. A* **97**, 046103. <https://doi.org/10.1103/PhysRevA.97.046103> (2018).
38. Banaszek, K., Horodecki, P., Karpinski, M. & Radzewicz, C. Quantum mechanical which-way experiment with an internal degree of freedom. *Nat. Commun.* **4**, 2594. <https://doi.org/10.1038/ncomms3594> (2013).
39. Łobjko, M., Mierzejewski, M. & Dajka, J. Interference of qubits in pure dephasing and almost pure dephasing environments. *J. Phys. A Math. Theor.* **48**, 275302. <https://doi.org/10.1088/1751-8113/48/27/275302> (2015).
40. Dajka, J., Mierzejewski, M. & Łuczka, J. Fidelity of asymmetric dephasing channels. *Phys. Rev. A* **79**, 012104. <https://doi.org/10.1103/PhysRevA.79.012104> (2009).
41. Schuster, D. I. *et al.* Resolving photon number states in a superconducting circuit. *Nature* **445**, 515–518. <https://doi.org/10.1038/nature05461> (2007).
42. Roszak, K. & Machnikowski, P. Complete disentanglement by partial pure dephasing. *Phys. Rev. A* **73**, 022313. <https://doi.org/10.1103/PhysRevA.73.022313> (2006).
43. Reina, J. H., Quiroga, L. & Johnson, N. F. Decoherence of quantum registers. *Phys. Rev. A* **65**, 032326. <https://doi.org/10.1103/PhysRevA.65.032326> (2002).
44. Kruchinin, S. Y. Non-markovian pure dephasing in a dielectric excited by a few-cycle laser pulse. *Phys. Rev. A* **100**, 043839. <https://doi.org/10.1103/PhysRevA.100.043839> (2019).
45. Chen, H.-B. *et al.* Quantifying the nonclassicality of pure dephasing. *Nat. Commun.* **10**, 3794. <https://doi.org/10.1038/s41467-019-11502-4> (2019).
46. Dajka, J. Scattering-like control of the cheshire cat effect in open quantum systems. *Quantum Rep.* **2**, 1–11. <https://doi.org/10.3390/quantum2010001> (2020).
47. Richter, M., Dziewit, B. & Dajka, J. The quantum cheshire cat effect in the presence of decoherence. *Adv. Math. Phys.* **2018**, 7060586. <https://doi.org/10.1155/2018/7060586> (2018).
48. Bratteli, O. & Robinson, D. W. *Operator Algebras and Quantum Statistical Mechanics: Equilibrium States. Models in Quantum Statistical Mechanics* (Springer, 2003).
49. Lendi, K. & van Wonderen, A. J. Davies theory for reservoir-induced entanglement in a bipartite system. *J. Phys. A Math. Theor.* **40**, 279–288. <https://doi.org/10.1088/1751-8113/40/2/007> (2006).
50. Dajka, J., Mierzejewski, M., Łuczka, J., Blattmann, R. & Hänggi, P. Negativity and quantum discord in Davies environments. *J. Phys. A Math. Theor.* **45**, 485306. <https://doi.org/10.1088/1751-8113/45/48/485306> (2012).
51. Dajka, J. & Łuczka, J. Quantum cloning disturbed by thermal Davies environment. *Quantum Inf. Process.* **15**, 2661. <https://doi.org/10.1007/s11128-016-1278-z> (2016).
52. Dajka, J., Łuczka, J. & Hänggi, P. Geometric phase as a determinant of a qubit–environment coupling. *Quantum Inf. Process.* **10**, 85–96. <https://doi.org/10.1007/s11128-010-0178-x> (2011).
53. Alicki, R. & Lendi, K. *Quantum Dynamical Semigroups and Applications* (Springer, Berlin, 2007).
54. Johansson, J., Nation, P. & Nori, F. Qutip 2: A python framework for the dynamics of open quantum systems. *Comput. Phys. Commun.* **184**, 1234–1240. <https://doi.org/10.1016/j.cpc.2012.11.019> (2013).
55. Johansson, J., Nation, P. & Nori, F. Qutip: An open-source python framework for the dynamics of open quantum systems. *Comput. Phys. Commun.* **183**, 1760–1772. <https://doi.org/10.1016/j.cpc.2012.02.021> (2012).

## Competing interests

The authors declare no competing interests.

## Additional information

**Correspondence** and requests for materials should be addressed to J.D.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021