




You have downloaded a document from
RE-BUS
repository of the University of Silesia in Katowice

Title: Elevated light intensity compensates for nitrogen deficiency during chrysanthemum growth by improving water and nitrogen use efficiency

Author: Sara Esmaeili, Sasan Aliniaiefard, Shirin Dianati Daylami, Soheil Karimi, Aida Shomali, Edyta Sierka i in.

Citation style: Esmaeili Sara, Aliniaiefard Sasan, Dianati Daylami Shirin, Karimi Soheil, Shomali Aida, Sierka Edyta i in. (2022). Elevated light intensity compensates for nitrogen deficiency during chrysanthemum growth by improving water and nitrogen use efficiency. "Scientific Reports" (Vol. 12 (2022), art. no. 10002), doi 10.1038/s41598-022-14163-4



Uznanie autorstwa - Licencja ta pozwala na kopiowanie, zmienianie, rozprowadzanie, przedstawianie i wykonywanie utworu jedynie pod warunkiem oznaczenia autorstwa.



UNIWERSYTET ŚLĄSKI
W KATOWICACH



Biblioteka
Uniwersytetu Śląskiego



Ministerstwo Nauki
i Szkolnictwa Wyższego



OPEN

Elevated light intensity compensates for nitrogen deficiency during chrysanthemum growth by improving water and nitrogen use efficiency

Sara Esmaeili¹, Sasan Aliniaiefard^{1✉}, Shirin Dianati Daylami², Soheil Karimi², Aida Shomali¹, Fardad Didaran¹, Arkadiusz Telesiński³, Edyta Sierka⁴ & Hazem M. Kalaji⁵

Identifying environmental factors that improve plant growth and development under nitrogen (N) constraint is essential for sustainable greenhouse production. In the present study, the role of light intensity and N concentrations on the biomass partitioning and physiology of chrysanthemum was investigated. Four light intensities [75, 150, 300, and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ photosynthetic photon flux density (PPFD)] and three N concentrations (5, 10, and 15 mM N L^{-1}) were used. Vegetative and generative growth traits were improved by increase in PPFD and N concentration. High N supply reduced stomatal size and g_s in plants under lowest PPFD. Under low PPFD, the share of biomass allocated to leaves and stem was higher than that of flower and roots while in plants grown under high PPFD, the share of biomass allocated to flower and root outweighed that of allocated to leaves and stem. As well, positive effects of high PPFD on chlorophyll content, photosynthesis, water use efficiency (WUE), Nitrogen use efficiency (NUE) were observed in N-deficient plants. Furthermore, photosynthetic functionality improved by raise in PPFD. In conclusion, high PPFD reduced the adverse effects of N deficiency by improving photosynthesis and stomatal functionality, NUE, WUE, and directing biomass partitioning toward the floral organs.

Nitrogen (N) is one of the most important nutritional elements for plants and the main constituent of many essential molecules such as chlorophyll, amino acids, and nucleic acids¹⁻³. It also plays a crucial role as signaling molecule in form of nitric oxide under stress condition⁴. The deficiency of N is often a limiting factor for plant growth and development^{5,6}. Modern agriculture approaches to maximize the efficiency of resources. N is known as the most consumed fertilizer used to increase crop production⁷. However, more than half of the N that used in agriculture is lost through leaching and evaporation⁸. Excessive use of chemical fertilizers, especially N fertilizers, increases the cost of crop production and greenhouse gas emissions, as well as soil and groundwater pollution^{9,10}. Therefore, optimization of crop production under nitrogen restriction conditions is a major challenge in agriculture¹¹.

Light is the primary energy source for photosynthesis. Light intensity, quality (spectrum), and duration (photoperiod) influence plant growth and development^{12,13}. The main absorption spectra of the chlorophyll pigments are in the range of blue and red light, therefore, these two spectra are mainly used for plant production in controlled environments¹³. Besides light quality, the intensity of the light also exerts a broad range of physiological effects on plant growth and performance¹⁴⁻¹⁶, including, N uptake and allocation¹⁷.

Light is among the environmental factors that affect the absorption and reduction of nitrogen in plants¹⁸. Light intensity and N concentration have been shown to play a crucial role in the N uptake and leaf N content

¹Photosynthesis Laboratory, Department of Horticulture, Aburaihan Campus, University of Tehran, Tehran 33916-53755, Iran. ²Department of Horticulture, Aburaihan Campus, University of Tehran, Tehran 33916-53755, Iran. ³Department of Bioengineering, West Pomeranian University of Technology in Szczecin, 17 Słowackiego Street, 71-434 Szczecin, Poland. ⁴Institute of Biology, Biotechnology and Environmental Protection, Faculty of Natural Sciences, University of Silesia in Katowice, 28 Jagiellonska, 40-032 Katowice, Poland. ⁵Department of Plant Physiology, Institute of Biology, Warsaw University of Life Sciences—SGGW, 02-787 Warsaw, Poland. ✉email: aliniaiefard@ut.ac.ir

of plants. It was shown that raise in light intensity increases the activity of enzymes involved in N metabolism¹⁹. In addition provision of adequate irradiance increases the N assimilation in plant leaves by providing enough energy for CO₂ fixation²⁰.

Today, there is a high tendency to produce ornamental plants in greenhouses and controlled environments and there is a need to determine the desirable PPFD for plant growth in an approach with the highest resource use efficiency. The nature and extent of plant response to light intensity are dependent on the N status of the plant and vice versa^{21,22}. Therefore, optimizing the N concentration of nutrient solution based on light intensity can be effective in achieving the best growth and yield of ornamental plants. To our knowledge, so far, no study has examined the effect of increasing light intensity as an effective strategy and environmentally compatible to reduce growth defects under N deficient conditions and increase nitrogen use efficiency (NUE).

Chrysanthemum is one of the most popular ornamental plants, which ranked second place after roses in the ornamental industry worldwide²³. Given that, both light and N are the most important limiting factors for the growth and yield of chrysanthemum. In this study, we hypothesized that light intensity may affect N metabolism and an increase in light intensity may compensate for N deficiency.

Here, we investigated the changes of the morphological and physiological characteristics, as well as, biomass partitioning of chrysanthemum in response to different combinations of light intensity and N concentrations of the nutrient solution. This study provides valuable insights into the interactive regulation of light intensity and N supply to improve the quality of chrysanthemum production.

Results

N constraint intervention in chlorophyll fluorescence is more pronounced in old leaves rather than young leaves under different PPFDs. To evaluate the effects of different N supplies under various PPFDs on photosynthesis functionality, F_v/F_m of chrysanthemum young and old leaves was measured. An increase in F_v/F_m was observed by the increase in PPFD while the effect of N concentration was negligible. However, the effect of N concentration on F_v/F_m on old leaves of plants grown under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$ was considerable compared to old leaves of plants grown under other PPFDs. Under low light intensity, F_v/F_m decreased by 6% and 17% by an increase in N concentration from 5 to 10 and 15 mM N L⁻¹, respectively (Fig. 1A). PI_{ABS} was increased by raise in PPFD but the effect of N concentration was negligible on young leaves. In old leaves, however, by the increase in N concentration from 5 to 10 and 15 mM N L⁻¹, 27% and 45% decline displayed by plants grown under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$. In contrast, the old leaves of plants grown under 150, 300, and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$, showed higher PI_{ABS} , when fed by 10 and 15 mM N L⁻¹ compared to that of plants under similar PPFD but fed by 5 mM N L⁻¹ (Fig. 1B). Considering the effects of PPFD on PI_{ABS} of plants under N constraint revealed that, increase in PPFD enhanced PI_{ABS} of plants indicated by 13%, 22%, and 32% increase in young leaves and 2%, 23%, and 38% increase in old leaves by raise in PPFD from 75 to 150, 300 and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$, respectively (Fig. 1B).

Chlorophyll concentration of leaves were strongly associated with nitrogen content under high PPFD but not under low PPFD. Chlorophyll content increased stepwise by the increase in both PPFD and N concentration (Fig. 2A). By increase in N concentration from 5 to 10 and 15 mM N L⁻¹, the increase in chlorophyll content was 24% and 38% under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$, 10% and 39% under 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$, 11% and 27% under 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and 19% and 30% under 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Supplementary Fig. 1). Furthermore, under N constraint (5 mM N L⁻¹), an increase in PPFD compensated for chlorophyll synthesis. By raise in PPFD from 75 to 150, 300 and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$, 40%, 60% and 76% increases were observed in chlorophyll content (Fig. 2A). Comparing the chlorophyll content of young and old leaves revealed that, regardless of N concentration and PPFD, the chlorophyll content of old leaves was smaller than young leaves. Moreover, in both young and old leaves increase in PPFD increased the chlorophyll content of the leaves regardless of N concentration. In young leaves, a raise in N concentration increased the chlorophyll content of plants under each PPFDs (Fig. 2A). In old leaves, however, a contrasting trend was observed in the chlorophyll content of leaves in plants grown under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and plants grown under higher PPFDs. The chlorophyll content of the old leaves of plants under PPFDs higher than 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$ increased by raise in N concentration, while its content in plants grown under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$ decreased by an increase in N concentration (Fig. 2A). The nitrogen content of the leaves increased by raise in PPFD in all N concentrations (Fig. 2B). The highest N content was observed in plants fed by 15 mM N L⁻¹ under 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and plants fed by 10 and 15 mM N L⁻¹ under 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$. N content in plants grown under 300 and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and 15 mM N L⁻¹ was 33% higher than that of plants under the same N concentration but exposed to 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 2B). When plants were fed by 10 mM N L⁻¹, the N content of leaves was 16%, 29%, and 49% higher in plants grown under 150, 300, and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ compared to the plants fed by 5 mM N L⁻¹. In plants fed by 5 mM N L⁻¹, the compensatory role of PPFD was conceived since 8%, 22%, and 31% increase in N content of leaves obtained by an increase in PPFD from 150 to 300 and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 2B).

Moreover, the correlation between N content of the leaves with chlorophyll content was negligible in plants exposed to 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$ but the chlorophyll content of plants grown under 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ was positively correlated with N content of the leaves. Nevertheless, when the PPFD reached 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ the correlation dropped drastically (Fig. 2C; Supplementary Figs. 2, 3).

Stomatal aperture and gas exchange enhanced by raise in PPFD but not affected by N unless the PPFD was low. Stomatal traits showed contrasting responses to N concentration when exposed to different light intensities. In plants grown under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$, stomatal aperture, stomatal length and width and stomatal conductance decreased by the increase in N concentration (Fig. 3A-E). However, the opposing trend




Figure 1. Quantum yield of PSII (F_v/F_m) (A), and performance index per absorbed light (PI_{ABS}) (B) of young (Y) and old (O) leaves of chrysanthemum plants grown under different light intensities (75, 150, 300, and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$) and nitrogen (N) concentrations (5, 10, and 15 mM N) at 70 days of cultivation. Vertical bars are means \pm SD. Within each graph, interactive effects of light intensity and nitrogen concentration are shown. Different letters (a–d) denote a significant difference between treatments ($P < 0.05$).

was observed when plants were exposed to PPFDs higher than 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$. In plants grown under 150 and 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$, stomatal length and stomatal pore length increased by the increase in N concentration while under 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$, increase in N concentration didn't affect stomatal traits. A negligible increase in stomatal width, stomatal pore width, and stomatal conductance was displayed by the increase in N concentration in plants grown under 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 3A–E).

Light intensity affect the accumulation and partitioning of biomass under contrasting nitrogen regimes. The total biomass was affected by PPFD and N concentration. By increase in N concentration, a stepwise elevation was observed in total biomass under each PPFD except for 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Regardless of N concentration, total biomass increased by raise in PPFD, except the biomass of plants under 5 mM N L⁻¹ that did not differ under 75 or 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 4A). Under N constraint, PPFD could compensate for N limitation for biomass accumulation. In plants fed by 5 mM N L⁻¹ total biomass of plants grown under 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ was twofold of the plants grown under 75 and 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and 1.5-fold of the total biomass of plants under 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 4A).

Partitioning of biomass to different plant organs was also affected by both light intensity and N concentration (Fig. 4B). Root biomass showed 52% and 41% decline by the increase in N concentration from 5 to 10 and 15 under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 4B). Increasing PPFD from 75 to 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$, partitioning of biomass to root decreased by 0.8% and 12% respectively by increase in N concentration from 5 to 10 and 15. Same as the plants exposed to 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$, by the increase in N concentration from 5 to 10 and 15, respectively 17% and 16% increase in the partitioning of biomass to the roots were detected in plants exposed to 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 4B). However, the reduction in the partitioning of biomass to root was more drastic under 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ in plants fed by 10 and 15 mM N L⁻¹ compared to the plants fed by 5 mM N L⁻¹, indicated by 45% and 55% reduction in root biomass, suggesting the compensatory effect of high PPFD on the partitioning of biomass to root under N constraint (Fig. 4B).

Partitioning of biomass to leaves increased when N concentration raised from 5 to 10 and 15 mM N L⁻¹ under all PPFDs. By increase in N concentration from 5 to 10 and 15 respectively, the increase in partitioning




Figure 2. Chlorophyll content of the leaves (A), Nitrogen content of leaves (B), and the correlation between N and chlorophyll content (C) of the leaves of chrysanthemum plants grown under different light intensities (75, 150, 300, and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$) and nitrogen (N) concentrations (5, 10, and 15 mM N) at 70 days of cultivation. Vertical bars are means \pm SD. Within each graph, interactive effects of light intensity and nitrogen concentration are shown. Different letters (a–d) denote a significant difference between treatments ($P < 0.05$).

of biomass to leaves were recorded as 7% and 38% in plants under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$, 38% and 17% in plant under 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$, 5% and 75% increase in plants under 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$, and 47% and 70% in plants under 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 4B).

Partitioning of biomass to stem increased by raise in N concentration from 5 to 10 and 15 mM N L⁻¹ by 6% in plants under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and 16% and 9% in plants under 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$. In contrast, in plants grown under 150 and 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$, a small reduction was observed in biomass partitioning to stem indicated by 10% and 3% decrease respectively by raise in N concentration from 5 to 10 and 15 mM N L⁻¹. A 5% decline in the partitioning of biomass to stem was also displayed by plants grown under 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and fed by 15 mM N L⁻¹ compared to plants under the same PPFD but 5 mM N L⁻¹ (Fig. 4B).

Partitioning of biomass to flower was also affected by N concentration and PPFD. In plants grown under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$, a 10% increase in the partitioning of biomass to flower was observed by raise in N concentration from 5 to 10 mM N L⁻¹. Whereas, an increase in N concentration from 5 to 15 mM N L⁻¹ resulted in a 38% decline in biomass partitioning to flower (Figs. 4B, 5). Increase in N concentration from 5 to 10 and 15 mM N L⁻¹ cut down on partitioning of biomass to flowers by 20% and 3% under 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$. As well, 33% increase in biomass partitioning to flowers observed in plants under 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ when N concentration increase from 5 to 15 mM N L⁻¹. In plants grown under 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$, an increase in N concentration from 5 to 10 mM N L⁻¹ reduced biomass partitioning to flower (Fig. 4B).

Overall, comparing the partitioning of biomass to different organs revealed that, under low PPFD, the share of biomass allocated to leaves (26%) and stem (41%) was higher than that of flower (19%) and root (13%), while, in plants grown under high PPFD, the share of biomass allocated to flower (27%) and root (23%) increased at the expense of biomass accumulation in leaves (23%) and stem (13%) (Fig. 4C,D).

WUE decreased by higher N concentration under low PPFD while increased by higher N concentration under high PPFD. WUE was affected by N concentration but came to contradictory effects in plants grown under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$ compared to plants grown under 150, 300 and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 6A). Under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$, by increase in N concentration form 5 to 10 and 15 mM N L⁻¹, 25 and 43% reduction in WUE was observed, conversely, 8% and 77% increase under 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$, 94% and 54% increase under 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and 15% and 72% increase under 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ were detected in WUE of plants




Figure 3. Stomatal length (A), stomatal pore length (B), stomatal width (C), stomatal pore width (D), and stomatal conductance (g_s) (E) of chrysanthemum plants grown under different light intensities (75, 150, 300, and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$) and nitrogen (N) concentrations (5, 10, and 15 mM N) at 70 days of cultivation. Vertical bars are means \pm SD. Within each graph, interactive effects of light intensity and nitrogen concentration are shown. Different letters (a–d) denote a significant difference between treatments ($P < 0.05$).

by the increase in the concentration of N from 5 to 10 and 15 mM N L^{-1} , respectively (Fig. 6A). Considering the effect of PPFD on WUE of plants grown under N constraint revealed that increase in PPFD from 75 to 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$ did not affect the WUE of plants and when PPFD was raised to 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ a negligible increase was observed, whereas when PPFD reached to 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$, WUE of plants increased to twofold of WUE of plants under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 6A). The correlation of total biomass with WUE also increased by the increase in PPFD under N constraint, indicating that the water uptaken by the plant were efficiently utilized for biomass accumulation. The correlation of PI_{ABS} with WUE was more profound on young leaves compared to old leaves; since the correlation was strongly negative under low PPFD and raised sharply under high PPFD in young leaves, while in old leaves the increase in correlation by raise in PPFD was slighter than that of young leaves (Fig. 6B). In addition, under N constraint, the correlation between PI_{ABS} and WUE was affected by PPFD.




Figure 4. Total biomass (A), biomass partitioning among different organs (B), biomass partitioning among different organs in plants grown under $75 \mu\text{mol m}^{-2} \text{s}^{-1}$ and 5 mM N (C) and biomass partitioning among different organs in plants grown under $600 \mu\text{mol m}^{-2} \text{s}^{-1}$ and 5 mM N (D) of chrysanthemum grown under different light intensities (75 , 150 , 300 , and $600 \mu\text{mol m}^{-2} \text{s}^{-1}$) and nitrogen (N) concentrations (5 , 10 , and 15 mM N) at 70 days of cultivation. Vertical bars are means \pm SD. Within each graph, interactive effects of light intensity and nitrogen concentration are shown. Different letters (a–d) denote a significant difference between treatments ($P < 0.05$).

WUE was negatively correlated with PI_{ABS} of young and old leaves under N constraint but an increase in PPFD strengthen the correlation and increase it from -76 to -50% in young and old leaves under $75 \mu\text{mol m}^{-2} \text{s}^{-1}$ to 99% and 61% under $600 \mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 6B). The effect of PPFD on increasing WUE under N constraint was more pronounced than that of old leaves. Moreover, according to correlation results, WUE and PI_{ABS} are more closely associated in young leaves compared to the old leaves (Fig. 6B; Supplementary Figs. 2, 3).

Carbohydrate level depleted by increasing N concentration of nutrient solution under low PPFD, while accumulated by exposure to higher PPFDs.

A contrasting trend was observed on the effect of N concentration on soluble carbohydrates under different light intensities (Fig. 7). In plants grown under $75 \mu\text{mol m}^{-2} \text{s}^{-1}$ the carbohydrate decreased by 44% and 57% by an increase in N concentration from 5 to 10 and 15 mM N L^{-1} , respectively (Fig. 7). In contrast, in plants grown under PPFDs higher than $75 \mu\text{mol m}^{-2} \text{s}^{-1}$, the carbohydrate content increased by raise in N concentration. Under $150 \mu\text{mol m}^{-2} \text{s}^{-1}$, 88% increase in carbohydrate was detected in plants fed by 10 and 15 mM N L^{-1} compared to 5 mM N L^{-1} (Fig. 7). Moreover, by raise in N concentrations from 5 to 10 and 15 mM N L^{-1} , 2 - and 3 -fold increases were observed in the carbohydrate of plants grown under 300 and $600 \mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 7). Regardless of N concentration, a raise in PPFD increased soluble carbohydrates. When plants were fed by 5 mM N L^{-1} , 9% , 32% , and 12% reduction were observed in carbohydrates by the increase in PPFD from 75 to 150 , 300 and $600 \mu\text{mol m}^{-2} \text{s}^{-1}$. However, in plants fed by 10 mM N L^{-1} , raise in PPFD from 75 to 150 , 300 and $600 \mu\text{mol m}^{-2} \text{s}^{-1}$ resulted in 2 -, 3 - and 5 -fold and under 15 mM N L^{-1} 2 -, 5 - and sevenfold increase in carbohydrate contents were observed by the increase in PPFD from 75 to 150 , 300 and $600 \mu\text{mol m}^{-2} \text{s}^{-1}$, suggesting a positive role of PPFD on carbohydrate synthesis (Fig. 7). Our data also revealed that a negative correlation existed between WUE and carbohydrate under N constraint and low light (-70%) and a raise in PPFD ameliorated the negative correlation, since under $600 \mu\text{mol m}^{-2} \text{s}^{-1}$ and N constraint, 61% correlation was detected between WUE and carbohydrate (Fig. 6B; Supplementary Figs. 2, 3).

NUE decreased by higher N concentration while increased by higher PPFDs.

NUE was increased by raise in PPFD, while declined by the increase in N concentration under all PPFDs (Fig. 8A). The highest NUE was associated with plants fed by the lowest N concentration and grown under the highest light intensity. When




Figure 5. Interaction effect of light intensity and concentration of nitrogen (N) on the flowering of chrysanthemum. Plants were exposed to different concentrations of nitrogen (N) (5, 10, and 15 mM N) and light intensities (75, 150, 300, and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$).

plants were fed by 5 mM N L^{-1} , NUE was increased by 77% and 169% by raise in PPFD from 75 to 300 and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and remained unchanged in plants grown under 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 8A). In other words, a raise in PPFD compensated for the adverse effects of N constraint on NUE since the highest PPFD was associated with the highest NUE in plants grown under 5 mM N L^{-1} . Under 300 and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$, NUE of plants fed by 10 mM N L^{-1} was threefold of NUE of plants grown under 75 and 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Under 150, 300, and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$, when plants were fed by 15 mM N L^{-1} , NUE was 2-, 3- and 7-fold of NUE of plants grown under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 8A).

C:N ratio was decreased by raise in N concentration from 5 to 10 and 15 mM N L^{-1} by 56% and 66% respectively, in plants under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and reduced by 14% and 7% in plant under 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Under 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ however, C:N ratio showed 34% and 38% increase by raise in N concentration from 5 to 10 and 15 mM N L^{-1} (Fig. 8B).

The relation between NUE and biomass accumulation was also affected by PPFD. Under N constraint (5 mM N L^{-1}) NUE was negatively correlated (-73%) with total biomass while, when PPFD was raised to 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$, a positive correlation (58%) was detected between NUE and total biomass (Fig. 8C). Chlorophyll content was also demonstrated a negative correlation (-78%) with NUE under low light and N constraint,




Figure 6. Water use efficiency (WUE) (A) of chrysanthemum plants grown under different light intensities (75, 150, 300, and 600 μmol m⁻² s⁻¹) and nitrogen (N) concentrations (5, 10, and 15 mM N) at 70 days of cultivation. The correlation between WUE with performance index per absorbed light (PI_{ABS}), total biomass, WSC, and C:N ratio of plants fed by 5 mM N with under differently light intensities (150, 300, and 600 μmol m⁻² s⁻¹) compared to that of plants under 75 μmol m⁻² s⁻¹ (B). Vertical bars are means ± SD. Within each graph, interactive effects of light intensity and nitrogen concentration are shown. Different letters (a–d) denote a significant difference between treatments (P < 0.05).




Figure 7. Water soluble carbohydrates (WSC) of chrysanthemum plants grown under different light intensities (75, 150, 300, and 600 μmol m⁻² s⁻¹) and nitrogen (N) concentrations (5, 10, and 15 mM N) at 70 days of cultivation. Vertical bars are means ± SD. Within each graph, interactive effects of light intensity and nitrogen concentration are shown. Different letters (a–d) denote a significant difference between treatments (P < 0.05).




Figure 8. NUE of leaves (A) C:N ratio (B), of chrysanthemum plants grown under different light intensities (75, 150, 300, and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$) and nitrogen (N) concentrations (5, 10, and 15 mM N) at 70 days of cultivation. The correlation between NUE of plants fed by 5 mM N with performance index per absorbed light (PI_{ABS}), total biomass, total chlorophyll (chl), C:N ration, water soluble carbohydrates (WSC), and nitrogen concentration under differently light intensities (150, 300, and 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$) compared to that of plants under 75 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (C). Vertical bars are means \pm SD. Within each graph, interactive effects of light intensity and nitrogen concentration are shown. Different letters (a–d) denote a significant difference between treatments ($P < 0.05$).

whereas, under 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and N constraint the negative correlation ameliorated to a 24% positive correlation (Fig. 8C; Supplementary Figs. 2, 3).

Discussion

F_v/F_m reflects the maximum photochemical quantum yield of PSII²⁴. In this research, variation in F_v/F_m under different treatments was negligible. However, the change in F_v/F_m was more detectable in old leaves under high N concentration compared to young leaves, suggesting that the dependency of F_v/F_m value to N concentration depends on leaf age and PPFD. The fact that plants respond to N supply by redistributing N from older leaves to the youngest can elaborate on the dependency of F_v/F_m to leaf age²⁵. Moreover, under the N constraint condition, the correlation between NUE and F_v/F_m showed opposite trends by the increase in PPFD. Under N constraint and low PPFD, a strong positive correlation was detected between NUE and F_v/F_m of young leaves, which reduced to a strongly negative correlation by raise in PPFD. Whereas, the correlation was strongly negative in old leaves under low PPFD and ameliorated to a small positive correlation under high PPFD. This finding may explain that under N constraint, old leaves invest more N for increasing photosynthesis efficiency compared to that of young leaves because young leaves invest N for other pathways like biomass accumulations or carbohydrate synthesis since the correlation of NUE with total biomass and carbohydrate was higher than the correlation between NUE and F_v/F_m under high PPFD and N constraints.

PI_{ABS} is a delicate parameter derived from chlorophyll fluorescence that indicates the photochemical performance of photosynthesis^{26–28}. The reduction in PI_{ABS} of old leaves under low PPFD by raise in N concentration indicates the decline in energy conversion ability and the photosynthesis apparatus activity, which is also portrayed by light curves of plants under different PPFDs and N concentrations (Supplementary Fig. 4). This reduction in energy conversion potential accounts for reduced NUE and WUE under low PPFD. This circumstance was accompanied by a reduction in soluble carbohydrates which is the output of the photosynthesis process. The correlation between PI_{ABS} with NUE and WUE was negative under low light and low N concentration, while under high PPFD a positive correlation was detected. This finding allows us to postulate that increase in PPFD

compensates for N limitation by increasing PI_{ABS} , denoting higher efficiency of energy conversion that further provides energy for N uptake and increases the requirements for water uptake.

The negative correlation between chlorophyll content and N content under high PPFDs may account for the increase in the use of N in other processes besides chlorophyll synthesis since by increase in PPFD the increase in the correlation between NUE and total chlorophyll was lower than the correlation between NUE and other parameters like biomass and C:N balance. This may explain, in part, the strong reduction in correlation between chlorophyll and N content under high PPFD. In this regard, previous reports also proposed the effect of PPFD on the allocation of N between photosynthetic and non-photosynthetic pools as well as different N allocation between Rubisco and light-harvesting^{29,30}. Besides, our data demonstrated a compensative role of PPFD for chlorophyll synthesis under N constraint conditions despite the negative correlation maintained among the N content of the leaves and chlorophyll content. This discrepancy suggests that high PPFD tends to tune the balance between the use of N among different pathways, while under low PPFD the negligible correlation between N content with chlorophyll, biomass, and C:N ratio explains the lack of balance between partitioning of N to different pathways under N constraint. Furthermore, the close relationship between N metabolism and light signals can be the other explanation for the role of PPFD in compensation of N constraint since N uptake is regulated by shoot-borne light signals, has shown to up-regulate the expression of root nitrogen transporters and enhance N uptake eventually^{31,32}. Also, an association between high light intensity, plant water uptake, transpiration and the uptake of soil nitrogen by roots were proposed which is governed by mass flow water movement³¹. Many studies have reported that chlorophyll a, chlorophyll b, and total chlorophyll content correlate strongly with leaf N concentration so that chlorophyll content is reduced with N deficiency³³.

The fact that an increase in stomatal aperture and stomatal conductance was accompanied by an increase in WUE explains the increase in PI_{ABS} and the highest biomass accumulation observed under high PPFD. Under N constraint, this circumstance accounts for improved growth and physiological status of plants, proposing a compensative role of PPFD for N constraint. An increase in N concentration and PPFD positively affected the N content of the leaves. The same effect was also reported on *Lolium perenne* L.³⁴. Light signaling within the root system may trigger N uptake through the mediation of root nitrogen transporters^{31,32}, further consequence in the enhancement of plant water uptake, transpiration, and the acquisition of soil nitrogen by roots under high light condition³¹. The correlation between leaf N concentration and NUE was negative under the N constraint condition, however, a raise in PPFD weakened this negative correlation, suggesting enhancing the role of PPFD for NUE under N constraint condition.

The increase in total biomass by the rise in PPFD and N concentration in plants grown under PPFDs higher than $75 \mu\text{mol m}^{-2} \text{s}^{-1}$ can be explained by improve in dry and fresh weight of plant organs (Supplementary Fig. 5) as well as the number of leaves, number of flowers, and plant height (Supplementary Fig. 6), which were the result of improved photosynthetic traits along with elevation of NUE and WUE. Under N constraint the correlation between total biomass with NUE and WUE increased sharply by raise in PPFD. The increased efficiency of water and nitrogen use may account for the ameliorative effect of PPFD on biomass accumulation under N constraint. Root biomass increased by the increase in light intensity and N concentration, which is in agreement with previous reports³⁵. Nevertheless, the share of root biomass in total biomass decreased by raise in N concentration at the expense of an increase in the partitioning of biomass to aerial organs indicated by an increase in biomass partitioning to leaves. The fact that partitioning of biomass to root was limited by raise in N concentrations can be elaborated by optimal partitioning theory which explains that the biomass is allocated to the organ of the plant that is exposed to the most limiting resource^{36–38}. This partitioning strategy is thought to minimize the stress imposed by the limiting resource³⁹. Contrary to N concentration, rise in PPFD has shown to increase the partitioning of biomass to root, which can be vindicated by increased carbon availability provided by higher photosynthesis functionality under high PPFDs⁴⁰. Concurringly, our data revealed a higher C:N ratio under high PPFD. Moreover, the correlation between NUE and C:N ratio was increased by the increase in PPFD, suggesting a balance in the increase in carbon assimilation and N uptake. An increase in PI_{ABS} along with WUE led to elevated carbon assimilation. In order to N uptake keep pace with carbon assimilation to maintain the balance of C:N, root biomass increased to maximize N uptake under N constraint. This circumstance elaborates on the underlying mechanism through which PPFD compensated for N uptake under N constraint. Concurred with our results, the positive effect of PPFD on root biomass reported on lettuce¹⁴ and chrysanthemum⁴¹. Regardless of PPFD, increase in N concentration reduced the partitioning of biomass to flowers. However, in the same N concentration, a larger proportion of biomass is allocated to flowers by the increase in PPFD, which can be explained by the increase in the number of flowers displayed by increased PPFD and N concentration (Supplementary Fig. 6).

We observed that the adverse effects of nitrogen deficiency extended to the reproductive growth stage and eventually prolonged the flower emergence period and decreased the biomass partitioning to the flower under low PPFD. The positive role of PPFD on the flower emergence period and the increasing number of flowers have been also declared in the previous study on chrysanthemum⁴¹. In addition, a strong genetic correlation between flowering date and NUE has been reported in wheat⁴². In the same line, our data denotes a close relationship between light intensity and NUE, which may in part explain the promoting effect of PPFD on flower emergence, regardless of N concentration, since under all N treatments the correlation between NUE and PPFD was high ($R^2 \geq 98\%$). Moreover, the correlation between NUE and day to flowering was negative under high PPFD and positive under low PPFD, suggesting that, increase in NUE shortens day to flowering under high PPFD and remains the opposite effect under low PPFD (Supplementary Figs. 2, 3).

Under high PPFD, WUE improved by the increase in N concentration, which is in line with the results of a previous study on *Robinia pseudoacacia*⁴³. However, according to our data, when PPFD was limited to $75 \mu\text{mol m}^{-2} \text{s}^{-1}$, WUE decreased by raise in N concentration. Consistently, in *Capsicum annum* L., the WUE of shaded plants was far lower than unshaded plants⁴⁴. On the other hand, under N constraint, PPFD compensated

Element	Full strength Hoagland solution (g/L)	Modified Hoagland solution containing 10 mL nitrogen (g/L)	Modified Hoagland solution containing 5 ml nitrogen (g/L)
KNO ₃	101.1	–	–
CaCl ₂	–	–	110.9
K ₂ SO ₄	–	87.1	87.1
Ca(NO ₃) ₂ ·4H ₂ O	236.1	236.1	236.1
MgSO ₄ ·7H ₂ O	246.5	246.5	246.5
KH ₂ PO ₄	136.1	136.1	136.1
H ₃ BO ₃	2.86	2.86	2.86
MnCl ₂ ·4H ₂ O	1.81	1.81	1.81
CuSO ₄ ·5H ₂ O	0.08	0.08	0.08
Na ₂ MoO ₄ ·2H ₂ O	0.12	0.12	0.12
H ₂ MoO ₄ ·H ₂ O	0.09	0.09	0.09
ZnSO ₄ ·5H ₂ O	0.22	0.22	0.22
FeNaEDTA	4.04	4.04	4.04

Table 1. The composition of nutrient solutions of different N levels were modified based on a full-strength Hoagland solution.

for WUE under N deficiency. The compensative effect of PPFD on WUE under N constraint is portrayed by increased PI_{ABS} , C:N ratio, and total biomass.

On contrary with the N content of the leaves, NUE decreased by an increase in N concentration however, a raise in PPFD compensated for NUE of plants under N constraint. The same results were also reported in previous experiments on three genotypes of rocket salad⁴⁵, corn⁴⁶, and rice⁴⁷. Being a complex physiological trait, NUE depends on N availability and the energy provided through photosynthesis to supply the energy required for N uptake⁴⁶. Under the N constraint condition, a negative correlation was detected between N concentration and NUE, however, this negative correlation ameliorated by increase in PPFD. Raise in PPFD under N constraint also increased the correlation of NUE with PI_{ABS} and C:N ratio. Suggesting a role for PPFD on enhancing NUE via improving energy provision through photosynthesis for N uptake and maintenance of C:N balance.

Materials and methods

Plant materials and growth conditions. The plant experiments were performed in accordance with relevant guidelines and regulations. Rooted cuttings (9 cm long with three leaves) of chrysanthemum (*Chrysanthemum morifolium* cv. Katinka) were sown in pots (14 × 10 cm) containing a mixture of cocopeat and perlite (2:1, v/v), then placed in four growth chambers (1 m × 1 m × 1 m) equipped with a fixed combination of red and blue LEDs with wavelength peaks at 660 ± 10 nm for red and 460 ± 10 nm for blue LEDs. Red and blue LEDs (70:30) were used because they are the main light spectra for photosynthesis and growth of chrysanthemum¹³. To provide different PPFDs, including 75, 150, 300, and 600 μmol m⁻² s⁻¹ and also to limit the production of heat by the light sources, LED light panels (provided by Iran Grow Light Co, Iran) were used. Light spectra and PPFDs were monitored using a Sekonic light meter (Sekonic C-7000, Japan). All plants were grown under the same climatic conditions, i.e. day/night temperature of 25/20 ± 2 °C, 50 ± 5% relative humidity (RH), and photoperiod 12/12 h light/ dark cycles. The nutrient solutions of different N levels were modified based on a full-strength Hoagland and Arnon solution and were applied to each pot three times per week (Table 1). The full-strength Hoagland solution, containing 15 mM N L⁻¹ was the control solution and Hoagland and Arnon solutions with modified N concentration (5 and 10 mM N L⁻¹) were used to apply N limitation.

The concentration of 15 mM N L⁻¹ (control) was achieved via the full-strength Hoagland and Arnon solution. Concentrations of 10 and 5 mM N L⁻¹ were achieved by total or partial substitution of KNO₃ and Ca(NO₃)₂ as sources of nitrogen in the solution. Potassium and calcium levels were equalized across treatments by adding K₂SO₄ and CaCl₂ into the solutions that had N limitations (Table 1)⁴⁸.

Chlorophyll fluorescence analysis. Fast induction of fluorescence transient (the so-called OJIP protocol) was performed on fully developed, mature leaves of plants after 45 days of growth. The samples were dark-adapted for 20 min. A Fluorpen FP 100-MAX (Photon Systems Instruments, Drasov, Czech Republic) was used for measuring the OJIP transients. Different biophysical and phenomenological parameters related to PSII status⁴⁹ were investigated by JIP-test according to the protocol described by⁵⁰.

The maximum quantum yield of PSII (F_v/F_M) and Performance index per absorbed light (PI_{ABS}) was calculated using the Eqs. (1) and (2), respectively:

$$F_v/F_M = (F_M - F_0)/F_M \quad (1)$$

$$(RC/ABS) \times (\phi P_0/(1 - \phi P_0)) \times (\psi_0/(1 - \psi_0)) \quad (2)$$

Determination of chlorophyll and nitrogen content. Total leaf nitrogen content was determined using 1 g samples of leaf tissue, based on the Kjeldahl method after 98% H₂SO₄ hot digestion. For measuring the chlorophyll content, Leaves were measured by using a SPAD-502 (Konica Minolta Corp., Solna, Sweden). This instrument determined the chlorophyll content in leaves non-destructively by considering leaf transmittance in red and near-infrared light spectra. three leaves were analyzed each time and three points were recorded per replicate leaf and were averaged⁵¹.

Determination of stomatal traits. Stomatal morphological parameters including stomatal length, stomatal width together with pore length, and aperture were measured on the young fully developed leaves. A section of the leaf midway between the tip and the base and equal distance from each edge were used for microscopic analysis. The abaxial surface of young developed leaves was coated by a thin layer of nail polish. The dry polish sample along with the adhered sticky tape was mounted on microscope slides, and stomatal morphological details were investigated under a light microscope. Images were taken by Omax top-view software version 3.5 and further analyzed using ImageJ software (U.S. National Institutes of Health, Bethesda, MD; <https://imagej.nih.gov/ij/>) to record stomatal length, stomatal width, pore length, and aperture⁵².

Stomatal conductance (g_s) was investigated according to the method described by Fanourakis et al.⁵³. In this calculation, stomatal pore depth was considered to be equal to the guard-cell width (i.e., stomatal width/2), assuming guard cells inflate to a circular cross-section. A 100 magnification was used to assess stomatal density. The number of stomata was counted on three randomly chosen areas of the same leaf from which stomatal size measurements were taken. Analysis was done on 1 mm² of the middle of the leaf on both sides of the main vein^{52,53}. Calculation of g_s was done based on the following equation:

$$g_s = \frac{(\text{diffusion coefficient}) \times (\text{stomatal density}) \times (\pi \times \text{pore aperture} \div 2 \times \text{pore length} \div 2)}{(\text{molar volume of air}) \times \left[(\text{pore depth}) + \sqrt{(\text{pore aperture} \div 2 \times \text{pore length} \div 2)} \right]} \quad (3)$$

Morphological and growth measurements. For biomass determination, the growing substrate was washed from the roots, and the plants were divided into flower, leaves, stem, and roots. The samples were weighed to determine their fresh weight (FW) and then dried in an oven at 60 °C for 72 h to reach a constant dry weight (DW).

Water use efficiency (WUE). To determine irrigation water use efficiency (WUE), the amount of water used during the growth period was recorded and after measuring the dry weight of the flowers, it was calculated with the following equation described by Karam et al.⁵⁴.

$$WUE = \frac{\text{Flower dry weight}}{\text{Consumed water}} \quad (4)$$

Nitrogen use efficiency (NUE). Nitrogen use efficiency (NUE) was calculated according to the equation below^{7,55,56}:

$$NUE = \frac{\text{Flower dry weight}}{\text{N supply}} \quad (5)$$

Measurements of carbohydrates. Determination of leaf soluble sugar was determined by the anthrone method using glucose as the standard⁵⁷. Fresh leaves (0.2 g) were extracted in 80% ethanol at 80 °C for 60 min. Then 3 mL of anthrone solution (150 mg anthrone in 100 mL 72% Sulfuric acid) was added to 0.1 mL of alcoholic extract. This mixture was placed in a water bath at 100 °C for 10 min and then cooled in an ice bath and the absorbance was spectrophotometrically recorded at 625 nm. The remaining solid part after extraction of soluble sugars was used to extract starch in 52% perchloric acid. Starch concentrations were determined by anthrone and were spectrophotometrically recorded at 630 nm as described by McCready⁵⁷.

Statistical analysis. The results represented the average mean values of six replications for each treatment. The data were analyzed using SAS software (version 9.4). The two-way analysis of variance (ANOVA) was performed to find the significant differences ($p \leq 0.05$) among treatments. Further, the Duncan multiple comparisons test was performed to compare the means. For analyzing chlorophyll fluorescence parameters, obtained data were subjected to two-way ANOVA, and for mean comparison, the Tukey multiple comparison tests were used. For stomatal characteristics, data obtained from one leaf were considered not independent, and for mean comparison, one-way ANOVA, as well as Tukey multiple comparison tests, were used.

Data availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Received: 21 January 2022; Accepted: 18 May 2022

Published online: 15 June 2022

References

- Mu, X., Chen, Q., Chen, F., Yuan, L. & Mi, G. Within-leaf nitrogen allocation in adaptation to low nitrogen supply in maize during grain-filling stage. *Front. Plant Sci.* **7**, 699 (2016).
- Shomali, A., Mashayekhi, K., Pahlevani, M. & Mousavizadeh, S. J. Effects of various nitrogen sources on synchronization of tomato somatic embryogenesis during induction and realization phases. *JPPB* **8**, 53–58 (2018).
- Hasanuzzaman, M. & Tanveer, M. *Salt and Drought Stress Tolerance in Plants: Signaling Networks and Adaptive Mechanisms* (Springer Nature, 2020).
- Aliniaiefard, S., Shomali, A., Seifikalhor, M. & Lastochkina, O. Calcium signaling in plants under drought. In *Salt and Drought Stress Tolerance in Plants: Signaling Networks and Adaptive Mechanisms*, 259–298 (Springer International Publishing, 2020).
- Zhao, D., Reddy, K. R., Kakani, V. G. & Reddy, V. Nitrogen deficiency effects on plant growth, leaf photosynthesis, and hyperspectral reflectance properties of sorghum. *Eur. J. Agron.* **22**, 391–403 (2005).
- Aboutalebijahromi, A. & Hosseini Farahi, M. Seed germination, vegetative growth and concentration of some elements in French marigold (*Tageta patula*) as influenced by salinity and ammonium nitrate. *Int. J. Hortic. Sci. Technol.* **3**, 199–209 (2016).
- Keeney, D. R. Nitrogen management for maximum efficiency and minimum pollution. *Nitrogen Agric. Soils* **22**, 605–649 (1982).
- Kano, Y., Nakagawa, H., Sekine, M., Goto, H. & Sugiura, A. Effect of nitrogen fertilizer on cell size and sugar accumulation in the leaves of cabbage (*Brassica oleracea* L.). *HortScience* **42**, 1490–1492 (2007).
- Hogewoning, S. W. *et al.* Blue light dose–responses of leaf photosynthesis, morphology, and chemical composition of *Cucumis sativus* grown under different combinations of red and blue light. *J. Exp. Bot.* **61**, 3107–3117 (2010).
- Quail, P. H. Phytochrome photosensory signalling networks. *Nat. Rev. Mol. Cell Biol.* **3**, 85–93 (2002).
- Wang, J. *et al.* Genome-wide analysis of PRR gene family uncovers their roles in circadian rhythmic changes and response to drought stress in *Gossypium hirsutum* L. *PeerJ* **8**, e9936 (2020).
- Bayat, L. *et al.* Effects of growth under different light spectra on the subsequent high light tolerance in rose plants. *AoB Plants* **10**, ply052 (2018).
- Seif, M. *et al.* Monochromatic red light during plant growth decreases the size and improves the functionality of stomata in chrysanthemum. *Funct. Plant Biol.* **48**, 515–528 (2021).
- Esmaili, M. *et al.* Assessment of adaptive neuro-fuzzy inference system (ANFIS) to predict production and water productivity of lettuce in response to different light intensities and CO₂ concentrations. *Agric. Water Manag.* **258**, 107201 (2021).
- Chen, Y. *et al.* Low UVA intensity during cultivation improves the lettuce shelf-life, an effect that is not sustained at higher intensity. *Postharvest Biol. Technol.* **172**, 111376 (2021).
- Ghorbanzadeh, P. *et al.* Dependency of growth, water use efficiency, chlorophyll fluorescence, and stomatal characteristics of lettuce plants to light intensity. *J. Plant Growth Regul.* **40**, 2191–2207 (2021).
- Frith, G. & Nichols, D. Nitrogen uptake by apple seedlings as affected by light, and nutrient stress in part of the root system. *Physiol. Plant.* **34**, 129–133 (1975).
- Mokhele, B., Zhan, X., Yang, G. & Zhang, X. Nitrogen assimilation in crop plants and its affecting factors. *Can. J. Plant Sci.* **92**, 399–405 (2012).
- Zhou, W. L., Liu, W. K. & Yang, Q. Quality changes in hydroponic lettuce grown under pre-harvest short-duration continuous light of different intensities. *J. Hortic. Sci. Biotechnol.* **87**, 429–434 (2012).
- Demšar, J., Osvald, J. E. & Vodnik, D. The effect of light-dependent application of nitrate on the growth of aeroponically grown lettuce (*Lactuca sativa* L.). *J. Am. Soc. Hortic. Sci.* **129**, 570–575 (2004).
- Casey, C. A., Mangan, F. X., Herbert, S. J., Barker, A. V. & Carter, A. K. In *XXVI International Horticultural Congress: The Future for Medicinal and Aromatic Plants* Vol. 629, 215–229.
- Wang, J.-F. & Feng, Y.-L. The effect of light intensity on biomass allocation, leaf morphology and relative growth rate of two invasive plants. *Chin. J. Plant Ecol.* **28**, 781 (2004).
- Su, J. *et al.* Current achievements and future prospects in the genetic breeding of chrysanthemum: A review. *Hortic. Res.* **6**, 1–19 (2019).
- Kalaji, H. M. *et al.* Chlorophyll a fluorescence as a tool to monitor physiological status of plants under abiotic stress conditions. *Acta Physiol. Plant* **38**, 102 (2016).
- Marschner, H. *Inorganic Plant Nutrition* 5–60 (Springer, 1983).
- Shomali, A. *et al.* Synergistic effects of melatonin and gamma-aminobutyric acid on protection of photosynthesis system in response to multiple abiotic stressors. *Cells* **10**, 1631 (2021).
- Kalaji, M. H. *Chlorophyll Fluorescence: Understanding Crop Performance—Basics and Applications* (CRC Press, 2017).
- Ashrestaghi, T. *et al.* Light intensity: The role player in cucumber response to cold stress. *Agronomy* **12**, 201 (2021).
- Croft, H. *et al.* Leaf chlorophyll content as a proxy for leaf photosynthetic capacity. *Glob. Chang. Biol.* **23**, 3513–3524 (2017).
- Kenzo, T. *et al.* Changes in photosynthesis and leaf characteristics with tree height in five dipterocarp species in a tropical rain forest. *Tree Physiol.* **26**, 865–873 (2006).
- Cohen, I., Rapaport, T., Chalifa-Caspi, V. & Rachmilevitch, S. Synergistic effects of abiotic stresses in plants: A case study of nitrogen limitation and saturating light intensity in *Arabidopsis thaliana*. *Physiol. Plant.* **165**, 755–767 (2019).
- Rachmilevitch, S., Cousins, A. B. & Bloom, A. J. Nitrate assimilation in plant shoots depends on photorespiration. *PNAS* **101**, 11506–11510 (2004).
- Boussadia, O. *et al.* Effects of nitrogen deficiency on leaf photosynthesis, carbohydrate status and biomass production in two olive cultivars 'Meski' and 'Koroneiki'. *Sci. Hortic.* **123**, 336–342 (2010).
- Alberda, T. The influence of temperature, light intensity and nitrate concentration on dry-matter production and chemical composition of *Lolium perenne* L. *Neth. J. Agric. Sci.* **13**, 335–360 (1965).
- Guenni, O., Seiter, S. & Figueroa, R. Growth responses of three *Brachiaria* species to light intensity and nitrogen supply. *Trop. Grassl.* **42**, 75 (2008).
- Mooney, H. Resource limitation in plants—An economic analogy. *Annu. Rev. Ecol. Evol. Syst.* **16**, 363–392 (1985).
- Thornley, J. A model to describe the partitioning of photosynthate during vegetative plant growth. *Ann. Bot.* **36**, 419–430 (1972).
- Friel, C. A. & Friesen, M. L. Legumes modulate allocation to rhizobial nitrogen fixation in response to factorial light and nitrogen manipulation. *Front. Plant Sci.* **10**, 1316 (2019).
- Cambui, C. A. *et al.* Patterns of plant biomass partitioning depend on nitrogen source. *PLoS ONE* **6**, e19211 (2011).
- Andrews, M., Raven, J. & Sprent, J. Environmental effects on dry matter partitioning between shoot and root of crop plants: Relations with growth and shoot protein concentration. *Ann. Appl. Biol.* **138**, 57–68 (2001).
- Hosseinzadeh, M., Aliniaiefard, S., Shomali, A. & Didaran, F. Interaction of light intensity and CO₂ concentration alters biomass partitioning in chrysanthemum. *Hortic. Res.* <https://doi.org/10.2478/johr-2021-0015> (2021).
- Guttieri, M. J., Frels, K., Regassa, T., Waters, B. M. & Baenziger, P. S. Variation for nitrogen use efficiency traits in current and historical great plains hard winter wheat. *Euphytica* **213**, 1–18 (2017).

43. Choi, D. *et al.* Photosynthetic characteristics and nitrogen allocation in the black locust (*Robinia pseudoacacia* L.) grown in a FACE system. *Acta Physiol. Plant* **39**, 71 (2017).
44. Díaz-Pérez, J. C. Bell pepper (*Capsicum annuum* L.) crop as affected by shade level: Microenvironment, plant growth, leaf gas exchange, and leaf mineral nutrient concentration. *HortScience* **48**, 175–182 (2013).
45. Santamaria, P., Elia, A. & Serio, F. Effect of solution nitrogen concentration on yield, leaf element content, and water and nitrogen use efficiency of three hydroponically-grown rocket salad genotypes. *J. Plant Nutr.* **25**, 245–258 (2002).
46. Biswas, D. K. & Ma, B.-L. Effect of nitrogen rate and fertilizer nitrogen source on physiology, yield, grain quality, and nitrogen use efficiency in corn. *Can. J. Plant Sci.* **96**, 392–403 (2016).
47. Li, Y., Yang, X., Ren, B., Shen, Q. & Guo, S. Why nitrogen use efficiency decreases under high nitrogen supply in rice (*Oryza sativa* L.) seedlings. *J. Plant Growth Regul.* **31**, 47–52 (2012).
48. Hernández, V., Hellín, P., Fenoll, J. & Flores, P. Interaction of nitrogen and shading on tomato yield and quality. *Sci. Hortic.* **255**, 255–259 (2019).
49. Strasser, B. & Strasser, R. In *Proceedings of the Xth International Photosynthesis Congress*. (Kluwer Academic Publishers).
50. Kalthor, M. S. *et al.* Enhanced salt tolerance and photosynthetic performance: Implication of γ -amino butyric acid application in salt-exposed lettuce (*Lactuca sativa* L.) plants. *Plant Physiol. Biochem.* **130**, 157–172 (2018).
51. Lichtenthaler, H. K. & Wellburn, A. R. (Portland Press Ltd., 1983).
52. Aliniaefard, S. & Van Meeteren, U. 71–76 (*Acta Hort*).
53. Fanourakis, D., Heuvelink, E. & Carvalho, S. M. A comprehensive analysis of the physiological and anatomical components involved in higher water loss rates after leaf development at high humidity. *J. Plant Physiol.* **170**, 890–898 (2013).
54. Karam, F. *et al.* Evapotranspiration, seed yield and water use efficiency of drip irrigated sunflower under full and deficit irrigation conditions. *Agric. Water Manag.* **90**, 213–223 (2007).
55. Dobermann, A. R. Nitrogen use efficiency-state of the art. *Agronomy--Faculty Publications*, 316 (2005).
56. Lei, Z.-Y. *et al.* Enhanced photosynthetic nitrogen use efficiency and increased nitrogen allocation to photosynthetic machinery under cotton domestication. *Photosynth. Res.* **150**, 239–250 (2021).
57. Irigoyen, J., Einerich, D. & Sánchez-Díaz, M. Water stress induced changes in concentrations of proline and total soluble sugars in nodulated alfalfa (*Medicago sativa*) plants. *Physiol. Plant.* **84**, 55–60 (1992).

Author contributions

S.E., S.A., S.D.D. and S.K. conceived and designed research. S.E. and F.D. conducted experiments, material preparation and data collection. S.E., and A.S. analyzed data. A.S., S.E., S.A. and H.M.K. wrote the manuscript. S.A. and H.M.K., A.T. and E.S. revised the manuscript. All authors read and approved the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-022-14163-4>.

Correspondence and requests for materials should be addressed to S.A.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022