



You have downloaded a document from
RE-BUS
repository of the University of Silesia in Katowice

Title: A study of O-H \cdots O hydrogen bonds along various isolines in 2-ethyl-1-hexanol. Temperature or pressure - which parameter controls their behavior?

Author: Barbara Hachuła, Kajetan Koperwas, Roman Wrzalik, Karolina Jurkiewicz, Magdalena Tarnacka, Sebastian Pawlus [i in.]

Citation style: Hachuła Barbara, Koperwas Kajetan, Wrzalik Roman, Jurkiewicz Karolina, Tarnacka Magdalena, Pawlus Sebastian [i in.]. (2022). A study of O-H \cdots O hydrogen bonds along various isolines in 2-ethyl-1-hexanol. Temperature or pressure - which parameter controls their behavior? "Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy" (2022), Pre-proof.
DOI: 10.1016/j.saa.2022.121726



Uznanie autorstwa - Licencja ta pozwala na kopiowanie, zmienianie, rozprowadzanie, przedstawianie i wykonywanie utworu jedynie pod warunkiem oznaczenia autorstwa.



UNIwersYTET ŚLĄSKI
W KATOWICACH



Biblioteka
Uniwersytetu Śląskiego



Ministerstwo Nauki
i Szkolnictwa Wyższego

Journal Pre-proofs

A study of O-H · · O hydrogen bonds along various isolines in 2-ethyl-1-hexanol. Temperature or pressure - which parameter controls their behavior?

Barbara Hachuła, Ewa Kamińska, Kajetan Koperwas, Roman Wrzalik, Karolina Jurkiewicz, Magdalena Tarnacka, Demetrio Scelta, Samuele Fanetti, Sebastian Pawlus, Marian Paluch, Kamil Kamiński

PII: S1386-1425(22)00875-7
DOI: <https://doi.org/10.1016/j.saa.2022.121726>
Reference: SAA 121726



To appear in: *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*

Received Date: 23 May 2022
Revised Date: 18 July 2022
Accepted Date: 5 August 2022

Please cite this article as: B. Hachuła, E. Kamińska, K. Koperwas, R. Wrzalik, K. Jurkiewicz, M. Tarnacka, D. Scelta, S. Fanetti, S. Pawlus, M. Paluch, K. Kamiński, A study of O-H · · O hydrogen bonds along various isolines in 2-ethyl-1-hexanol. Temperature or pressure - which parameter controls their behavior?, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy* (2022), doi: <https://doi.org/10.1016/j.saa.2022.121726>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A study of O-H···O hydrogen bonds along various isolines in 2-ethyl-1-hexanol. Temperature or pressure - which parameter controls their behavior?

Journal Pre-proofs

Barbara Hachula,^{1,*} Ewa Kamińska,² Kajetan Koperwas,³ Roman Wrzalik,³

Karolina Jurkiewicz,³ Magdalena Tarnacka,³ Demetrio Scelta,^{4,5} Samuele Fanetti,^{4,5}

Sebastian Pawlus,^{3,*} Marian Paluch,³ Kamil Kamiński³

¹ *Institute of Chemistry, Faculty of Science and Technology, University of Silesia in Katowice, Szkolna 9, 40-006 Katowice, Poland*

² *Department of Pharmacognosy and Phytochemistry, Faculty of Pharmaceutical Sciences in Sosnowiec, Medical University of Silesia in Katowice, Jagiellońska 4, 41-200 Sosnowiec, Poland*

³ *Institute of Physics, Faculty of Science and Technology, University of Silesia in Katowice, 75 Pułku Piechoty 1, 41-500 Chorzów, Poland*

⁴ *LENS, European Laboratory for Nonlinear Spectroscopy, 50019 Sesto Fiorentino, Italy*

⁵ *ICCOMCNR, Institute of Chemistry of OrganoMetallic Compounds, National Research Council of Italy, I-50019 Sesto Fiorentino, Firenze, Italy*

*Corresponding authors: (B.H.) barbara.hachula@us.edu.pl, (S.P.) sebastian.pawlus@us.edu.pl

Abstract

The nature of hydrogen bond (HB) interactions is still far from being understood despite intense experimental and theoretical studies on this subject carried out by the leading research centers. In this paper, by a combination of unique high-pressure infrared, dielectric and volumetric data, the intramolecular dynamics of hydroxyl moieties (which provides direct information about H-bonds) was studied along various isolines, i.e., isotherms, isobars, isochrones, and isochores, in a simple monohydroxy alcohol (2-ethyl-1-hexanol). This allowed us to discover that the temperature controls the intermolecular hydrogen bonds, which then affect the intramolecular dynamics of O-H units. Although the role of density fluctuations gets stronger as temperature rises. We also demonstrated a clear connection between the intra- and intermolecular dynamics of the associating liquid at high pressure. The data reported herein open a new perspective to explore this important aspect of the glass transition phenomenon and understand H-bonding interactions at varying thermodynamic conditions.

Keywords: Hydrogen bond, Temperature, Pressure, Isolines, Intramolecular dynamics.

Introduction

Pressure (p) is a fundamental thermodynamic variable that affects the distances between molecules (molecular packing), depth, and shape of the potential well describing intermolecular

Journal Pre-proofs

the studied matter. As an example, one can mention that previous studies performed at varying thermodynamic conditions demonstrated that it is possible to shorten reaction times and synthesize materials of unique properties or geometry not attainable at ambient pressure [4,5]. What is more, the experiments allowed inducing quite an extraordinary variation in physical features of the compressed matter. In this context, it is worth reminding of the most spectacular ones, such as gas metallization, the transition from insulator to superconductor, ionization of water, or ammonia that occur at high compression [6,7,8].

Furthermore, the studies at elevated pressure were the key to better understanding the essence of H-bonds (HBs) – one of the most fundamental and crucial electrostatic interactions occurring in nature. Briefly, one can recall that this type of molecular interactions i) underlies all-important processes undergoing in the living organisms, as well as the self-assembly phenomenon, ii) determines the biological activity of proteins, or iii) the peculiar behavior of water and other associating liquids. Herein, it should be added that extensive high-pressure (HP) studies on the H-bonded liquids clearly indicated that their dynamical and thermodynamical properties, such as pressure evolution of the relaxation times, diffusion, pressure coefficient of the glass transition temperature, activation volume, bulk moduli, and thermal expansion coefficient, are much different with respect to the van der Waals systems [9,10,11,12]. What is more, it was experimentally observed that the thermodynamic scaling (TS) - a hotly debated concept that links dynamical properties (relaxation times) with the exponent describing intermolecular repulsion between molecules, is usually not satisfied for liquids forming HBs. However, it is not a rule, and the TS seems to depend on the strength of H-bonds, variation in the architecture of nanoassociates, range of p , etc. A good illustration of a deviation from the TS rule is 2-ethyl-1-hexanol (2E1H) - a simple monohydroxy alcohol, for which the scaling exponent is not constant above 500 MPa [13,14,15]. Interestingly, this extraordinary behavior was linked with the change of the H-bonding pattern in the sample. Nevertheless, it should be emphasized that the mentioned example and others described above are devoted to investigating the behavior of associated liquids at high p , not HBs themselves. Over recent years, using indirect experimental methods was the most common way and serious drawback of probing HBs under high p . As a consequence of that, there are different points of


Journal Pre-proofs

view on the impact of high compression on the strength and population of this kind of bonds at different temperature and pressure conditions. Based on dielectric data, it was postulated that HBs are being broken, and their population decreases with the increase in p [16,17,18]. On the other hand, the simulation data obtained for various systems indicated no effect of pressure or the enhancement of these interactions with compression [19,20]. Finally, Fourier-Transform Infrared (FTIR) and Raman studies revealed a monotonic increase in the strength of HBs with p [21,22]. In fact, the former technique is the most sensitive and fully dedicated to probing the properties of H-bonded associates at varying thermodynamics conditions. It is worth emphasizing that the length, as well as the angle between the proton donor (X-H) and acceptor (Y) groups, influence the frequency of the stretching vibration of the hydroxyl units (ν_{O-H}) and the shape of the related band [23]. Therefore, any variation in the spectroscopic parameters, namely frequency, the full width at half maximum (FWHM), or the integral intensity of the ν_{X-H} band, may provide unique information on the real effect of compression on the behavior of HBs. Nevertheless, due to the high intensity of the mentioned band in FTIR experiments under high pressure, performed usually in diamond anvil cells (DAC) [24,25], determination of the credible parameters describing these specific interactions is quite difficult.

In this paper, we applied the vibrational spectroscopy technique combined with the DAC to measure FTIR spectra of 2E1H at high p . To obtain samples of uniform thickness and absorbance value, DAC was filled with the tested alcohol and very thin polyethylene foil. The collected data yielded fundamental information about the impact of pressure on HBs. Furthermore, by combining dielectric and volume data with the ones obtained from FTIR investigations, we were able to make credible conclusions about the behavior of H-bonding interactions along different isolines, namely isotherms (temperature, $T=\text{const.}$), isobars ($p=\text{const.}$), isochrones (relaxation times, $\tau_\alpha=\text{const.}$) and isochores (volume, $V=\text{const.}$). Finally, for the first time, we evaluated the effect of density and temperature on the energy/strength of HBs.

Materials and Methods

2-ethyl-1-hexanol (2E1H) was purchased from Sigma-Aldrich (purity > 99%). Before use, the alcohol was dried under a stream of liquid nitrogen. The chemical structure of the studied compound is presented in **Scheme 1**.



Scheme 1. The scheme of chemical structure of the H-bonded 2E1H under investigation.

High-pressure (HP) Fourier Transform Infrared (FTIR) spectroscopy

FTIR absorption spectra were recorded using a Bruker-IFS 120 HR spectrometer suitably modified for experiments in DAC, with an instrumental resolution set to 1 cm^{-1} [26]. The sample pressure was measured by the ruby fluorescence method using a few mW of a 532 nm laser line from a doubled Nd:YAG laser source. 2E1H was loaded into a membrane diamond anvil cell (MDAC) equipped with type IIa synthetic diamonds from Almax easyLab Inc., together with a small ruby chip as a pressure gauge. A thin pellet of polyethylene was used to reduce the sample thickness and to have an O-H absorbance from 0.1 to 1. All samples were laterally contained by steel gaskets, drilled to have an initial sample diameter of $150\text{ }\mu\text{m}$ and a thickness of about $50\text{ }\mu\text{m}$. High-temperature runs were carried out using resistive heaters, and the temperature was measured by means of a K-type thermocouple with an accuracy of $\pm 0.1\text{ K}$.

It should be added that the analysis of spectral parameters (the peak position, FWHM, integrated intensity) for the bands measured at HP was a challenging task due to the interference fringes disturbing the data. This well-known and common effect occurring in the HP experiments results from multiple internal reflections inside the film with two nearly parallel surfaces leading to signatures of constructive and destructive interference. For this reason, the O-H peak positions were determined using the Fityk software (a data processing and nonlinear curve fitting program; version: 1.3.1.; url: <https://fityk.nieto.pl/>) by applying VoightA peak profile functions adjusting the intensity and the width of the fitting curves [27]. It should also be noted that the band component analysis of the C-H stretching region, occurring at ca. $3050\text{--}2700\text{ cm}^{-1}$, was excluded from the deconvolution of the O-H band due to the saturation of C-H

bands. Moreover, this spectral region was strongly disturbed by absorption signals originating from the polyethylene spacers used to maintain a reliable absorbance value of the $\nu_{\text{O-H}}$ band. Moreover, the fitting of poorly resolved and broad O-H bands with saturated components of C-H bands led to a large error, only worsening the final deconvolution result. One can stress that the sinusoidal background was determined from the particular fringes observed in each IR spectrum individually and then was fitted to the measured data in order to facilitate the interpretation of the IR spectra without losing any spectral information. The experimental error for the band position in the HP FTIR spectra was estimated to be ca. $\pm 5 \text{ cm}^{-1}$, although for higher p ($>3 \text{ GPa}$), it was ca. $\pm 10 \text{ cm}^{-1}$ since the O-H band profile overlapped the C-H band region with increasing compression. Illustrative plots and details for the fitted spectra are shown in **Figure S1** in the **Supplementary Material** (SM).

Fourier Transform Infrared (FTIR) spectroscopy

FTIR spectra measurements were also performed at atmospheric pressure in the temperature range 123–293 K using a Nicolet iS50 FTIR spectrometer (Thermo Scientific) with a Linkam THMS 600 heating/cooling stage (Linkam Scientific Instruments Ltd., Surrey, U.K.) mounted inside the sample stage of FTIR spectrometer. A sample of the studied alcohol was placed between two CaF_2 windows, separated by a $15 \text{ }\mu\text{m}$ thick spacer, which ensured the uniform thickness of the sample and warranted the constant geometry of the system. The spectra were recorded in the range of $4000\text{--}400 \text{ cm}^{-1}$ at a resolution of 4 cm^{-1} . For every spectrum, 16 scans were averaged. The temperature stabilization accuracy was equal to $\pm 0.1 \text{ K}$.

Results and discussion

Despite the universal nature and ubiquity of HBs, so far, it has not been experimentally shown which thermodynamic parameter (temperature or pressure) has the most significant impact on their dynamics, pattern, and strength. Furthermore, it was not proven if there is any connection between the behavior of HBs and dynamics of associating liquids - an assumption made a priori by leading research groups. To gain an insight into these fundamental problems, we have carried out HP FTIR measurements on 2E1H, model alcohol, for which dielectric as well as volumetric data obtained at varying thermodynamic conditions are available in the literature [15, 28]. IR spectra of 2E1H were collected upon compression runs up to $p \sim 6 \text{ GPa}$

at four different $T = 293, 308, 323,$ and 348 K. A separate IR experiment was performed at ambient pressure as a reference to track the changes in the behavior of HBs as a function of T . Herein, it must be stressed that to avoid the saturation effect of the stretching vibration of hydroxyl units at elevated p , a very thin polyethylene foil has been introduced between diamonds in an anvil cell. As a consequence, the dynamics and the spectral parameters originating from the vibrations of C-H units were highly affected by those coming from polyethylene. Therefore, this spectral range was excluded from further analysis. What is more, the disturbance of the C-H bands also had a negative effect on the deconvolution of the ν_{O-H} bands, as this absorption signal moved towards the C-H bands with increasing pressure. Another problem was the interference, which introduced a disturbance in the shape (the subtle structure) of the ν_{O-H} band, influencing its spectral parameters (the peak position, FWHM, and integrated intensity). Data analysis revealed that the O-H band frequency is the least affected parameter due to interference and saturation effects. Hence, the evolution of this parameter as a function of varying thermodynamics conditions will be analyzed in more detail in this paper.

The temperature and pressure evolution of the IR spectra in the region of $3600-2500$ cm^{-1} is shown in **Figure 1** and **Figure S2** in the SM file. As can be seen, there is a broad absorption signal between 3600 and 3050 cm^{-1} assigned to the stretching vibration of the O-H group, whereas the lower frequency band ($3050-2700$ cm^{-1}) corresponds to the C-H stretching vibrations. As mentioned above, the observed C-H bands were not taken for further analysis due to the saturation effect and overlapping with the vibrations originating from the polyethylene spacers.




Figure 1. FTIR spectra of 2E1H measured at (a) $T = 293$ - 123 K and $p = 0.1$ MPa and b) $T = 293$ K and $p = 400$ - 6170 MPa in the frequency region of 3600 - 2500 cm^{-1} .

At first, we analyzed the IR spectra measured during isobaric cooling. As shown in **Figure 1(a)**, the stretching frequencies of the covalent O-H bonds are red-shifted (shifted to lower wavenumbers) with decreasing T due to the elongation of the O-H bond lengths (the strengthening of the H-bond interactions). This spectral modification, connected with a change in the X-H peak position of the H-bond donor molecule (where X is either O, Cl, C, N, or F), accompanying different external factors, i.e., T or p , is an indicator of the strength of HBs. It is a well-known correlation that the larger the red-shift of the X-H band position, the stronger the hydrogen bond [29]. It should be mentioned that the high-precision *ab initio* methods of calculations applied recently on water dimer showed that the H-bond ($\text{H}\cdots\text{O}$) becomes weaker with the contraction of H-bond length and $\text{O}\cdots\text{O}$ distance, which is attributed to the exchange repulsion of electrons [Refs. 30, 31, 32].

Next, the effect of isothermal compression on the position of the band was considered. As can be observed in **Figure 1(b)**, with increasing pressure, the $\nu_{\text{O-H}}$ signal red-shifts and shows the monotonic change in the frequency with no discontinuities or changes in slope indicative of a phase change or crystallization. An analogous spectroscopic effect of lowering the frequency of the O-H band with the compression was observed for other IR spectra recorded

as a function of p at 308, 323, and 348 K (see **Figure S2** in the SM file). Thus, the studied alcohol exhibits a typical pressure-induced behavior, i.e., the red-shift of the IR active O-H vibration with increasing pressure suggesting the strengthening of intermolecular HBs. A similar scenario was detected in previous HP studies on methanol, ethanol, or ethylene glycol [22,33].

At first glance, one can see that the pressure had a greater effect on the position of the O-H band than temperature (**Figure 1**). The O-H peak frequency for the temperature-dependent IR spectra moved to the red regime by 62 cm^{-1} (from 293 to 123 K) compared with the 135 cm^{-1} shift obtained during the isothermal compression at 293 K (from 0.1 to 6170 MPa) (for higher temperatures: $\Delta\nu=121\text{ cm}^{-1}/6140\text{ MPa}$ at 308 K; $\Delta\nu=166\text{ cm}^{-1}/6130\text{ MPa}$ at 323 K; $\Delta\nu=145\text{ cm}^{-1}/6290\text{ MPa}$ at 348 K). The much longer elongation of the O-H bond for an isobar compared to that of an isotherm indicates stronger H-bonding interactions occurring at high p than at 123 K. This experimental finding suggests that isothermal compression (density) has a greater effect on the H-bond dynamics than isobaric cooling (combined effect of temperature and density fluctuations). However, this effect may be apparent since it is very difficult to compare directly pressurization effects (density packing) with a thermal variation. Hence to gain deeper insight into this problem, further analysis was performed.




Figure 2. FTIR spectra of 2E1H in the frequency region of 3600-2600 cm^{-1} obtained after the deconvolution process, using sinusoidal and VoigtA formulas, at (a) 293 K as a function of pressure, (b) constant specific volume ($V_s = 0.88 \text{ cm}^3 \cdot \text{g}^{-1}$), and (c) at isochronal conditions determined from the pressure dependences of the glass transition temperature for the alcohol under investigation. The spectra have been shifted for clarity.

Considering dielectric and volumetric data measured for 2E1H at various thermodynamic conditions (given in Refs. [15] and [28]), we decided to take a unique opportunity to monitor the behavior of H-bonding interactions along different isolines, namely isotherms ($T=\text{const}$), isobar ($p=\text{const}$), isochrones ($\tau_\alpha=\text{const}$) and isochores ($V=\text{const}$). Representative FTIR spectra at the following constant condition of (a) temperature, (b) specific volume ($V_s=0.88 \text{ cm}^3 \cdot \text{g}^{-1}$) and (c) relaxation times (viscosity) are shown in **Figure 2**. To determine the values of V for the given T and p conditions the Tait equation of state was used [15]:

$$V(T, p) = V_0 \exp(a_0 T) \left\{ 1 - C \ln \left[1 + \frac{p}{b_0 \exp(-b_1 T)} \right] \right\} \quad (1)$$

where: $V_0 = 1.1729 \pm 10^{-4} \text{ cm}^3 \cdot \text{g}^{-1}$, $a_0 = (8.873 \pm 0.008) \cdot 10^{-4} \text{ C}^{-1}$, $b_0 = 137.5 \pm 0.1 \text{ MPa}$, $b_1 = (6.19 \pm 0.02) \cdot 10^{-3} \text{ C}^{-1}$, and $C = 0.0894$ are constants taken from Ref. [15]. It should be noted that the temperature and pressure dependence of V_s was determined experimentally up to 200 MPa (under the given T and n conditions). Hence to obtain $V(T, p)$ at much higher pressures, corresponding to those where FTIR spectra were collected, we calculated them from the Tait equation of state (**Eq. 1**). Thus, by combining the isothermal and isobaric FTIR data with the results of volumetric measurements modeled using the Tait formula, the correlation between the V_s s and the O-H peak positions for any conditions of T and p were estimated. IR spectra measured at isochoric point ($V_s = 0.88 \text{ cm}^3 \cdot \text{g}^{-1}$) are shown in **Figure 2(b)**. To monitor the evolution of the O-H band at isochronal conditions ($\tau_\alpha = \text{const}$), we used dielectric data reported in Ref. [28]. In this paper, a pressure evolution of the glass transition temperature (T_g) was evaluated (please see **Figure S3**). Taking into account that vitrification occurs at constant structural relaxation times (usually, it is defined as temperature or pressure at which $\tau_\alpha = 100 \text{ s}$), one can easily determine isochrones along various thermodynamic conditions. Herein, it should be noted that the pressure evolution of the T_g for the studied alcohol was obtained up to 1570 MPa. Hence, to evaluate T_g vs. p_g variation at much higher pressures, comparable to those applied during FTIR experiments, this dependency was further modeled by the well-known empirical Anderson–Anderson formula (see **Figure S3** in the SM file) [34]. Having all these analyses done, one can follow the shift of the O-H peak position under the isobaric, isochoric, and isochronal conditions in **Figure 2**. At first sight, it is visible that there is a red shift of the $\nu_{\text{O-H}}$ band, the blue-shift one, and the non-monotonic change of this band frequency for isobaric, isochoric, and isochronal conditions, respectively. Nevertheless, to explore these spectral changes in detail, the determined O-H peak positions along different isolines were plotted vs. p and T in **Figure 3**.




Figure 3. The variation of the O-H band frequency along a) isotherm at 323 K, b) isobar at 0.1 MPa, c) isochore at $V_s = 0.8, 0.88$ and $1.0 \text{ cm}^3 \cdot \text{g}^{-1}$, and d) isochrone ($\tau_\alpha = 100 \text{ s}$) along $T_g(p_g)$.

Examination of the plots presented in **Figure 3** revealed: (a) the red shift of the $\nu_{\text{O-H}}$ band frequency from 3329 cm^{-1} (460 MPa) to 3189 cm^{-1} (6130 MPa) during isothermal compression at 323 K; (b) the reduction of the $\nu_{\text{O-H}}$ band frequency from 3331 cm^{-1} (293 K) to 3269 cm^{-1} (123 K) during cooling at ambient pressure; (c) the blue-shift of the $\nu_{\text{O-H}}$ band frequency from 3231 cm^{-1} (293 K; 2110 MPa) to 3268 cm^{-1} (348 K, 2350 MPa) with an increase of temperature at constant volume ($V_s = 0.88 \text{ cm}^3 \cdot \text{g}^{-1}$); and (d) a drastic decrease in the $\nu_{\text{O-H}}$ band frequency from 3277 to 3214 cm^{-1} along $T_g(p_g)$ line from $T = 143 \text{ K}$ to $T = 293 \text{ K}$, then the non-monotonically rise of this parameter up to 348 K (3241 cm^{-1}). It was also observed that for higher specific volume ($V_s = 1.00 \text{ cm}^3 \cdot \text{g}^{-1}$), a smaller change in the O-H peak position for 293 K and 348 K ($\Delta\nu \sim 27 \text{ cm}^{-1}$) occurs than for the lower V_s : 0.80 and $0.88 \text{ cm}^3 \cdot \text{g}^{-1}$ ($\Delta\nu \sim 36 \text{ cm}^{-1}$). This observation may suggest that for the higher compression, the temperature fluctuations become more important in controlling the strength of HBs.

However, to quantify this observation, as a final point of our discussion, we analyzed the pressure dependence of the ratio of the isochoric and isobaric activation energies, E_V/E_p (where $E_V = R \partial \log(x) / \partial (1/T)|_V$, $E_p = R \partial \log(x) / \partial (1/T)|_p$), which is a fundamental tool to

verify whether the temperature or density fluctuations govern the x parameter, which refers to the dynamics. Herein, it should be noted that this kind of analysis was commonly adopted to determine the influence of thermal and molecular packing on the structural and segmental dynamics in low- and high-molecular-weight liquids [9]. Briefly, one can stress that as this ratio equals 0 or 1, the mobility is governed solely either by volume or temperature, respectively. The schematic representation of the discussed scenarios is presented in **Figure 4(a)** for the ν_{O-H} band. The blue arrow depicts the situation, in which the volume controls the shift of ν_{O-H} band. In this case, the change in the temperature at isochoric conditions does not affect the ν_{O-H} position. Consequently, one can observe the horizontal $\nu_{O-H}(1/T)$ dependence, ($E_V/E_p = 0$). Contrary, the red arrow reflects the scenario, in which exclusively the temperature controls HBs dynamics. In this situation, the isochronal $\nu_{O-H}(1/T)$ overlaps the isobaric one ($E_V/E_p = 1$). Hence, to find out which scenario prevails, we plotted the pressure dependencies of the ν_{O-H} band position at four different isotherms and described them by the exponential functions (see solid lines in **Figure 4(b)**). This procedure allowed us to estimate ν_{O-H} at any pressure at $T=293, 308, 323,$ and 348 K. Subsequently, employing the equation of state, we predicted the evolution of pressure, when temperature increases at $V=\text{const}$. Hence, it was possible to estimate the position of the O-H band along the isochore. We performed these computations starting with a density corresponding to the point measured at $T=293$ K and $p=0.1$ MPa, see **Figure 4(a)**. Interestingly, the obtained isochoric dependency of the ν_{O-H} band almost perfectly overlaps the experimental points measured at ambient pressure (isobar $p=0.1$ MPa). It means the E_V/E_p ratio should be close to the unity. In fact, the calculated E_V/E_p equals 0.97. From this analysis, it becomes clear that temperature is the main parameter controlling the intermolecular hydrogen bonds that influence the intramolecular dynamics of O-H moieties.




Figure 4. Panel (a) depicts an evolution of the logarithm of the O-H band frequency (ν) as a function of the inverse temperature ($1000/T$). Arrows schematically present the behavior of O-H band frequencies corresponding to the situations in which the dynamics of HBs is governed by the volume (blue arrow, $E_V/E_p = 0$), and the temperature (red arrow, $E_V/E_p = 1$). The black arrow illustrates the results of our calculations. Panel (b) presents O-H band frequencies (ν) as a function of p . The pressure dependence of E_V/E_p is shown in the inset of panel (b).

Further deeper analysis of the data revealed that the E_V/E_p parameter changes non-monotonically with the pressure for all isotherms up to the values close to the unity (from 0.86-0.97 at 0.1 MPa to 0.96-0.98 at 5800 MPa). Moreover, the characteristic minimum occurs at ca. 600 MPa. Interestingly, such pressure corresponds to the upper limit of p , for which the TS rule for 2E1H is obeyed [13]. One can add that below and above this p , one can scale the relaxation times measured at different thermodynamic conditions with constant and τ_α -dependent scaling exponent [13], respectively. What is more, the E_V/E_p approaches unity with the compression. It means that temperature plays a dominant role in controlling the intramolecular dynamics of

O-H units. Moreover, panel (b) in **Figure 4** revealed that as the temperature of the isotherm increases, the E_V/E_p decreases, suggesting the increasing role of density. The other interesting result is that the weaker HBs (the higher values of the O-H band frequency) are more sensitive to changes in the volume than the stronger HBs. Finally, it should be noted that the obtained values of E_V/E_p ratio are close to those determined from dielectric measurements for the Debye relaxation ($E_V/E_p = 0.76$), which is regarded as a manifestation of the dynamics of HBs [15]. Therefore, the analysis of the O-H stretch band position at high p clearly indicates that there is a close correlation between the dynamics of the Debye relaxation observed in dielectric loss spectra and the intramolecular dynamics of O-H units or the strength of HBs. What is more, the value of E_V/E_p obtained herein is also comparable to those estimated for the structural relaxation in other H-bonded liquids with more than one hydroxyl group, e.g., glycerol ($E_V/E_p = 0.97$), sorbitol ($E_V/E_p = 0.87$), glycol derivatives (propylene glycol: $E_V/E_p = 0.88$, dipropylene glycol: $E_V/E_p = 0.86$, and tripropylene glycol: $E_V/E_p = 0.83$) [15]. Herein it should be noted that the value of this parameter for the non-associated materials, such as ortho-terphenyl ($E_V/E_p = 0.55$), salol ($E_V/E_p = 0.43$), propylene carbonate ($E_V/E_p = 0.64$), and phenylphthalindimethylether ($E_V/E_p = 0.53$) is much lower [15]. These simple examples indicate that so high values of E_V/E_p for liquids associating via HBs are connected to the intramolecular dynamics of O-H units.

Conclusions

In this paper, we considered a fundamental question from the perspective of physics, chemistry, and biology: which parameter does determine the properties of HBs – temperature or density (at given thermodynamic conditions)? For this purpose, we have analyzed the variation of the strength of HBs in 2E1H, directly through the shift of the O-H band position in the FTIR spectra, upon T and p variation. Then, the obtained FTIR results were combined with the dielectric and volumetric data. In this way, we were able to describe the temperature and pressure dependence of H-bonding energy along various isolines. What is more, we calculated the E_V/E_p parameter, which provides a direct insight into the impact of density and temperature on the dynamics of an associating system. Adopting this kind of analysis for the O-H band frequencies of simple alcohol – 2E1H allowed us to demonstrate that T controls the intermolecular hydrogen bonds affecting the intramolecular dynamics of O-H moieties. However, it must be added that the impact of density grows together with increasing T and

weakening of H-bonding interactions. It was also found that the temperature plays a dominant role in the case of stronger HBs, while for the weaker ones, the density effect on the association process via HBs gets stronger. Thus, the presented studies enable unprecedented insight into the behavior of HBs and the dynamics of associating liquids. What is more, they open completely new pathways to explore HBs in various kinds of materials and an even better understanding of these fundamental electrostatic interactions.

Journal Pre-proofs

AUTHOR INFORMATION

NOTES. The authors declare no competing financial interests.

SUPPLEMENTARY MATERIAL. The Supplementary Material file is available free of charge at <http://pubs.acs.org>. It contains additional figures, including the details of the deconvolution of the $\nu_{\text{O-H}}$ band, the pressure-dependent FTIR spectra measured at $T=308$, 323, and 348 K, and the pressure dependence of the glass transition temperature for 2E1H.

ACKNOWLEDGMENT. The research leading to these results has received funding from LASERLABEUROPE (Grant agreement No. 654148, European Union's Horizon 2020 research and innovation programme; Proposal No. LENS002738). B.H, K.J., K.K., and S.P. are thankful for the Polish National Science Centre's financial support within the OPUS project (Dec. no UMO-2019/35/B/ST3/02670).

REFERENCES

- ¹ S. Fanetti, M. Citroni, K. Dziubek, M. M. Nobrega, R. Bini, The role of H-bond in the high-pressure chemistry of model molecules, *J. Phys. Condens. Matter* 30 (2018) 094001. <https://doi.org/10.1088/1361-648X/aaa8cf>.
- ² S. K. Sikka, S. M. Sharma, The hydrogen bond under pressure, *Phase Transit.* 81 (2008) 907–934. <https://doi.org/10.1080/01411590802098864>.
- ³ G. Nielsen, H. Schwalbe, Hydrogen bonds under pressure, *Nat. Chem.* 4 (2012) 693–695. <https://doi.org/10.1038/nchem.1443>.
- ⁴ M. Citroni, M. Ceppatelli, R. Bini, V. Schettino, Laser-induced selectivity for dimerization versus polymerization of butadiene under pressure, *Science* 295 (2002) 2058–2060. <https://doi.org/10.1126/science.1068451>.
- ⁵ D. Chelazzi, M. Ceppatelli, M. Santoro, R. Bini, V. Schettino, High-pressure synthesis of crystalline polyethylene using optical catalysis, *Nat. Mater.* 3 (2004) 470–475. <https://doi.org/10.1038/nmat1147>.
- ⁶ P. F. McMillan, New materials from high-pressure experiments, *Nat. Mater.* 1 (2002) 19–25. <https://doi.org/10.1038/nmat716>.
- ⁷ L. Zhang, Y. Wang, J. Lv, Y. Ma, Materials discovery at high pressures, *Nat. Rev. Mater.* 2 (2017) 17005. <https://doi.org/10.1038/natrevmats.2017.5>.
- ⁸ C. Buzea, K. Robbie, Assembling the puzzle of superconducting elements: a review, *Supercond. Sci. Technol.* 18 (2005) R1–R8. <https://doi.org/10.1088/0953-2048/18/1/R01>.
- ⁹ G. Floudas, M. Paluch, A. Grzybowski, K. Ngai, *Molecular dynamics of glass-Forming systems: Effects of pressure* (Springer-Verlag, Berlin, Heidelberg, 2011).

- ¹⁰ C. M. Roland, M. Paluch, T. Pakula, R. Casalini, Volume and temperature as control parameters for the dielectric a relaxation of polymers and molecular glass formers, *J. Phil. Mag.* 84 (2004) 1573–1581. <https://doi.org/10.1080/14786430310001644350>.
- ¹¹ T. Atake, C. A. Angell, Pressure dependence of the glass transition temperature in molecular liquids and plastic crystals, *J. Phys. Chem.* 8 (1979) 3218–3223. <https://doi.org/10.1021/j100488a007>.
- ¹² H. S. Ashbaugh, Pressure-Induced Changes in the Dynamics of Hydrogen Bonding in Liquid Methanol, *J. Chem. Phys.* 117 (2002) 10101. <https://doi.org/10.1063/1.1619938>.
- ¹³ S. Pawlus, M. Paluch, A. Grzybowski, Communication: Thermodynamic scaling of the Debye process in primary alcohols, *J. Chem. Phys.* 134 (2011) 041103. <https://doi.org/10.1063/1.3540636>.
- ¹⁴ A. Reiser, G. Kasper, C. Gainaru, R. Böhmer, Communications: High-pressure dielectric scaling study of a monohydroxy alcohol, *J. Chem. Phys.*, 132 (2010) 181101. <https://doi.org/10.1063/1.3421555>.
- ¹⁵ D. Fragiadakis, C. M. Roland, R. Casalini, Insights on the origin of the Debye process in monoalcohols from dielectric spectroscopy under extreme pressure conditions, *J. Chem. Phys.* 132 (2010) 144505. <https://doi.org/10.1063/1.3374820>.
- ¹⁶ C. M. Roland, S. Hensel-Bielowka, M. Paluch, R. Casalini, Supercooled dynamics of glass-forming liquids and polymers under hydrostatic pressure, *Rep. Prog. Phys.* 68 (2005) 1405. <https://doi.org/10.1088/0034-4885/68/6/R03>
- ¹⁷ K. L. Ngai, S. Capaccioli, Impact of the application of pressure on the fundamental understanding of glass transition, *J. Phys.: Condens. Matter* 20 (2008) 244101. <https://doi.org/10.1088/0953-8984/20/24/244101>.
- ¹⁸ C. M. Roland, R. Casalini, M. Paluch, Isochronal temperature–pressure superpositioning of the α -relaxation in type-A glass formers, *Chem. Phys. Lett.* 367 (2003) 259–264. [https://doi.org/10.1016/S0009-2614\(02\)01655-X](https://doi.org/10.1016/S0009-2614(02)01655-X).
- ¹⁹ W. L. Jorgensen, M. Ibrahim, Pressure dependence of hydrogen bonding in liquid methanol, *J. Am. Chem. Soc.* (1982) 373–378. <https://doi.org/10.1021/ja00366a001>.
- ²⁰ L. J. Root and B. J. Berne, Effect of pressure on hydrogen bonding in glycerol: A molecular dynamics investigation, *J. Chem. Phys.* 107 (1997) 4350–4357. <https://doi.org/10.1063/1.474776>.
- ²¹ A. Arencibia, M. Taravillo, F. J. Perez, J. Nunez, V. G. Baonza, Effect of pressure on hydrogen bonding in liquid methanol, *Phys. Rev. Lett.* 89 (2002) 195504. <https://doi.org/10.1103/PhysRevLett.89.195504>.
- ²² J. F. Mammone, S. K Sharma, M. Nicol, Raman spectra of methanol and ethanol at pressures up to 100 Kbar, *J. Phys. Chem.* 84 (1980) 3130–3134. <https://doi.org/10.1021/j100460a032>.
- ²³ P. Griffiths, J. A. de Haseth, *Fourier Transform Infrared Spectrometry* (2nd ed.), Wiley-Blackwell, 2007.
- ²⁴ P. A. Guńka, A. Olejniczak, S. Fanetti, R. Bini, I. E. Collings, V. Svitlyk, K. F. Dziubek, Crystal structure and non-hydrostatic stress-induced phase transition of urotropine under high pressure, *Chem. Eur. J.* 27 (2021) 1094–1102. <https://doi.org/10.1002/chem.202003928>.
- ²⁵ W. Dzwolak, M. Kato and Y. Taniguchi, Fourier transform infrared spectroscopy in high-pressure studies on proteins, *Biochimica et Biophysica Acta (BBA) - Protein Structure and Molecular Enzymology.* 1595 (2002) 131–144. [https://doi.org/10.1016/s0167-4838\(01\)00340-5](https://doi.org/10.1016/s0167-4838(01)00340-5).
- ²⁶ R. Bini, R. Ballerini, G. Prates, H. Jodl, Experimental setup for Fourier transform infrared spectroscopy studies in condensed matter at high pressure and low temperatures, *J. Rev. Sci. Instrum.* 68 (1997) 3154–3160. <https://doi.org/10.1063/1.1148261>.
- ²⁷ M. Wojdyr, Fityk: a general-purpose peak fitting program, *J. Appl. Cryst.* 43 (2010) 1126–1128. <https://doi.org/10.1107/S0021889810030499>.
- ²⁸ S. Pawlus, M. Paluch, M. Dzida, Molecular dynamics changes induced by hydrostatic pressure in a supercooled primary alcohol, *J. Phys. Chem. Lett.* 1 (2010) 3249–3253. <https://doi.org/10.1021/jz101288v>.
- ²⁹ G. A. Jeffrey, *An Introduction to Hydrogen Bonding*, Oxford University Press, 1997.
- ³⁰ D. Li, Z. Zhang, W. Jiang, Y. Zhu, Y. Gao, Z. Wang, Uncooperative effect of hydrogen bond on water dimer, *Chin. Phys. Lett.* 38 (2021) 013101. <https://doi.org/10.1088/0256-307X/38/1/013101>.
- ³¹ J. C. Howard, J. L. Gray, A. J. Hardwick, L. T. Nguyen, G. S. Tschumper, Getting down to the fundamentals of hydrogen bonding: Anharmonic vibrational frequencies of (HF)₂ and (H₂O)₂ from ab initio electronic structure computations, *J. Chem. Theory Comput.* 10 (2014) 5426–5435. <https://doi.org/10.1021/ct500860v>.
- ³² J. R. Lane, CCSDTQ optimized geometry of water dimer, *J. Chem. Theory Comput.* 9 (2013) 316–323. <https://doi.org/10.1021/ct300832f>.
- ³³ Y. Song, Novel pressure-induced molecular transformations probed by in situ vibrational spectroscopy, *Applications of Molecular Spectroscopy to Current Research in the Chemical and Biological Sciences*, Edited by Mark T. Stauffer, 2016.
- ³⁴ M. Tarnacka, W. K. Kipnusu, E. Kaminska, S. Pawlus, K. Kaminski, M. Paluch, The peculiar behavior of the molecular dynamics of a glass-forming liquid confined in native porous materials – the role of negative pressure, *Phys. Chem. Chem. Phys.* 18 (2016) 23709–23714. <https://doi.org/10.1039/C6CP03923E>.

Highlights:

Journal Pre-proofs


T and p dependence of H-bonding energy along various isolines was studied.

A phase diagram of 2E1H, identifying isolines in the T - p plane, was constructed.

T controls the intramolecular dynamics of O-H units in 2E1H.

The impact of density fluctuation grows with increasing T and weakening of HBs.

The intramolecular dynamics of OH units



Declaration of interests

- The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.
- The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

CRediT author statement:

B. Hachuła: Writing – original draft, Writing - Review & Editing, Investigation, Formal analysis, Visualization, Methodology, Conceptualization.

E. Kamińska: Writing – original draft.

K. Koperwas: Formal Analysis.

R. Wrzalik: Formal Analysis.

K. Jurkiewicz: Visualization.

M. Tarnacka: Investigation.

D. Scelta: Investigation.

S. Fanetti: Investigation.

S. Pawlus: Investigation, Funding acquisition.

M. Paluch: Supervision.

K. Kamiński: Conceptualization, Supervision.