



You have downloaded a document from
RE-BUS
repository of the University of Silesia in Katowice

Title: Complete electroweak two-loop corrections to Z boson production and decay

Author: Ievgen Dubovyk, Ayres Freitas, Janusz Gluza, Tord Riemann, Johann Usovitsch

Citation style: Dubovyk Ievgen, Freitas Ayres, Gluza Janusz, Riemann Tord, Usovitsch Johann. (2018). Complete electroweak two-loop corrections to Z boson production and decay. "Physics Letters, B" (Vol. 783 (2018), s. 86-94), doi 10.1016/j.physletb.2018.06.037



Uznanie autorstwa - Licencja ta pozwala na kopiowanie, zmienianie, rozprowadzanie, przedstawianie i wykonywanie utworu jedynie pod warunkiem oznaczenia autorstwa.



UNIWERSYTET ŚLĄSKI
W KATOWICACH



Biblioteka
Uniwersytetu Śląskiego



Ministerstwo Nauki
i Szkolnictwa Wyższego



Complete electroweak two-loop corrections to Z boson production and decay



Ievgen Dubovyk^a, Ayres Freitas^{b,*}, Janusz Gluza^c, Tord Riemann^c, Johann Usovitsch^d

^a II. Institut für Theoretische Physik, Universität Hamburg, 22761 Hamburg, Germany

^b Department of Physics & Astronomy, University of Pittsburgh, Pittsburgh, PA 15260, USA

^c Institute of Physics, University of Silesia, 40-007 Katowice, Poland

^d Trinity College Dublin (TCD) – School of Mathematics, Dublin 2, Ireland

ARTICLE INFO

Article history:

Received 18 May 2018

Accepted 18 June 2018

Available online 21 June 2018

Editor: A. Ringwald

ABSTRACT

This article presents results for the last unknown two-loop contributions to the Z-boson partial widths and Z-peak cross-section. These are the so-called bosonic electroweak two-loop corrections, where “bosonic” refers to diagrams without closed fermion loops. Together with the corresponding results for the Z-pole asymmetries A_l, A_b , which have been presented earlier, this completes the theoretical description of Z-boson precision observables at full two-loop precision within the Standard Model. The calculation has been achieved through a combination of different methods: (a) numerical integration of Mellin–Barnes representations with contour rotations and contour shifts to improve convergence; (b) sector decomposition with numerical integration over Feynman parameters; (c) dispersion relations for sub-loop insertions. Numerical results are presented in the form of simple parameterization formulae for the total width, Γ_Z , partial decay widths $\Gamma_{e,\mu}, \Gamma_\tau, \Gamma_\nu, \Gamma_u, \Gamma_c, \Gamma_{d,s}, \Gamma_b$, branching ratios R_l, R_c, R_b and the hadronic peak cross-section, σ_{had}^0 . Theoretical intrinsic uncertainties from missing higher orders are also discussed.

© 2018 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The number of Z bosons collected at LEP in the 1990s, 1.7×10^7 , together with SLD data made it possible to determine electroweak pseudo-observables (EWPOs) with high precision: the Z-boson mass M_Z , its decay width Γ_Z , branching ratios R , forward–backward and left–right asymmetries (or equivalently A_f or $\sin^2 \theta_{\text{eff}}^f$) [1]. At that time, theoretical calculations, which included complete one-loop Standard Model corrections, selected higher order QCD corrections, and partial electroweak two-loop results with intricate QED resummations, were accurate enough to go hand-in-hand with experimental demands [2,3]. However, up to 5×10^{12} Z-boson decays are planned to be observed at projected future e^+e^- machines (ILC, FCC-ee, CEPC) running at the Z-boson resonance [4–7]. These statistics are several orders of magnitude larger than at LEP and would lead to very accurate experimental measurements of EWPOs. Limitations will come from experimental systematics, but they are in many cases estimated to

be improved by more than an order of magnitude compared to the LEP experiments [4–7]. This raises a new situation and theoretical calculations must be much more precise than assumed before [8, 9]. The improved precision will provide a platform for deep tests of the quantum structure of nature and unprecedented sensitivity to heavy or super-weakly coupled new physics.

As an important step towards that goal, this article reports on the completion of the two-loop corrections to Z-pole observables in the Glashow–Weinberg–Salam gauge theory, known as the Standard Model (SM) [10–12]. This work extends and complements previous results for the full electroweak two-loop contributions to the leptonic [13–15] and b-quark [16] effective weak mixing angles.

The first non-trivial study of electroweak (EW) loop effects was the calculation of the large quadratic top quark mass contribution to the Z and W propagators at one-loop order [17]. A few years later, the on-shell renormalization scheme as it is used today [18] and the notion of effective weak mixing angles [19] were introduced, and the scheme was used for calculations of the W^\pm and Z boson masses [20]. The complete one-loop corrections to the Z decay parameters were derived in Refs. [21–24], and those to the W^\pm width in Refs. [23,25,26]. Through the years of LEP and SLC

* Corresponding author.

E-mail address: afreitas@pitt.edu (A. Freitas).

studies, the effects of EW quantum corrections became visible in global fits of the SM parameters [1–3,27]. Global fits to EW precision measurements allowed to predict the mass of the top quark and the Higgs boson prior to their discoveries at Tevatron in 1995 [28,29] and at the LHC in 2012 [30].

At future e^+e^- colliders, EWPOs will again play a crucial role. These include the total and partial widths of the Z boson and the Z -boson couplings. The latter can be extracted from measurements of the cross-section and polarization and angular asymmetries of the processes $e^+e^- \rightarrow (Z) \rightarrow f\bar{f}$. Here f stands for any SM lepton or quark, except the top quark, whereas the notation (Z) is supposed to indicate that the amplitude is dominated by the s -channel Z -boson resonance, but there is contamination from photon and two-boson backgrounds.

Already for the precision achieved at LEP and SLC, the calculation of loop corrections beyond the one-loop order was necessary to keep theory uncertainties under control. Specifically, these included two-loop $\mathcal{O}(\alpha\alpha_s)$ [31–35] and fermionic $\mathcal{O}(\alpha^2)$ [36–50] corrections to the Fermi constant, which can be used to predict the W -boson mass, and to the Z -pole parameters. Here α refers to an electroweak loop order, whereas “fermionic” denotes contributions from diagrams with at least one closed fermion loop. In addition, leading three- and four-loop results, enhanced by powers of the top Yukawa coupling y_t , were obtained at order $\mathcal{O}(\alpha_t\alpha_s^2)$ [51,52], $\mathcal{O}(\alpha_t^2\alpha_s)$, $\mathcal{O}(\alpha_t^3)$ [53,54], and $\mathcal{O}(\alpha_t\alpha_s^3)$ [55–57], where $\alpha_t = y_t^2/(4\pi)$.

For the EW two-loop corrections, the calculation of the fermionic contributions was a natural first step, since these are numerically enhanced by the numbers of flavors and colors and by powers of y_t . Moreover, the fermionic two-loop diagrams are relatively simpler than the full set. For example, the latter includes non-planar vertex topologies, which are absent in the former. The remaining bosonic two-loop corrections to the Fermi constant and the leptonic effective weak mixing angle, $\sin^2\theta_{\text{eff}}^\ell$, have subsequently been presented in Refs. [13–15,58–61], and more recently also for the weak mixing angle in the $b\bar{b}$ channel [16].

While the numerical effects of the bosonic two-loop corrections are relatively small compared to the current experimental precision from LEP and SLC, their inclusion will become mandatory for future e^+e^- colliders. Thus the computation of the full two-loop corrections for all Z -pole EWPOs is an important goal. This article completes this goal by presenting the remaining bosonic $\mathcal{O}(\alpha^2)$ contributions to the Z -boson total and partial widths, and the hadronic Z -peak cross-section within the SM. This has been achieved by using the numerical integration methods discussed in Ref. [16], with some technical improvements.

The paper is organized as follows. After a brief review of the field theoretic definition of the relevant observables in section 2, the technical aspects of the two-loop calculation are described in section 3. The numerical impact of the bosonic EW two-loop corrections is demonstrated in section 4. In particular, results for the total and partial Z widths, several commonly used branching ratios, and the hadronic Z -peak cross-section are given in terms of simple parameterization formulae, which provide an accurate description of the full results within the currently allowed ranges of the input parameters. Finally, the theory uncertainty from missing three- and four-loop contributions is estimated in section 5, before concluding in section 6.

2. Definition of the observables

The amplitude for $e^+e^- \rightarrow f\bar{f}$ near the Z pole, $\sqrt{s} \approx M_Z$, can be written in a theoretically well-defined way as a Laurent expansion around the complex pole $s_0 \equiv \bar{M}_Z^2 - i\bar{M}_Z\bar{\Gamma}_Z$ (see Refs. [62–65]),

$$\mathcal{A}[e^+e^- \rightarrow f\bar{f}] = \frac{R}{s-s_0} + S + (s-s_0)S' + \dots, \quad (1)$$

where \bar{M}_Z and $\bar{\Gamma}_Z$ are the on-shell mass and width of the Z boson, respectively. According to eq. (1), the approximate line shape of the cross-section near the Z pole is given by $\sigma \propto [(s-\bar{M}_Z^2)^2 + \bar{M}_Z^2\bar{\Gamma}_Z^2]^{-1}$. It is important to note that this differs from the line shape ansatz used in experimental analyses, which for historical reasons is of the form $\sigma \propto [(s-M_Z^2)^2 + s^2\Gamma_Z^2/M_Z^2]^{-1}$. As a result, the parameters in eq. (1) differ from the experimental mass M_Z and width Γ_Z from LEP by a fixed factor [66]:

$$\begin{aligned} \bar{M}_Z &= M_Z / \sqrt{1 + \Gamma_Z^2/M_Z^2}, \\ \bar{\Gamma}_Z &= \Gamma_Z / \sqrt{1 + \Gamma_Z^2/M_Z^2}. \end{aligned} \quad (2)$$

Numerically, this leads to $\bar{M}_Z \approx M_Z - 34$ MeV and $\bar{\Gamma}_Z \approx \Gamma_Z - 0.9$ MeV.

The total width, $\bar{\Gamma}_Z$, can be extracted from the condition that the Z propagator has a pole at $s = s_0$, leading to

$$\bar{\Gamma}_Z = \frac{1}{M_Z} \text{Im} \Sigma_Z(s_0), \quad (3)$$

where $\Sigma_Z(s)$ is the transverse part of the Z self-energy. Using the optical theorem, it can also be written as [49,50]

$$\bar{\Gamma}_Z = \sum_f \bar{\Gamma}_f, \quad (4)$$

$$\bar{\Gamma}_f = \frac{N_c^f \bar{M}_Z}{12\pi} \left[\mathcal{R}_V^f F_V^f + \mathcal{R}_A^f F_A^f \right]_{s=\bar{M}_Z^2}. \quad (5)$$

Here the sum runs over all fermion types besides the top quark, $f = e, \mu, \tau, \nu_e, \nu_\mu, \nu_\tau, u, d, c, s, b$, and $N_c^f = 3(1)$ for quarks (leptons). The radiator functions $\mathcal{R}_{V,A}$ capture the effect of final-state QED and QCD corrections. They are known up to $\mathcal{O}(\alpha_s^4)$ and $\mathcal{O}(\alpha^2)$ for massless external fermions and $\mathcal{O}(\alpha_s^3)$ for the kinematic mass corrections [67–69]. For the results shown in this article, the explicit form given in the appendix of Ref. [50] has been used.

The remaining radiative corrections are IR finite and contained in the form factors $F_{V,A}^f$. These include massive EW corrections as well as mixed EW–QCD and EW–QED corrections. The bosonic two-loop contributions, which are of interest for this article, contribute according to [50]:

$$\begin{aligned} F_{V(2)}^f &= 2 \text{Re}(v_{f(0)} v_{f(2)}) + |v_{f(1)}|^2 \\ &\quad - v_{f(0)}^2 [\text{Re} \Sigma'_{Z(2)} - (\text{Re} \Sigma'_{Z(1)})^2] \\ &\quad - 2 \text{Re}(v_{f(0)} v_{f(1)}) \text{Re} \Sigma'_{Z(1)}, \end{aligned} \quad (6)$$

$$\begin{aligned} F_{A(2)}^f &= 2 \text{Re}(a_{f(0)} a_{f(2)}) + |a_{f(1)}|^2 \\ &\quad - a_{f(0)}^2 [\text{Re} \Sigma'_{Z(2)} - (\text{Re} \Sigma'_{Z(1)})^2] \\ &\quad - 2 \text{Re}(a_{f(0)} a_{f(1)}) \text{Re} \Sigma'_{Z(1)}, \end{aligned} \quad (7)$$

where v_f and a_f are the effective vector and axial-vector couplings, respectively, which include $Zf\bar{f}$ vertex corrections and Z - γ mixing contributions. Σ'_Z denotes the derivative of Σ_Z , and the loop order is indicated by the subscript (n) .

It should be pointed out that v_f , a_f and Σ_Z as defined above include γ - Z mixing contributions, *i.e.*

$$v_f(s) = v_f^Z(s) - v_f^\gamma(s) \frac{\Sigma_{\gamma Z}(s)}{s + \Sigma_{\gamma\gamma}(s)}, \quad (8)$$

$$a_f(s) = a_f^Z(s) - a_f^\gamma(s) \frac{\Sigma_{\gamma Z}(s)}{s + \Sigma_{\gamma\gamma}(s)}, \quad (9)$$

$$\Sigma_Z(s) = \Sigma_{ZZ}(s) - \frac{[\Sigma_{\gamma Z}(s)]^2}{s + \Sigma_{\gamma\gamma}(s)}. \quad (10)$$

Here v_f^Z and a_f^Z are the one-particle irreducible $Zf\bar{f}$ vector- and axial-vector vertex contributions, respectively, whereas v_f^γ and a_f^γ are their counterpart for the $\gamma f\bar{f}$ vertex. Furthermore, $\Sigma_{V_1 V_2}$ denotes the one-particle irreducible V_1 - V_2 self-energy.

Another important quantity is the hadronic peak cross section, σ_{had}^0 , which is defined as the total cross section for $e^+e^- \rightarrow (Z) \rightarrow$ hadrons for $s = M_Z^2$, after removal of s -channel photon exchange and box diagram contributions, as well as after the de-convolution of initial-state and initial-final interference QED effects [1,2]. The impact of the bosonic two-loop vertex corrections on σ_{had}^0 is given by [49,50]

$$\sigma_{\text{had}(2)}^0 = \sum_{f=u,d,c,s,b} \frac{12\pi}{M_Z^2} \left[\frac{\bar{\Gamma}_{e(0)}\bar{\Gamma}_{f(2)} + \bar{\Gamma}_{e(2)}\bar{\Gamma}_{f(0)}}{\bar{\Gamma}_Z^2(0)} - 2 \frac{\bar{\Gamma}_{e(0)}\bar{\Gamma}_{f(0)}\bar{\Gamma}_Z^2(2)}{\bar{\Gamma}_Z^2(0)} \right]. \quad (11)$$

The form factors $F_{V,A}^f$ are understood to include appropriate counterterms such that they are UV finite. Throughout this work, the on-shell renormalization scheme is being used, which defines all particle masses in terms of their (complex) propagator poles and the electromagnetic coupling in terms of the photon-electron vertex in the Thomson limit. A more detailed discussion of the relevant counterterms can be found in Ref. [44].

As a consequence of this renormalization scheme, the EW corrections are organized as a series in the electromagnetic coupling α , rather than the Fermi constant G_μ . Instead, G_μ will be used to compute M_W within the SM, including appropriate two-loop and partial higher-loop corrections. After this step, the remaining input parameters for the prediction of the Z coupling form factors are M_Z , M_H , m_t , G_μ , α , α_s and $\Delta\alpha$. Here $\Delta\alpha$ captures the running of the electromagnetic coupling induced by light fermion loops. It is defined through $\alpha(M_Z^2) = \alpha(0)/(1 - \Delta\alpha)$, where $\alpha(q^2)$ is the coupling at scale q^2 . The contribution from leptons to $\Delta\alpha$ can be computed perturbatively and is known at the three-loop level [70], $\Delta\alpha_{\text{lept}}(M_Z) = 0.0314976$. On the other hand, the quark contribution is non-perturbative at low scales and thus is commonly derived from experimental data. For recent evaluations of $\Delta\alpha_{\text{had}}^{(5)}$, see Refs. [71–73]. As a reference value, $\Delta\alpha_{\text{had}}^{(5)} = 0.02750$ is used in this work.

Additionally, Γ_Z and Γ_W are needed as inputs to convert M_Z and M_W to the complex pole scheme, see eq. (2). Furthermore, the radiator functions $\mathcal{R}_{V,A}^f$ depend on $m_b^{\overline{\text{MS}}}$, $m_c^{\overline{\text{MS}}}$ and m_τ to account for kinematic fermion mass effects in the final state, whereas the masses of electron, muon, neutrinos, and $u/d/s$ quarks can be taken as zero to very good approximation. In contrast to all other masses in this work, the $\overline{\text{MS}}$ masses are used for the bottom and charm quarks, since their on-shell counterparts are poorly defined.

3. Calculation of two-loop vertex corrections

For the calculations we followed the strategy developed in Ref. [16], where the two-loop bosonic corrections to the bottom

quark weak mixing angle, $\sin^2\theta_{\text{eff}}^b$, were obtained. In fact, the $Zb\bar{b}$ vertex is the technically most difficult case due to the larger number of mass scales in that problem compared to other flavors. Details are described there and also in [74–76]. On the other hand, for the computation of the Z width we are faced not only with ratios $v_{f(2)}/a_{f(2)}$, but also with sums of powers of $v_{f(2)}$ and $a_{f(2)}$, see (6) and (7). This leads to the occurrence of extra integrals which cancel out in the ratios v/a .

The complete set of two-loop diagrams required for this calculation has been generated with the computer algebra package `FeynArts 3.3` [77]. They can be divided into several categories. The renormalization counterterms require two-loop self-energies with Minkowskian external momenta, $p^2 = M_i^2 + i\varepsilon$, $M_i = M_W, M_Z$. In addition, there are two-loop vertex integrals with one non-vanishing external momentum squared, $s = M_Z^2 + i\varepsilon$. The two-loop self-energy integrals needed for the renormalization procedure and the vertex integrals with self-energy sub-loops have been computed using the dispersion relation technique described in Refs. [14,78,79]. The remaining bosonic two-loop diagrams amount to about one thousand integrals with a planar or non-planar vertex topology.

We did not try to reduce these integrals to a minimal set of master integrals, except for trivial cancellations of numerator and denominator terms. This means that tensors of rank $R \leq 3$ were calculated directly. For this purpose, two numerical approaches were used. Firstly, sector decomposition (SD) [80] was applied, with the packages `SecDec` [81,82] and `FIESTA 3` [83]. Secondly, Mellin Barnes (MB) representations [84–86] were derived and evaluated at the `MBSuite`, consisting of software packages available at the `MBtools` webpage in the `hepforge` archive [87]: `MB` [88], `MBresolve` [89], `AMBRE 1` [90], `barnesroutines` (D. Kosower) and `PlanarityTest` [91], `AMBRE 2` [92] and `AMBRE 3` [93], as well as `MBsums` [94], which are available from the `AMBRE` webpage [95]. The numerical package `MBnumerics` is being developed since 2015 [96]. It is of special importance for Minkowskian kinematics as encountered here. For the numerical integrations, `MBSuite` calls the `CUHRE` routine of the `CUBA` library [97,98].

Some new classes of integrals compared to the $\sin^2\theta_{\text{eff}}^b$ case are met. They are simpler from a numerical point of view than those solved in Ref. [16]. For instance, there are various one- and two-scale integrals with internal W propagators, which improves the singular threshold behavior of integrals with only Z propagators. There are altogether about one hundred integrals of this kind with different permutations of propagators, including the tensor integrals. As an example of one of the most difficult cases, the SD method for integrals from Fig. 1 in [16] gives an accuracy of up to four relevant digits. Using the MB method, these diagrams are equivalent to up to 4-dimensional MB integrals, which can be calculated efficiently with eight relevant digits by `MBnumerics`.

In selected cases, like those described above, the MB approach is uniquely powerful. This statement applies to several hundred integrals. In the majority of integrals, though, the SD method is presently more efficient than the MB approach, mainly due to the smaller number of integration variables. For our semi-automatized calculation of massive 2-loop vertices the availability of two complementary numerical methods with a large overlap was crucial.

4. Numerical results

In this section, numerical results for bosonic two-loop corrections are compared to and combined with all other known corrections to the $Zf\bar{f}$ vertices. These are

Table 1

Input parameters used in the numerical analysis, from [105], except for $\Delta\alpha$, for which a value close to several recent evaluations [71–73] has been chosen.

Parameter	Value
M_Z	91.1876 GeV
Γ_Z	2.4952 GeV
M_W	80.385 GeV
Γ_W	2.085 GeV
M_H	125.1 GeV
m_t	173.2 GeV
$m_b^{\overline{\text{MS}}}$	4.20 GeV
$m_c^{\overline{\text{MS}}}$	1.275 GeV
m_τ	1.777 GeV
$m_e, m_\mu, m_u, m_d, m_s$	0
$\Delta\alpha$	0.05900
$\alpha_s(M_Z)$	0.1184
G_μ	$1.16638 \times 10^{-5} \text{ GeV}^{-2}$

- Complete one-loop EW contributions [21] (which have been re-evaluated for this work) and fermionic $\mathcal{O}(\alpha^2)$ contributions [49,50];
- Mixed QCD-EW corrections to internal gauge-boson self-energies of order $\mathcal{O}(\alpha\alpha_s)$ [31–35] (where again we use our own re-evaluation of these terms);
- Higher-loop corrections in the large- m_t limit, of order $\mathcal{O}(\alpha_t\alpha_s^2)$ [51,52], $\mathcal{O}(\alpha_t^2\alpha_s)$, $\mathcal{O}(\alpha_t^3)$ [53,54], and $\mathcal{O}(\alpha_t\alpha_s^3)$ [55–57], where $\alpha_t \equiv y_t^2/(4\pi)$ and y_t is the top Yukawa coupling;
- Final-state QED radiation and, for quark final states, QCD radiation up to $\mathcal{O}(\alpha^2)$, $\mathcal{O}(\alpha\alpha_s)$ and $\mathcal{O}(\alpha_s^4)$ [67–69]; incorporated through the radiator functions $\mathcal{R}_{V,A}$ in (6) and (7);
- Non-factorizable $\mathcal{O}(\alpha\alpha_s)$ vertex contributions [99–104], which cannot be written as a product of EW form factors $F_{V,A}$ and final-state radiator functions $\mathcal{R}_{V,A}$, but instead are added separately to the formula in (5).

These are applied to a range of EWPOs: The partial Z widths, $\Gamma_f \equiv \Gamma(Z \rightarrow f\bar{f})$, as well as total width, Γ_Z , various branching ratios, and the hadronic peak cross-section σ_{had} . The full electroweak two-loop corrections for the leptonic and bottom-quark asymmetries have been published previously [13–16] and are not repeated here. Nevertheless, as a cross-check we reproduced the result for the leptonic asymmetry and found agreement with Refs. [13,15] within intrinsic numerical uncertainties. Moreover, with the methods described here we can produce results for the bosonic two-loop corrections to $\sin^2\theta_{\text{eff}}^\ell$ with four robust digits of precision, which exceeds the accuracy obtained with asymptotic expansions as in Ref. [13].

As discussed above, the gauge-boson mass renormalization has been performed in accordance with the complex-pole scheme in eq. (1). However, for the sake of comparison with the wider literature, the numerical results below are presented after translating to the scheme with an s -dependent width. In other words, results are shown for un-barred quantities, such as Γ_Z in eq. (2).

Light fermion masses m_f , $f \neq t$, have been neglected throughout, except for a non-zero bottom quark mass in the $\mathcal{O}(\alpha)$ and $\mathcal{O}(\alpha\alpha_s)$ vertex contributions, as well as for non-zero m_b , m_c and m_τ in the radiators $\mathcal{R}_{V,A}$. The numerical input values used in this section are listed in Table 1.

4.1. Partial widths

Let us begin by presenting results for a fixed value of M_W as input, instead of calculating M_W from G_μ . This more clearly il-

lustrates the impact of the newly completed $\mathcal{O}(\alpha_{\text{bos}}^2)$ corrections. Table 2 shows the contributions from different loop orders to the SM prediction of various partial Z widths. As is evident from the table, the two-loop EW corrections are significant and larger than the current experimental uncertainty (2.3 GeV for Γ_Z [1]). The newly calculated bosonic corrections $\mathcal{O}(\alpha_{\text{bos}}^2)$ are smaller but still noteworthy. They amount to half of all known leading three-loop QCD corrections $\mathcal{O}(\alpha_t\alpha_s^2, \alpha_t\alpha_s^3, \alpha_t^2\alpha_s, \alpha_t^3)$, even though the latter are enhanced by powers of α_s , α_t and N_f .

Table 3 shows the SM predictions obtained if one uses G_μ as an input to compute M_W , based on the results of [43,44,58–61]. Each line of the table adds an additional order of perturbation theory to the previous line, using the same order for the $Zf\bar{f}$ vertex corrections and the calculation of the W mass.¹

The $\mathcal{O}(\alpha_{\text{bos}}^2)$ correction to Γ_Z , corresponding to the difference between the last two rows in Table 3, amounts to 0.34 MeV, which is more than three times larger than its previous estimation [50]. An updated discussion on how this knowledge changes the intrinsic error estimations will be given in section 5.

4.2. Ratios

The experimental results from LEP and SLC are typically not presented in terms of partial widths for the different final states. Instead, this information is captured in the form of various branching ratios. The most relevant ones are

$$R_\ell \equiv \Gamma_{\text{had}}/\Gamma_\ell, \quad R_c \equiv \Gamma_c/\Gamma_{\text{had}}, \quad R_b \equiv \Gamma_b/\Gamma_{\text{had}}, \quad (12)$$

where $\Gamma_\ell = \frac{1}{3}(\Gamma_e + \Gamma_\mu + \Gamma_\tau)$, and Γ_{had} is the partial width into hadronic final states, which at the parton level is equivalent to $\sum_q \Gamma_q$ ($q = u, d, c, s, b$). In addition, the hadronic peak cross-section (11) is, to a good approximation, defined as a ratio of partial widths and the total Z width.

Numerical results for σ_{had}^0 and the ratios in (12) are given in Table 3 and Table 4, respectively, again broken down to different orders of radiative corrections. These quantities are less sensitive to higher loop effects than Γ_Z , since there is a partial cancellation between the corrections in the numerators and denominators of the ratios. Thus the influence of the new bosonic corrections on all branching ratios R_ℓ, R_c, R_b and on σ_{had}^0 is about 0.02% or less, which is far below the current experimental errors: $R_\ell = 20.767 \pm 0.025$, $R_c = 0.1721 \pm 0.0030$, $R_b = 0.21629 \pm 0.00066$, and $\sigma_{\text{had}}^0 = 41.541 \pm 0.037 \text{ nb}$ [1]. However, these are at the level of sensitivity of proposed measurements of R_b at future e^+e^- colliders [4–7].

4.3. Parameterization formulae

While the tables above only contain numbers for a single benchmark point, the results for a range of input values can be conveniently expressed in terms of simple parameterization formulae. The coefficients of these formulae have been fitted to the full calculation results on a grid that spans the currently allowed experimental ranges for each input parameter. Here the full calculation includes all higher-order corrections listed at the beginning of section 4 for the partial widths, branching ratios and the peak

¹ Note that the value in the next-to-last line of Table 3 differs slightly from Ref. [49]. This is because in Ref. [49] the “best value” prediction of M_W was carried with the full (fermionic plus bosonic) EW two-loop predictions included. Here, however, we are interested in a clear distinction of fermionic and bosonic two-loop terms in all contributions, including the M_W prediction.

Table 2

Contributions of different orders in perturbation theory to the partial and total Z widths. A fixed value of M_W has been used as input, instead of G_μ . N_f and N_f^2 refer to corrections with one and two closed fermion loops, respectively, whereas α_{bos}^2 denotes contributions without closed fermion loops. Furthermore, $\alpha_t = y_t^2/(4\pi)$. In all rows the radiator functions $\mathcal{R}_{V,A}$ with known contributions through $\mathcal{O}(\alpha_s^4)$, $\mathcal{O}(\alpha^2)$ and $\mathcal{O}(\alpha\alpha_s)$ are included.

Γ_i [MeV]	Γ_e	Γ_ν	Γ_d	Γ_u	Γ_b	Γ_Z
Born	81.142	160.096	371.141	292.445	369.562	2420.19
$\mathcal{O}(\alpha)$	2.273	6.174	9.717	5.799	3.857	60.22
$\mathcal{O}(\alpha\alpha_s)$	0.288	0.458	1.276	1.156	2.006	9.11
$\mathcal{O}(\alpha_t\alpha_s^2, \alpha_t\alpha_s^3, \alpha_t^2\alpha_s, \alpha_t^3)$	0.038	0.059	0.191	0.170	0.190	1.20
$\mathcal{O}(N_f^2\alpha^2)$	0.244	0.416	0.698	0.528	0.694	5.13
$\mathcal{O}(N_f\alpha^2)$	0.120	0.185	0.493	0.494	0.144	3.04
$\mathcal{O}(\alpha_{\text{bos}}^2)$	0.017	0.019	0.059	0.058	0.167	0.51

Table 3

Results for Γ_Z and σ_{had}^0 , with M_W calculated from G_μ using the same order of perturbation theory as indicated in each line. In all cases, the complete radiator functions $\mathcal{R}_{V,A}$ are included.

	Γ_Z [GeV]	σ_{had}^0 [nb]
Born	2.53601	41.6171
+ $\mathcal{O}(\alpha)$	2.49770	41.4687
+ $\mathcal{O}(\alpha\alpha_s)$	2.49649	41.4758
+ $\mathcal{O}(\alpha_t\alpha_s^2, \alpha_t\alpha_s^3, \alpha_t^2\alpha_s, \alpha_t^3)$	2.49560	41.4770
+ $\mathcal{O}(N_f^2\alpha^2, N_f\alpha^2)$	2.49441	41.4883
+ $\mathcal{O}(\alpha_{\text{bos}}^2)$	2.49475	41.4896

Table 4

Results for the ratios R_ℓ , R_c and R_b , with M_W calculated from G_μ to the same order as indicated in each line. In all cases, the complete radiator functions $\mathcal{R}_{V,A}$ are included.

	R_ℓ	R_c	R_b
Born	21.0272	0.17306	0.21733
+ $\mathcal{O}(\alpha)$	20.8031	0.17230	0.21558
+ $\mathcal{O}(\alpha\alpha_s)$	20.7963	0.17222	0.21593
+ $\mathcal{O}(\alpha_t\alpha_s^2, \alpha_t\alpha_s^3, \alpha_t^2\alpha_s, \alpha_t^3)$	20.7943	0.17222	0.21593
+ $\mathcal{O}(N_f^2\alpha^2, N_f\alpha^2)$	20.7512	0.17223	0.21580
+ $\mathcal{O}(\alpha_{\text{bos}}^2)$	20.7516	0.17222	0.21585

cross-sections, and with M_W calculated from G_μ to the same precision.² For all EWPOs reported here, the same form of parameterization formula is utilized:

$$X = X_0 + c_1 L_H + c_2 \Delta_t + c_3 \Delta_{\alpha_s} + c_4 \Delta_{\alpha_s}^2 + c_5 \Delta_{\alpha_s} \Delta_t + c_6 \Delta_\alpha + c_7 \Delta_Z, \quad (13)$$

$$L_H = \log \frac{M_H}{125.7 \text{ GeV}}, \quad \Delta_t = \left(\frac{m_t}{173.2 \text{ GeV}} \right)^2 - 1,$$

$$\Delta_{\alpha_s} = \frac{\alpha_s(M_Z)}{0.1184} - 1, \quad \Delta_\alpha = \frac{\Delta\alpha}{0.059} - 1,$$

$$\Delta_Z = \frac{M_Z}{91.1876 \text{ GeV}} - 1.$$

As before, M_H , M_Z , m_t and $\Delta\alpha$ are defined in the on-shell scheme, using the s -dependent width scheme for M_Z (to match the published experimental values), while α_s is defined in the $\overline{\text{MS}}$ scheme. The dependence on m_b , m_c and m_τ is negligible within the allowed ranges for these quantities.

² Fit formulae for the leptonic and bottom-quark asymmetries can be found in Refs. [13,14,16].

The fit values of the coefficients for the different EWPOs are given in Table 5. With these parameters, the formulae provide very good approximations to the full results within the ranges $M_H = 125.1 \pm 5.0$ GeV, $m_t = 173.2 \pm 4.0$ GeV, $\alpha_s = 0.1184 \pm 0.0050$, $\Delta\alpha = 0.0590 \pm 0.0005$ and $M_Z = 91.1876 \pm 0.0042$ GeV, with maximal deviations as quoted in the last column of Table 5. As can be seen from the latter, the accuracies of the fit formulae are sufficient for the foreseeable future.

5. Error estimates

In addition to the dependence on the input parameters, the accuracy of the results presented here is limited by unknown three- and four-loop contributions. The numerically leading missing pieces are the $\mathcal{O}(\alpha^3)$, $\mathcal{O}(\alpha^2\alpha_s)$, $\mathcal{O}(\alpha\alpha_s^2)$ and $\mathcal{O}(\alpha\alpha_s^3)$ corrections beyond the known leading y_t^n terms from Refs. [51–57].

Following Refs. [50,106], the size of these terms may be estimated by assuming that the perturbation series approximately is a geometric series. In this way one obtains

$$\begin{aligned} \mathcal{O}(\alpha^3) - \mathcal{O}(\alpha_t^3) &\sim \frac{\mathcal{O}(\alpha^2) - \mathcal{O}(\alpha_t^2)}{\mathcal{O}(\alpha)} \mathcal{O}(\alpha^2), \\ \mathcal{O}(\alpha^2\alpha_s) - \mathcal{O}(\alpha_t^2\alpha_s) &\sim \frac{\mathcal{O}(\alpha^2) - \mathcal{O}(\alpha_t^2)}{\mathcal{O}(\alpha)} \mathcal{O}(\alpha\alpha_s), \\ \mathcal{O}(\alpha\alpha_s^2) - \mathcal{O}(\alpha_t\alpha_s^2) &\sim \frac{\mathcal{O}(\alpha\alpha_s) - \mathcal{O}(\alpha_t\alpha_s)}{\mathcal{O}(\alpha)} \mathcal{O}(\alpha\alpha_s), \\ \mathcal{O}(\alpha\alpha_s^3) - \mathcal{O}(\alpha_t\alpha_s^3) &\sim \frac{\mathcal{O}(\alpha\alpha_s) - \mathcal{O}(\alpha_t\alpha_s)}{\mathcal{O}(\alpha)} \mathcal{O}(\alpha\alpha_s^2), \end{aligned} \quad (14)$$

where the known leading large- m_t approximations have been subtracted in the numerators. For the example of the total Z width, these expressions lead to

$$\begin{aligned} \Gamma_Z: \quad \mathcal{O}(\alpha^3) - \mathcal{O}(\alpha_t^3) &\sim 0.20 \text{ MeV}, \\ \mathcal{O}(\alpha^2\alpha_s) - \mathcal{O}(\alpha_t^2\alpha_s) &\sim 0.21 \text{ MeV}, \\ \mathcal{O}(\alpha\alpha_s^2) - \mathcal{O}(\alpha_t\alpha_s^2) &\sim 0.23 \text{ MeV}, \\ \mathcal{O}(\alpha\alpha_s^3) - \mathcal{O}(\alpha_t\alpha_s^3) &\sim 0.035 \text{ MeV}. \end{aligned} \quad (15)$$

An additional source of theoretical uncertainty stems from the unknown $\mathcal{O}(\alpha_s^2)$ final-state QCD corrections and three-loop mixed QED/QCD final-state corrections of order $\mathcal{O}(\alpha\alpha_s^2)$ and $\mathcal{O}(\alpha^2\alpha_s)$. In [50] they were found to be sub-dominant, and the estimates can be taken over from there without change. Combining these findings with eqs. (15) in quadrature, the total theory error adds up to $\delta\Gamma_Z \approx 0.4$ MeV. Compared to the previous theory error estimate $\delta\Gamma_Z \approx 0.5$ MeV [50] one observes a slight decrease due to the knowledge of the bosonic corrections calculated in this work.

Table 5

Coefficients for the parameterization formula (13) for various observables (X). Within the ranges $M_H = 125.1 \pm 5.0$ GeV, $m_t = 173.2 \pm 4.0$ GeV, $\alpha_s = 0.1184 \pm 0.0050$, $\Delta\alpha = 0.0590 \pm 0.0005$ and $M_Z = 91.1876 \pm 0.0042$ GeV, the formulae approximate the full results with maximal deviations given in the last column.

Observable	X_0	c_1	c_2	c_3	c_4	c_5	c_6	c_7	max. dev.
$\Gamma_{e,\mu}$ [MeV]	83.983	-0.061	0.810	-0.096	-0.01	0.25	-1.1	286	< 0.001
Γ_τ [MeV]	83.793	-0.060	0.810	-0.095	-0.01	0.25	-1.1	285	< 0.001
Γ_ν [MeV]	167.176	-0.071	1.26	-0.19	-0.02	0.36	-0.1	504	< 0.001
Γ_u [MeV]	299.993	-0.38	4.08	14.27	1.6	1.8	-11.1	1253	< 0.002
Γ_c [MeV]	299.916	-0.38	4.08	14.27	1.6	1.8	-11.1	1253	< 0.002
$\Gamma_{d,s}$ [MeV]	382.828	-0.39	3.83	10.20	-2.4	0.67	-10.1	1470	< 0.002
Γ_b [MeV]	375.889	-0.36	-2.14	10.53	-2.4	1.2	-10.1	1459	< 0.006
Γ_Z [MeV]	2494.74	-2.3	19.9	58.61	-4.0	8.0	-56.0	9273	< 0.012
R_ℓ [10^{-3}]	20751.6	-7.8	-37	732.3	-44	5.5	-358	11696	< 0.1
R_c [10^{-3}]	172.22	-0.031	1.0	2.3	1.3	0.38	-1.2	37	< 0.01
R_b [10^{-3}]	215.85	0.029	-2.92	-1.32	-0.84	0.032	0.72	-18	< 0.01
σ_{had}^0 [pb]	41489.6	1.6	60.0	-579.6	38	7.3	85	-86011	< 0.1

Table 6

Theory uncertainty estimates for the partial and total Z widths and branching ratios from missing 3-loop and higher orders. See text for details.

$\Gamma_{e,\mu\tau}$	0.018 MeV	$\Gamma_{u,c}$	0.11 MeV	R_ℓ	$6 \cdot 10^{-3}$
Γ_ν	0.016 MeV	Γ_b	0.18 MeV	R_c	$5 \cdot 10^{-5}$
$\Gamma_{d,s}$	0.08 MeV	Γ_Z	0.4 MeV	R_b	$1 \cdot 10^{-4}$

In addition to the elimination of an uncertainty associated with the previous unknown $\mathcal{O}(\alpha_{\text{bos}}^2)$ corrections, the values in the first and second rows of (15) also shifted since the full $\mathcal{O}(\alpha^2)$ corrections used in (14) were not available before. These shifts conspire to result in a reduction of the uncertainty estimate for these two error sources.

The corresponding error estimates for the partial widths are shown in Table 6. For the ratios (R_ℓ , R_c and R_b), the theory uncertainty has been estimated from the partial widths using simple Gaussian error propagation.

The theory uncertainty for the hadronic peak cross-section is dominated by a non-factorizable contribution stemming from the imaginary part of the Z -boson self-energy [50]. This non-factorizable term does not receive any bosonic two-loop corrections, so that its error estimate can be taken from Ref. [50] without change:

$$\sigma_{\text{had}}^0 : \mathcal{O}(\alpha^3) \sim 3.7 \text{ pb}, \quad \mathcal{O}(\alpha^2\alpha_s) \sim 4.2 \text{ pb}. \quad (16)$$

Adding these in quadrature leads to the overall uncertainty estimate of $\delta\sigma_{\text{had}}^0 \approx 6$ pb.

6. Summary

In this work the bosonic two-loop electroweak corrections, $\mathcal{O}(\alpha_{\text{bos}}^2)$, to Z boson production and decay parameters are presented for the first time. These corrections are comparable in size to the leading three-loop corrections of $\mathcal{O}(\alpha_t\alpha_s^2)$, $\mathcal{O}(\alpha_t\alpha_s^3)$, $\mathcal{O}(\alpha_t^2\alpha_s)$, $\mathcal{O}(\alpha_t^3)$. This is especially pronounced for Γ_b , see Table 2, and for σ_{had}^0 , see Table 3. The bosonic corrections shift the value of Γ_Z by 0.51 MeV when using M_W as input and 0.34 MeV when using G_μ as input, which is large from the point of view of future colliders. Similarly, the bosonic corrections are important for R_b , see Table 4. Due to the high accuracy of the numerical loop integrations, the results obtained here are stable enough even in the context of potential future experimental precisions.

Updated theory error estimations are given, which are slightly reduced due to the newly available full two-loop corrections. We expect that the numerical integration methods used here can be extended to compute the full three-loop and possibly part of the four-loop corrections to Z -pole EWPOs. These improvements will require substantial technical work and ingenuity, but the challenges are well understood and thus could in principle be overcome. The ultimate achievable precision is limited by two factors: (a) the complexity of the analytic properties of multi-loop diagram topologies and their translation into well-defined numerical integrals, and (b) the numerical precision with which these integrals can be evaluated. If these can be driven to allow the computation of mixed EW/QCD and leading EW 4-loop topologies with at least two digits precision, theory uncertainties will not be a limiting factor for future e^+e^- experiments. For a more detailed discussion of future projections, see Ref. [8,9].

It should be noted that the $\mathcal{O}(\alpha_{\text{bos}}^2)$ correction for the total Z decay width appears to be relatively large compared to previous estimates based on the knowledge of the lower order result $\mathcal{O}(\alpha_{\text{bos}})$. A similar observation concerns the bosonic two-loop corrections to A_b . This means that all estimations at this level of accuracy should be taken with a grain of salt. Therefore, explicit calculations are important even for contributions that were previously estimated to be subdominant.

At this point we should mention that we did not consider the theoretical efforts needed to unfold the large QED corrections from the measured real cross sections in the Z peak region and to extract the EWPOs studied here in detail. For LEP, this was based on tools such as the ZFITTER package [107–109] and was discussed carefully e.g. in Refs. [1,2,110]. The correct unfolding framework for extracting $2 \rightarrow 2$ observables at accuracies amounting to about 1/20 of the LEP era certainly has to rely on the correct treatment of Laurent series for the Z line shape as is discussed e.g. in [111–114].

The 1-loop corrections to the Z boson parameters were determined in the 1980s [21]. Today, 33 years later, while the present study finalizes the determination of the electroweak two-loop corrections to the Z -boson parameters, we are already faced with the need of more precision in the future.

Acknowledgements

The work of I.D. is supported by a research grant (no. 91530979/2014) of Deutscher Akademischer Austauschdienst (DAAD) and by Deutsches Elektronen-Synchrotron (DESY). The work of A.F. is sup-

ported in part by the National Science Foundation under grant PHY-1519175. The work of J.G. is supported in part by the Polish National Science Centre under grant no. 2017/25/B/ST2/01987 and COST Action CA16201 PARTICLEFACE. The work of T.R. is supported in part by DESY and an Alexander von Humboldt Polish Honorary Research Fellowship. J.U. received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under grant agreement no. 647356 (CutLoops). We would like to thank Peter Uwer and his group "Phenomenology of Elementary Particle Physics beyond the Standard Model" at Humboldt-Universität zu Berlin for providing computer resources.

References

- [1] S. Schael, et al., Precision electroweak measurements on the Z resonance, Phys. Rep. 427 (2006) 257–454, <https://doi.org/10.1016/j.physrep.2005.12.006>, arXiv:hep-ex/0509008.
- [2] D. Bardin, W. Beenakker, M. Bilenky, W. Hollik, M. Martinez, G. Montagna, O. Nicrosini, V. Novikov, L. Okun, A. Olshchovsky, G. Passarino, F. Piccinini, S. Riemann, T. Riemann, A. Rozaanov, F. Teubert, M. Vysotsky, Electroweak Working Group Report 7-162, CERN 95-03A. In [3], arXiv:hep-ph/9709229.
- [3] D. Bardin, W. Hollik, G. Passarino (Eds.), Reports of the Working Group on Precision Calculations for the Z Resonance, report CERN 95-03, <http://cds.cern.ch/record/280836/files/CERN-95-03.pdf?version=2>, 31 March 1995.
- [4] H. Baer, T. Barklow, K. Fujii, Y. Gao, A. Hoang, S. Kanemura, J. List, H.E. Logan, A. Nomerotski, M. Perelstein, et al., The International Linear Collider Technical Design Report - Volume 2: Physics, arXiv:1306.6352, 2013.
- [5] M. Bicer, et al., First look at the physics case of tLEP, J. High Energy Phys. 01 (2014) 164, [https://doi.org/10.1007/JHEP01\(2014\)164](https://doi.org/10.1007/JHEP01(2014)164), arXiv:1308.6176.
- [6] D. d'Enterria, Physics at the FCC-ee, in: Proceedings of the 17th Lomonosov Conference on Elementary Particle Physics, Moscow, Russia, 20–26 August 2015, 2017, pp. 182–191, arXiv:1602.05043, <http://inspirehep.net/record/1421932/files/arXiv:1602.05043.pdf>.
- [7] CEPC-SPPC Study Group, CEPC-SPPC Preliminary Conceptual Design Report. 1. Physics and Detector, IHEP-CEPC-DR-2015-01, IHEP-TH-2015-01, IHEP-EP-2015-01, http://inspirehep.net/record/1395734/files/main_preCDR.pdf, 2015.
- [8] A. Blondel, J. Gluza, P. Janot (org.), Mini Workshop on Precision EW and QCD Calculations for the FCC Studies: Methods and Techniques, 12–13 Jan 2018, CERN, webpage <https://indico.cern.ch/event/669224/>; A. Blondel, J. Gluza, S. Jadach, P. Janot, T. Riemann (Eds.), Report, in preparation.
- [9] I. Dubovyk, A. Freitas, J. Gluza, K. Grzanka, S. Jadach, T. Riemann, J. Usovitsch, Precision calculations for the Z line shape at the FCC-ee, Poster presented at the FCC Week 2018, 9–13 April 2018, Amsterdam, https://indico.cern.ch/event/656491/contributions/2947663/attachments/1622685/2582801/Poster-FCC-Amsterdam_Sjadach_et_al.pdf.
- [10] S. Weinberg, A model of leptons, Phys. Rev. Lett. 19 (1967) 1264–1266, <https://doi.org/10.1103/PhysRevLett.19.1264>.
- [11] S. Glashow, Partial symmetries of weak interactions, Nucl. Phys. 22 (1961) 579–588, [https://doi.org/10.1016/0029-5582\(61\)90469-2](https://doi.org/10.1016/0029-5582(61)90469-2).
- [12] A. Salam, Weak and electromagnetic interactions, originally printed in: Svartholm (Ed.), Elementary Particle Theory, Proceedings of the Nobel Symposium held 1968 at Lerum, Sweden, Stockholm 1968, 1968, pp. 367–377; see also: <http://inspirehep.net/record/53083?ln=de>.
- [13] M. Awramik, M. Czakon, A. Freitas, Bosonic corrections to the effective weak mixing angle at $O(\alpha^2)$, Phys. Lett. B 642 (2006) 563–566, <https://doi.org/10.1016/j.physletb.2006.07.035>, arXiv:hep-ph/0605339.
- [14] M. Awramik, M. Czakon, A. Freitas, Electroweak two-loop corrections to the effective weak mixing angle, J. High Energy Phys. 11 (2006) 048, <https://doi.org/10.1088/1126-6708/2006/11/048>, arXiv:hep-ph/0608099.
- [15] W. Hollik, U. Meier, S. Uccirati, The effective electroweak mixing angle $\sin^2 \theta_{\text{eff}}$ with two-loop bosonic contributions, Nucl. Phys. B 765 (2007) 154–165, <https://doi.org/10.1016/j.nuclphysb.2006.12.001>, arXiv:hep-ph/0610312.
- [16] I. Dubovyk, A. Freitas, J. Gluza, T. Riemann, J. Usovitsch, The two-loop electroweak bosonic corrections to $\sin^2 \theta_{\text{eff}}^b$, Phys. Lett. B 762 (2016) 184–189, <https://doi.org/10.1016/j.physletb.2016.09.012>, arXiv:1607.08375.
- [17] M. Veltman, Limit on mass differences in the Weinberg model, Nucl. Phys. B 123 (1977) 89, [https://doi.org/10.1016/0550-3213\(77\)90342-X](https://doi.org/10.1016/0550-3213(77)90342-X).
- [18] A. Sirlin, Radiative corrections in the $SU(2)_L \times U(1)$ theory: a simple renormalization framework, Phys. Rev. D 22 (1980) 971–981, <https://doi.org/10.1103/PhysRevD.22.971>.
- [19] W.J. Marciano, A. Sirlin, Radiative corrections to neutrino induced neutral current phenomena in the $SU(2)_L \times U(1)$ theory, Phys. Rev. D 22 (1980) 2695, <https://doi.org/10.1103/PhysRevD.22.2695>, Phys. Rev. D 31 (1985) 213, <https://doi.org/10.1103/PhysRevD.31.213> (Erratum).
- [20] W.J. Marciano, A. Sirlin, Testing the Standard Model by precise determinations of W^\pm and Z masses, Phys. Rev. D 29 (1984) 945, <https://doi.org/10.1103/PhysRevD.29.945>, Phys. Rev. D 31 (1985) 213, <https://doi.org/10.1103/PhysRevD.31.213.3> (Erratum).
- [21] A. Akhundov, D. Bardin, T. Riemann, Electroweak one loop corrections to the decay of the neutral vector boson, Nucl. Phys. B 276 (1986) 1, [https://doi.org/10.1016/0550-3213\(86\)90014-3](https://doi.org/10.1016/0550-3213(86)90014-3).
- [22] W. Beenakker, W. Hollik, The width of the Z boson, Z. Phys. C 40 (1988) 141, <https://doi.org/10.1007/BF01559728>.
- [23] F. Jegerlehner, Precision tests of electroweak interaction parameters, in: R. Manka, M. Zralek (Eds.), Proc. 11th Int. School of Theoretical Physics, Testing the Standard Model, Szczyrk, Poland, 18–22 Sep. 1987, World Scientific, Singapore, 1988, pp. 33–108, <http://ccdb5fs.kek.jp/cgi-bin/img/allpdf?198801263>.
- [24] J. Bernabeu, A. Pich, A. Santamaria, $\Gamma(Z \rightarrow b\bar{b})$: a signature of hard mass terms for a heavy top, Phys. Lett. B 200 (1988) 569, [https://doi.org/10.1016/0370-2693\(88\)90173-6](https://doi.org/10.1016/0370-2693(88)90173-6).
- [25] D. Bardin, S. Riemann, T. Riemann, Electroweak one loop corrections to the decay of the charged vector boson, Z. Phys. C 32 (1986) 121–125, <https://doi.org/10.1007/BF01441360>.
- [26] A. Denner, T. Sack, The W boson width, Z. Phys. C 46 (1990) 653–663, <https://doi.org/10.1007/BF01560267>.
- [27] K. Hikasa, et al., Review of particle properties. Particle Data Group, Phys. Rev. D 45 (1992) S1, <https://doi.org/10.1103/PhysRevD.45.S1>, Phys. Rev. D 46 (1992) 5210, <https://doi.org/10.1103/PhysRevD.46.5210> (Erratum).
- [28] F. Abe, et al., Observation of top quark production in $\bar{p}p$ collisions, Phys. Rev. Lett. 74 (1995) 2626–2631, <https://doi.org/10.1103/PhysRevLett.74.2626>, arXiv:hep-ex/9503002.
- [29] S. Abachi, et al., Observation of the top quark, Phys. Rev. Lett. 74 (1995) 2632–2637, <https://doi.org/10.1103/PhysRevLett.74.2632>, arXiv:hep-ex/9503003.
- [30] G. Aad, et al., Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC, Phys. Lett. B 716 (2012) 1–29, <https://doi.org/10.1016/j.physletb.2012.08.020>, arXiv:1207.7214.
- [31] A. Djouadi, C. Verzegnassi, Virtual very heavy top effects in LEP/SLC precision measurements, Phys. Lett. B 195 (1987) 265–271, [https://doi.org/10.1016/0370-2693\(87\)91206-8](https://doi.org/10.1016/0370-2693(87)91206-8).
- [32] A. Djouadi, $O(\alpha_s)$ vacuum polarization functions of the standard model gauge bosons, Nuovo Cimento A 100 (1988) 357, <https://doi.org/10.1007/BF02812964>.
- [33] B.A. Kniehl, Two loop corrections to the vacuum polarizations in perturbative QCD, Nucl. Phys. B 347 (1990) 86–104, [https://doi.org/10.1016/0550-3213\(90\)90552-0](https://doi.org/10.1016/0550-3213(90)90552-0).
- [34] B.A. Kniehl, A. Sirlin, Dispersion relations for vacuum polarization functions in electroweak physics, Nucl. Phys. B 371 (1992) 141–148, [https://doi.org/10.1016/0550-3213\(92\)90232-Z](https://doi.org/10.1016/0550-3213(92)90232-Z).
- [35] A. Djouadi, P. Gambino, Electroweak gauge bosons selfenergies: complete QCD corrections, Phys. Rev. D 49 (1994) 3499–3511, <https://doi.org/10.1103/PhysRevD.49.3499>, arXiv:hep-ph/9309298, Phys. Rev. D 53 (1996) 4111, <https://doi.org/10.1103/PhysRevD.53.4111> (Erratum).
- [36] R. Barbieri, M. Beccaria, P. Ciafaloni, G. Curci, A. Vicere, Radiative correction effects of a very heavy top, Phys. Lett. B 288 (1992) 95–98, [https://doi.org/10.1016/0370-2693\(92\)91960-H](https://doi.org/10.1016/0370-2693(92)91960-H), arXiv:hep-ph/9205238, Phys. Lett. B 312 (1993) 511, [https://doi.org/10.1016/0370-2693\(93\)90990-Y](https://doi.org/10.1016/0370-2693(93)90990-Y) (Erratum).
- [37] R. Barbieri, M. Beccaria, P. Ciafaloni, G. Curci, A. Vicere, Two loop heavy top effects in the Standard Model, Nucl. Phys. B 409 (1993) 105–127, [https://doi.org/10.1016/0550-3213\(93\)90448-X](https://doi.org/10.1016/0550-3213(93)90448-X).
- [38] J. Fleischer, O.V. Tarasov, F. Jegerlehner, Two loop heavy top corrections to the rho parameter: a simple formula valid for arbitrary Higgs mass, Phys. Lett. B 319 (1993) 249–256, [https://doi.org/10.1016/0370-2693\(93\)90810-5](https://doi.org/10.1016/0370-2693(93)90810-5).
- [39] J. Fleischer, O.V. Tarasov, F. Jegerlehner, Two loop large top mass corrections to electroweak parameters: analytic results valid for arbitrary Higgs mass, Phys. Rev. D 51 (1995) 3820–3837, <https://doi.org/10.1103/PhysRevD.51.3820>.
- [40] G. Degrossi, P. Gambino, A. Vicini, Two loop heavy top effects on the $M_Z - M_W$ interdependence, Phys. Lett. B 383 (1996) 219–226, [https://doi.org/10.1016/0370-2693\(96\)00720-4](https://doi.org/10.1016/0370-2693(96)00720-4), arXiv:hep-ph/9603374.
- [41] G. Degrossi, P. Gambino, A. Sirlin, Precise calculation of M_W , $\sin^2 \hat{\theta}_W(M_Z)$, and $\sin^2 \theta_{\text{eff}}^{\text{lep}}$, Phys. Lett. B 394 (1997) 188–194, [https://doi.org/10.1016/S0370-2693\(96\)01677-2](https://doi.org/10.1016/S0370-2693(96)01677-2), arXiv:hep-ph/9611363.
- [42] G. Degrossi, P. Gambino, Two loop heavy top corrections to the Z^0 boson partial widths, Nucl. Phys. B 567 (2000) 3–31, [https://doi.org/10.1016/S0550-3213\(99\)00729-4](https://doi.org/10.1016/S0550-3213(99)00729-4), arXiv:hep-ph/9905472.
- [43] A. Freitas, W. Hollik, W. Walter, G. Weiglein, Complete fermionic two loop results for the $M_W - M_Z$ interdependence, Phys. Lett. B 495 (2000) 338–346, [https://doi.org/10.1016/S0370-2693\(00\)01263-6](https://doi.org/10.1016/S0370-2693(00)01263-6), arXiv:hep-ph/0007091, Phys. Lett. B 570 (2003) 265, <https://doi.org/10.1016/j.physletb.2003.08.006> (Erratum).
- [44] A. Freitas, W. Hollik, W. Walter, G. Weiglein, Electroweak two loop corrections to the $M_W - M_Z$ mass correlation in the Standard Model, Nucl. Phys. B 632 (2002) 189–218, [https://doi.org/10.1016/S0550-3213\(02\)00243-2](https://doi.org/10.1016/S0550-3213(02)00243-2), arXiv:hep-ph/0202131, Nucl. Phys. B 666 (2003) 305 (Erratum).

- [45] M. Awramik, M. Czakon, A. Freitas, G. Weiglein, Complete two-loop electroweak fermionic corrections to $\sin^2 \theta_{\text{eff}}^{\text{lep}}$ and indirect determination of the Higgs boson mass, *Phys. Rev. Lett.* 93 (2004) 201805, <https://doi.org/10.1103/PhysRevLett.93.201805>, arXiv:hep-ph/0407317.
- [46] W. Hollik, U. Meier, S. Uccirati, The effective electroweak mixing angle $\sin^2 \theta_{\text{eff}}$ with two-loop fermionic contributions, *Nucl. Phys. B* 731 (2005) 213–224, <https://doi.org/10.1016/j.nuclphysb.2005.10.015>, arXiv:hep-ph/0507158.
- [47] M. Awramik, M. Czakon, A. Freitas, B. Kniehl, Two-loop electroweak fermionic corrections to $\sin^2 \theta_{\text{eff}}^{\text{bb}}$, *Nucl. Phys. B* 813 (2009) 174–187, <https://doi.org/10.1016/j.nuclphysb.2008.12.031>, arXiv:0811.1364.
- [48] A. Freitas, Y.-C. Huang, Electroweak two-loop corrections to $\sin^2 \theta_{\text{eff}}^{\text{bb}}$ and R_b using numerical Mellin–Barnes integrals, *J. High Energy Phys.* 1208 (2012) 050, [https://doi.org/10.1007/JHEP08\(2012\)050](https://doi.org/10.1007/JHEP08(2012)050), arXiv:1205.0299.
- [49] A. Freitas, Two-loop fermionic electroweak corrections to the Z-boson width and production rate, *Phys. Lett. B* 730 (2014) 50–52, <https://doi.org/10.1016/j.physletb.2014.01.017>, arXiv:1310.2256.
- [50] A. Freitas, Higher-order electroweak corrections to the partial widths and branching ratios of the Z boson, *J. High Energy Phys.* 1404 (2014) 070, [https://doi.org/10.1007/JHEP04\(2014\)070](https://doi.org/10.1007/JHEP04(2014)070), arXiv:1401.2447.
- [51] L. Avdeev, J. Fleischer, S. Mikhailov, O. Tarasov, $O(\alpha_s^2)$ correction to the electroweak ρ parameter, *Phys. Lett. B* 336 (1994) 560–566, [https://doi.org/10.1016/0370-2693\(94\)90573-8](https://doi.org/10.1016/0370-2693(94)90573-8), arXiv:hep-ph/9406363, *Phys. Lett. B* 349 (1995) 597 (Erratum).
- [52] K. Chetyrkin, J.H. Kühn, M. Steinhauser, Corrections of order $O(G_F M_t^2 \alpha_s^2)$ to the ρ parameter, *Phys. Lett. B* 351 (1995) 331–338, [https://doi.org/10.1016/0370-2693\(95\)00380-4](https://doi.org/10.1016/0370-2693(95)00380-4), arXiv:hep-ph/9502291.
- [53] J.J. van der Bij, K.G. Chetyrkin, M. Faisst, G. Jikia, T. Seidensticker, Three loop leading top mass contributions to the ρ parameter, *Phys. Lett. B* 498 (2001) 156–162, [https://doi.org/10.1016/S0370-2693\(01\)00002-8](https://doi.org/10.1016/S0370-2693(01)00002-8), arXiv:hep-ph/0011373.
- [54] M. Faisst, J.H. Kühn, T. Seidensticker, O. Veretin, Three loop top quark contributions to the ρ parameter, *Nucl. Phys. B* 665 (2003) 649–662, [https://doi.org/10.1016/S0550-3213\(03\)00450-4](https://doi.org/10.1016/S0550-3213(03)00450-4), arXiv:hep-ph/0302275.
- [55] Y. Schröder, M. Steinhauser, Four-loop singlet contribution to the ρ parameter, *Phys. Lett. B* 622 (2005) 124–130, <https://doi.org/10.1016/j.physletb.2005.06.085>, arXiv:hep-ph/0504055.
- [56] K.G. Chetyrkin, M. Faisst, J.H. Kühn, P. Maierhofer, C. Sturm, Four-loop QCD corrections to the ρ parameter, *Phys. Rev. Lett.* 97 (2006) 102003, <https://doi.org/10.1103/PhysRevLett.97.102003>, arXiv:hep-ph/0605201.
- [57] R. Boughezal, M. Czakon, Single scale tadpoles and $O(G_F M_t^2 \alpha_s^2)$ corrections to the ρ parameter, *Nucl. Phys. B* 755 (2006) 221–238, <https://doi.org/10.1016/j.nuclphysb.2006.08.007>, arXiv:hep-ph/0606232.
- [58] M. Awramik, M. Czakon, Complete two loop bosonic contributions to the muon lifetime in the Standard Model, *Phys. Rev. Lett.* 89 (2002) 241801, <https://doi.org/10.1103/PhysRevLett.89.241801>, arXiv:hep-ph/0208113.
- [59] M. Awramik, M. Czakon, Complete two loop electroweak contributions to the muon lifetime in the Standard Model, *Phys. Lett. B* 568 (2003) 48–54, <https://doi.org/10.1016/j.physletb.2003.06.007>, arXiv:hep-ph/0305248.
- [60] A. Onishchenko, O. Veretin, Two loop bosonic electroweak corrections to the muon lifetime and $M_Z - M_W$ interdependence, *Phys. Lett. B* 551 (2003) 111–114, [https://doi.org/10.1016/S0370-2693\(02\)03004-6](https://doi.org/10.1016/S0370-2693(02)03004-6), arXiv:hep-ph/0209010.
- [61] M. Awramik, M. Czakon, A. Freitas, G. Weiglein, Precise prediction for the W boson mass in the Standard Model, *Phys. Rev. D* 69 (2004) 053006, <https://doi.org/10.1103/PhysRevD.69.053006>, arXiv:hep-ph/0311148.
- [62] S. Willenbrock, G. Valencia, On the definition of the Z boson mass, *Phys. Lett. B* 259 (1991) 373–376, [https://doi.org/10.1016/0370-2693\(91\)90843-F](https://doi.org/10.1016/0370-2693(91)90843-F).
- [63] A. Sirlin, Theoretical considerations concerning the Z^0 mass, *Phys. Rev. Lett.* 67 (1991) 2127–2130, <https://doi.org/10.1103/PhysRevLett.67.2127>.
- [64] R. Stuart, Gauge invariance, analyticity and physical observables at the Z^0 resonance, *Phys. Lett. B* 262 (1991) 113–119, [https://doi.org/10.1016/0370-2693\(91\)90653-8](https://doi.org/10.1016/0370-2693(91)90653-8).
- [65] H.G.J. Veltman, Mass and width of unstable gauge bosons, *Z. Phys. C* 62 (1994) 35–52, <https://doi.org/10.1007/BF01559523>.
- [66] D. Bardin, A. Leike, T. Riemann, M. Sachwitz, Energy dependent width effects in e^+e^- annihilation near the Z boson pole, *Phys. Lett. B* 206 (1988) 539–542, [https://doi.org/10.1016/0370-2693\(88\)91625-5](https://doi.org/10.1016/0370-2693(88)91625-5).
- [67] K. Chetyrkin, J.H. Kühn, A. Kwiatkowski, QCD corrections to the e^+e^- cross-section and the Z boson decay rate, in: *Reports of the Working Group on Precision Calculations for the Z Resonance*, Geneva, 1994, CERN 95-03, 31 March 1996, pp. 175–263. See also: QCD corrections to the e^+e^- cross-section and the Z boson decay rate: concepts and results, March 1996, 105 pages, LBL-36678-REV and *Phys. Rep.* 277 (1996) 189–281, arXiv:hep-ph/9503396.
- [68] P. Baikov, K. Chetyrkin, J. Kühn, Order α_s^4 QCD corrections to Z and τ decays, *Phys. Rev. Lett.* 101 (2008) 012002, <https://doi.org/10.1103/PhysRevLett.101.012002>, arXiv:0801.1821.
- [69] A.L. Kataev, Higher order $O(\alpha^2)$ and $O(\alpha_s)$ corrections to $\sigma_{\text{tot}}(e^+e^- \rightarrow \text{hadrons})$ and Z boson decay rate, *Phys. Lett. B* 287 (1992) 209–212, [https://doi.org/10.1016/0370-2693\(92\)91901-K](https://doi.org/10.1016/0370-2693(92)91901-K).
- [70] M. Steinhauser, Leptonic contribution to the effective electromagnetic coupling constant up to three loops, *Phys. Lett. B* 429 (1998) 158–161, [https://doi.org/10.1016/S0370-2693\(98\)00503-6](https://doi.org/10.1016/S0370-2693(98)00503-6), arXiv:hep-ph/9803313.
- [71] M. Davier, A. Hoecker, B. Malaescu, Z. Zhang, Reevaluation of the hadronic vacuum polarisation contributions to the Standard Model predictions of the muon $g-2$ and $\alpha(m_Z^2)$ using newest hadronic cross-section data, *Eur. Phys. J. C* 77 (12) (2017) 827, <https://doi.org/10.1140/epjc/s10052-017-5161-6>, arXiv:1706.09436.
- [72] F. Jegerlehner, Variations on photon vacuum polarization, arXiv:1711.06089.
- [73] A. Keshavarzi, D. Nomura, T. Teubner, The muon $g-2$ and $\alpha(M_Z^2)$: a new data-based analysis, arXiv:1802.02995.
- [74] I. Dubovyk, J. Gluza, T. Riemann, J. Usovitsch, Numerical integration of massive two-loop Mellin–Barnes integrals in Minkowskian regions, *PoS LL2016* (2016) 034, <https://pos.sissa.it/260/034/pdf>, arXiv:1607.07538.
- [75] I. Dubovyk, A. Freitas, J. Gluza, T. Riemann, J. Usovitsch, 30 years, some 700 integrals, and 1 dessert, or: Electroweak two-loop corrections to the $Z\bar{b}b$ vertex, *PoS LL2016* (2016) 075, <https://pos.sissa.it/260/075/pdf>, arXiv:1610.07059.
- [76] I. Dubovyk, J. Gluza, T. Jelinski, T. Riemann, J. Usovitsch, New prospects for the numerical calculation of Mellin–Barnes integrals in Minkowskian kinematics, *Acta Phys. Pol. B* 48 (2017) 995, <https://doi.org/10.5506/APhysPolB.48.995>, arXiv:1704.02288.
- [77] T. Hahn, Generating Feynman diagrams and amplitudes with FeynArts 3, *Comput. Phys. Commun.* 140 (2001) 418–431, [https://doi.org/10.1016/S0010-4655\(01\)00290-9](https://doi.org/10.1016/S0010-4655(01)00290-9), arXiv:hep-ph/0012260.
- [78] S. Bauberger, F.A. Berends, M. Böhm, M. Buza, Analytical and numerical methods for massive two loop selfenergy diagrams, *Nucl. Phys. B* 434 (1995) 383–407, [https://doi.org/10.1016/0550-3213\(94\)00475-T](https://doi.org/10.1016/0550-3213(94)00475-T), arXiv:hep-ph/9409388.
- [79] S. Bauberger, M. Böhm, Simple one-dimensional integral representations for two loop selfenergies: the master diagram, *Nucl. Phys. B* 445 (1995) 25–48, [https://doi.org/10.1016/0550-3213\(95\)00199-3](https://doi.org/10.1016/0550-3213(95)00199-3), arXiv:hep-ph/9501201.
- [80] K. Hepp, Proof of the Bogolyubov–Parasiuk theorem on renormalization, *Commun. Math. Phys.* 2 (1966) 301–326, <https://doi.org/10.1007/BF01773358>.
- [81] T. Binoth, G. Heinrich, An automatized algorithm to compute infrared divergent multiloop integrals, *Nucl. Phys. B* 585 (2000) 741–759, [https://doi.org/10.1016/S0550-3213\(00\)00429-6](https://doi.org/10.1016/S0550-3213(00)00429-6), arXiv:hep-ph/0004013.
- [82] S. Borowka, G. Heinrich, S.P. Jones, M. Kerner, J. Schlenk, T. Zirke, SecDec 3.0: numerical evaluation of multi-scale integrals beyond one loop, *Comput. Phys. Commun.* 196 (2015) 470–491, <https://doi.org/10.1016/j.cpc.2015.05.022>, arXiv:1502.06595.
- [83] A.V. Smirnov, FIESTA 3: cluster-parallelizable multiloop numerical calculations in physical regions, *Comput. Phys. Commun.* 185 (2014) 2090–2100, <https://doi.org/10.1016/j.cpc.2014.03.015>, arXiv:1312.3186.
- [84] N.I. Usyukina, On a representation for three point function, *Teor. Mat. Fiz.* 22 (1975) 300–306, <https://doi.org/10.1007/BF01037795>.
- [85] V.A. Smirnov, Analytical result for dimensionally regularized massless on shell double box, *Phys. Lett. B* 460 (1999) 397–404, [https://doi.org/10.1016/S0370-2693\(99\)00777-7](https://doi.org/10.1016/S0370-2693(99)00777-7), arXiv:hep-ph/9905323.
- [86] J. Tausk, Nonplanar massless two loop Feynman diagrams with four on-shell legs, *Phys. Lett. B* 469 (1999) 225–234, [https://doi.org/10.1016/S0370-2693\(99\)01277-0](https://doi.org/10.1016/S0370-2693(99)01277-0), arXiv:hep-ph/9909506.
- [87] M. Czakon (MB MBAsymptotics), D. Kosower (barnesroutines), A. Smirnov, V. Smirnov (MBresolve), K. Bielas, I. Dubovyk, J. Gluza, K. Kajda, T. Riemann (AMBRE PlanarityTest), MBtools webpage <https://mbtools.hepforge.org/>.
- [88] M. Czakon, Automatized analytic continuation of Mellin–Barnes integrals, *Comput. Phys. Commun.* 175 (2006) 559–571, <https://doi.org/10.1016/j.cpc.2006.07.002>, arXiv:hep-ph/0511200.
- [89] A. Smirnov, V. Smirnov, On the resolution of singularities of multiple Mellin–Barnes integrals, *Eur. Phys. J. C* 62 (2009) 445–449, <https://doi.org/10.1140/epjc/s10052-009-1039-6>, arXiv:0901.0386.
- [90] J. Gluza, K. Kajda, T. Riemann, AMBRE – a Mathematica package for the construction of Mellin–Barnes representations for Feynman integrals, *Comput. Phys. Commun.* 177 (2007) 879–893, <https://doi.org/10.1016/j.cpc.2007.07.001>, arXiv:0704.2423.
- [91] K. Bielas, I. Dubovyk, J. Gluza, T. Riemann, Some remarks on non-planar Feynman diagrams, *Acta Phys. Pol. B* 44 (2013) 2249–2255, <https://doi.org/10.5506/APhysPolB.44.2249>, arXiv:1312.5603.
- [92] J. Gluza, K. Kajda, T. Riemann, V. Yundin, Numerical evaluation of tensor Feynman integrals in Euclidean kinematics, *Eur. Phys. J. C* 71 (2011) 1516, <https://doi.org/10.1140/epjc/s10052-010-1516-y>, arXiv:1010.1667.
- [93] I. Dubovyk, J. Gluza, T. Riemann, Non-planar Feynman diagrams and Mellin–Barnes representations with AMBRE 3.0, *J. Phys. Conf. Ser.* 608 (2015) 012070, <https://doi.org/10.1088/1742-6596/608/1/012070>, DESY 14–174 (13 April 2016).
- [94] M. Ochman, T. Riemann, MBsums – a Mathematica package for the representation of Mellin–Barnes integrals by multiple sums, *Acta Phys. Pol. B* 46 (2015) 2117, <https://doi.org/10.5506/APhysPolB.46.2117>, arXiv:1511.01323.
- [95] Silesian University at Katowice, webpage <http://prac.us.edu.pl/~gluza/ambre/>.

- [96] I. Dubovyk, T. Riemann, J. Usovitsch, Numerical calculation of multiple MB-integral representations for Feynman integrals, webpage <http://prac.us.edu.pl/~gluza/ambre/>;
J. Usovitsch, Mathematica/C++ package MBnumerics, 2018, in preparation.
- [97] T. Hahn, CUBA: a library for multidimensional numerical integration, *Comput. Phys. Commun.* 168 (2005) 78–95, <https://doi.org/10.1016/j.cpc.2005.01.010>, arXiv:hep-ph/0404043.
- [98] T. Hahn, Concurrent CUBA, *J. Phys. Conf. Ser.* 608 (2015) 012066, <https://doi.org/10.1088/1742-6596/608/1/012066>, arXiv:1408.6373.
- [99] A. Czarnecki, J.H. Kühn, Nonfactorizable QCD and electroweak corrections to the hadronic Z boson decay rate, *Phys. Rev. Lett.* 77 (1996) 3955–3958, <https://doi.org/10.1103/PhysRevLett.77.3955>, arXiv:hep-ph/9608366.
- [100] R. Harlander, T. Seidensticker, M. Steinhauser, Complete corrections of order $O(\alpha_s)$ to the decay of the Z boson into bottom quarks, *Phys. Lett. B* 426 (1998) 125–132, [https://doi.org/10.1016/S0370-2693\(98\)00220-2](https://doi.org/10.1016/S0370-2693(98)00220-2), arXiv:hep-ph/9712228.
- [101] J. Fleischer, O.V. Tarasov, F. Jegerlehner, P. Raczka, Two loop $O(\alpha_s G_\mu m_t^2)$ corrections to the partial decay width of the Z^0 into $b\bar{b}$ final states in the large top mass limit, *Phys. Lett. B* 293 (1992) 437–444, [https://doi.org/10.1016/0370-2693\(92\)90909-N](https://doi.org/10.1016/0370-2693(92)90909-N).
- [102] G. Buchalla, A.J. Buras, QCD corrections to the $\bar{s}dZ$ vertex for arbitrary top quark mass, *Nucl. Phys. B* 398 (1993) 285–300, [https://doi.org/10.1016/0550-3213\(93\)90110-B](https://doi.org/10.1016/0550-3213(93)90110-B).
- [103] G. Degrossi, Current algebra approach to heavy top effects in $Z \rightarrow b + \bar{b}$, *Nucl. Phys. B* 407 (1993) 271–289, [https://doi.org/10.1016/0550-3213\(93\)90058-W](https://doi.org/10.1016/0550-3213(93)90058-W), arXiv:hep-ph/9302288.
- [104] K. Chetyrkin, A. Kwiatkowski, M. Steinhauser, Leading top mass corrections of order $O(\alpha_s m_t^2 / M_W^2)$ to partial decay rate $\Gamma(Z \rightarrow b\bar{b})$, *Mod. Phys. Lett. A* 8 (1993) 2785–2792, <https://doi.org/10.1142/S0217732393003172>.
- [105] C. Patrignani, et al., Review of Particle Physics, *Chin. Phys. C* 40 (10) (2016) 100001, <https://doi.org/10.1088/1674-1137/40/10/100001>.
- [106] A. Freitas, Numerical multi-loop integrals and applications, *Prog. Part. Nucl. Phys.* 90 (2016) 201–240, <https://doi.org/10.1016/j.pnpnp.2016.06.004>, arXiv:1604.00406.
- [107] D. Bardin, P. Christova, M. Jack, L. Kalinovskaya, A. Olchevski, S. Riemann, T. Riemann, ZFITTER v. 6.21: a semianalytical program for fermion pair production in e^+e^- annihilation, *Comput. Phys. Commun.* 133 (2001) 229–395, [https://doi.org/10.1016/S0010-4655\(00\)00152-1](https://doi.org/10.1016/S0010-4655(00)00152-1), arXiv:hep-ph/9908433.
- [108] A. Arbuzov, M. Awramik, M. Czakon, A. Freitas, M. Grünewald, K. Mönig, S. Riemann, T. Riemann, ZFITTER: a semi-analytical program for fermion pair production in e^+e^- annihilation, from version 6.21 to version 6.42, *Comput. Phys. Commun.* 174 (2006) 728–758, <https://doi.org/10.1016/j.cpc.2005.12.009>, arXiv:hep-ph/0507146.
- [109] A. Akhundov, A. Arbuzov, S. Riemann, T. Riemann, The ZFITTER project, *Phys. Part. Nucl.* 45 (3) (2014) 529–549, <https://doi.org/10.1134/S1063779614030022>, arXiv:1302.1395.
- [110] D. Bardin, M. Grünewald, G. Passarino, Precision calculation project report, arXiv:hep-ph/9902452, 1999.
- [111] A. Leike, T. Riemann, J. Rose, S matrix approach to the Z line shape, *Phys. Lett. B* 273 (1991) 513–518, [https://doi.org/10.1016/0370-2693\(91\)90307-C](https://doi.org/10.1016/0370-2693(91)90307-C), arXiv:hep-ph/9508390.
- [112] T. Riemann, Cross-section asymmetries around the Z peak, *Phys. Lett. B* 293 (1992) 451–456, [https://doi.org/10.1016/0370-2693\(92\)90911-M](https://doi.org/10.1016/0370-2693(92)90911-M), arXiv:hep-ph/9506382.
- [113] T. Riemann, S-matrix Approach to the Z Resonance, *Acta Phys. Pol. B* 46 (11) (2015) 2235, <https://doi.org/10.5506/APhysPolB.46.2235>, arXiv:1610.04501.
- [114] T. Riemann, et al., Unfolding real observables at the Z peak, to appear as contribution to the report on [8], <https://indico.cern.ch/event/669224/contributions/2805424/attachments/1581663/2499614/2018-cern-ceex.pdf>, and, <https://indico.cern.ch/event/669224/contributions/2805418/attachments/1581648/2504620/riemann-FCCeeminiCERN-short.pdf>.